

# Solved exercises on advanced nonlinear differential equations

Ana Carpio, Universidad Complutense de Madrid

December, 2020

## 1 Contents

- Partial Differential Equations
  - Elliptic problems: 1-9
  - Hyperbolic problems and conservation laws: 10-11, 21-24
  - Parabolic problems: 9, 15-20
  - Navier-Stokes, vorticity equations: 12-14
  - Elasticity equations: 25-28
- Integrodifferential Equations: 1-5
- Numerical methods
  - Shooting: 1
  - Finite differences: 3-6
  - Variational techniques: 7-9
  - Spectral basis methods: 10
  - Particle methods: 11
  - Voronoi discretizations: 12
- Differential-Difference Equations
  - Front like solutions, pinned and traveling, depinning: 2, 3-8
  - Pulse like solutions, propagation failure, synchronization: 9-10
  - Two dimensional problems: 11-13
  - Problems with inertia: 1, 15-17
  - Blow-up: 18-19
  - Kinetic problems: 20-21

References

## 2 Partial Differential Equations

1. Consider the problem

$$\begin{cases} \nabla \cdot \gamma_e \nabla u = 0 & \text{in } \Omega \setminus \overline{\Omega}_i, & \nabla \cdot \gamma_i \nabla u = 0 & \text{in } \Omega_i, \\ u^- - u^+ = 0 & \text{on } \partial\Omega_i, & \gamma_i \nabla u^- \cdot \mathbf{n} - \gamma_e \nabla u^+ \cdot \mathbf{n} = 0 & \text{on } \partial\Omega_i, \\ \gamma_e \nabla u \cdot \mathbf{n} = j & \text{on } \partial\Omega. \end{cases}$$

with continuous and positive  $\gamma_e, \gamma_i$ , up to the boundary. We assume  $\Omega_i \subset \Omega$ , domains with smooth boundaries. The unit normal  $\mathbf{n}$  points outside  $\Omega_e$  but inside  $\Omega_i$  and  $u^-$  and  $u^+$  denote the limit values of  $u$  on  $\partial\Omega_i$  from outside and inside  $\Omega_i$ , respectively. Can we expect to have solutions for any  $j \in L^2(\partial\Omega)$ ? Can we expect uniqueness of solutions?

Taken from [57]. Integrating over  $\Omega$  and applying the divergence theorem, we find

$$\begin{aligned} & \int_{\Omega \setminus \overline{\Omega}_i} \nabla \cdot \gamma_e \nabla u \, d\mathbf{x} + \int_{\Omega_i} \nabla \cdot \gamma_i \nabla u \, d\mathbf{x} \\ &= \int_{\partial\Omega} \gamma_e \nabla u \cdot \mathbf{n} \, dl = \int_{\partial\Omega} j \, dl = 0. \end{aligned}$$

We have a constraint on the boundary integral of  $j$  to be able to construct solutions. Once this constant is satisfied, possible solutions are not unique, since addition of any constant provides another solution.

2. Calculate the solution of

$$\begin{aligned} \Delta p + \lambda^2 p &= a \delta_\Gamma \quad x \in \mathbb{R}^N, \\ \lim_{r \rightarrow \infty} r^{\frac{N-1}{2}} \left( \frac{\partial p}{\partial r} - i\lambda p \right) &= 0, \quad r = |\mathbf{x}|, \end{aligned}$$

where  $\delta_\Gamma$  is a Dirac mass supported at a curve  $\Gamma$ .

Taken from [63, 62]. The fundamental solution for the Helmholtz equation  $\Delta G + \lambda^2 G = -\delta$  in the whole space satisfying this condition at infinity (outgoing Sommerfeld radiation condition) is known in explicit form. The solution for this particular right hand side is obtained by convolution

$$\begin{aligned} p(\mathbf{x}) &= \int_{\mathbb{R}^N} G(\mathbf{x} - \mathbf{y}) a(\mathbf{y}) \delta_\Gamma(\mathbf{y}) \, d\mathbf{y} = \\ &= \int_\Gamma G(\mathbf{x} - \mathbf{y}) a(\mathbf{y}) \, d\mathbf{y}. \end{aligned}$$

3. Given a continuous function  $a$ , find an explicit expression for the solution of the problem

$$\begin{aligned} \Delta P(\mathbf{x}) + k_e^2 P(\mathbf{x}) &= a(\mathbf{x}) \delta_{\mathbf{x}_0} \quad \mathbf{x} \in \mathbb{R}^3, \\ \lim_{r \rightarrow 0} r \left( \frac{\partial P}{\partial r} + ik_e P \right) &= 0, \quad r = |\mathbf{x}|, \quad \mathbf{x}_0 \in \mathbb{R}^3. \end{aligned}$$

Taken from [69]. The complex conjugate  $Q = \overline{P}$  satisfies a Helmholtz equation with outgoing radiation condition at infinity:

$$\begin{aligned} \Delta Q(\mathbf{x}) + k_e^2 Q(\mathbf{x}) &= \overline{a(\mathbf{x})} \delta_{\mathbf{x}_0}, \quad \mathbf{x} \in \mathbb{R}^3, \\ \lim_{r \rightarrow 0} r \left( \frac{\partial Q}{\partial r} - ik_e Q \right) &= 0, \quad r = |\mathbf{x}|, \quad \mathbf{x}_0 \in \mathbb{R}^3. \end{aligned}$$

The fundamental solution is known to be  $F(\mathbf{x}) = \frac{e^{ik_e|\mathbf{x}|}}{4\pi|\mathbf{x}|}$ . Thus  $\overline{P} = -F * \overline{a} \delta_{\mathbf{x}_0}$ ,  $\delta_{x_0}$  being a Dirac mass supported at  $\mathbf{x}_0$  and

$$P(\mathbf{x}) = -\frac{e^{-ik_e|\mathbf{x}-\mathbf{x}_0|}}{4\pi|\mathbf{x}-\mathbf{x}_0|} a(\mathbf{x}_0).$$

4. Find an explicit expression for the solution of

$$\begin{aligned} \mathbf{curl}(\mathbf{curl} \mathbf{P}) - k_e^2 \mathbf{P} &= \mathbf{d}(\mathbf{x}) \delta_{\mathbf{x}_0} \quad \text{in } \mathbb{R}^3, \\ \lim_{|\mathbf{x}| \rightarrow \infty} |\mathbf{x}| |\mathbf{curl} \mathbf{P} \times \hat{\mathbf{x}} - ik_e \mathbf{P}| &= 0. \end{aligned}$$

Taken from [77]. We take the divergence of the equation. Since  $\text{div}(\mathbf{curl} \mathbf{A}) = 0$  for any vector  $\mathbf{A}$ , we find  $\text{div} \mathbf{P} = -\frac{1}{k_e^2} \text{div} \mathbf{d} \delta_{\mathbf{x}_0}$ . Making use of the vector identity  $\mathbf{curl}(\mathbf{curl} \mathbf{P}) = \nabla(\text{div} \mathbf{P}) - \Delta \mathbf{P}$  we have

$$-\Delta \mathbf{P} - k_e^2 \mathbf{P} = \delta_{\mathbf{x}_0} \mathbf{d} + \frac{1}{k_e^2} \nabla(\text{div} \mathbf{d} \delta_{\mathbf{x}_0}).$$

We can solve the equations by convolution with the Green function of Helmholtz equation:

$$\mathbf{P} = G_{k_e} * \mathbf{d} \delta_{\mathbf{x}_0} + \frac{1}{k_e^2} G_{k_e} * \nabla(\text{div} \mathbf{d} \delta_{\mathbf{x}_0}).$$

Notice that the right hand side can be rewritten as  $G_{k_e} * \mathbf{d} \delta_{\mathbf{x}_0} + \frac{1}{k_e^2} G_{k_e} * [\mathbf{curl} \mathbf{curl} \mathbf{d} \delta_{\mathbf{x}_0} + \Delta \mathbf{d} \delta_{\mathbf{x}_0}]$ . Interchanging the derivatives in the convolution we find

$$\mathbf{P}(\mathbf{x}) = \frac{1}{k_e^2} \mathbf{curl} \mathbf{curl} G_{k_e}(\mathbf{x} - \mathbf{x}_0) \mathbf{d}(\mathbf{x}_0).$$

for  $\mathbf{x} \neq \mathbf{x}_0$ .

5. Given a smooth semicircle  $\Omega$ , with curved upper boundary  $\partial\Omega^+$  and lower straight boundary  $\partial\Omega^-$ , consider the problem

$$\begin{aligned} d\Delta c &= k_s \frac{c}{c + K_s}, \quad \mathbf{x} \in \Omega \\ c &= c_0 > 0, \quad \mathbf{x} \in \partial\Omega^- \\ \frac{\partial c}{\partial \mathbf{n}} &= 0, \quad \mathbf{x} \in \partial\Omega^+, \end{aligned}$$

with positive parameters  $d, k_s, K_s$ . Prove that this problem has a nonnegative solution  $c \in H^1(\Omega)$ .

Taken from [60]. The solution  $c$  can be constructed as the limit of iterates  $c^{(m)}$  solution of linearized problems

$$\begin{aligned} d\Delta c^{(m)} &= \frac{k_s}{c^{(m-1)} + K_s} c^{(m)}, & \mathbf{x} \in \Omega \\ c^{(m)} &= c_0 > 0, & \mathbf{x} \in \partial\Omega^- \\ \frac{\partial c^{(m)}}{\partial \mathbf{n}} &= 0, & \mathbf{x} \in \partial\Omega^+, \end{aligned}$$

starting from  $c^{(0)} = c_0$ . Lax Milgram's Theorem implies existence of a unique solution  $c^{(m)} \in H^1(\Omega)$ . Set  $a_{m-1} = \frac{k_s}{c^{(m-1)} + K_s}$ . We multiply the equation by the negative part of  $c^{(m)}$ ,  $c^{(m)-}$

$$d \int_{\Omega} |\nabla c^{(m)-}|^2 d\mathbf{x} + \int_{\Omega} a_{m-1} |c^{(m)-}|^2 d\mathbf{x} = 0,$$

because  $\int_{\Omega} \frac{\partial c^{(m)}}{\partial \mathbf{n}} c_0^- dl = 0$ . Initially,  $a_0 > 0$ . Thus,  $c^{(1)-} = 0$  and  $c^{(1)} \geq 0$ , which implies  $a_1$ . By induction, we conclude that  $c^{(m)} \geq 0$ ,  $a_m \geq 0$  and  $a_m \leq k_s/K_s$ . Writing  $c^{(m)} = \tilde{c}^{(m)} + c_0$ , with  $\tilde{c}^{(m)} \in H_0^1(\Omega)$ , we get

$$\begin{aligned} d\Delta \tilde{c}^{(m)} &= a_{m-1} \tilde{c}^{(m)} + a_{m-1} c_0, & \mathbf{x} \in \Omega \\ \tilde{c}^{(m)} &= 0 > 0, & \mathbf{x} \in \partial\Omega^- \\ \frac{\partial \tilde{c}^{(m)}}{\partial \mathbf{n}} &= 0, & \mathbf{x} \in \partial\Omega^+. \end{aligned}$$

Multiplying by  $\tilde{c}^{(m)}$  and integrating, we find

$$d \int_{\Omega} |\nabla \tilde{c}^{(m)}|^2 d\mathbf{x} + \int_{\Omega} a_{m-1} |\tilde{c}^{(m)}|^2 d\mathbf{x} = \int_{\Omega} a_{m-1} c_0 \tilde{c}^{(m)} d\mathbf{x}.$$

Using Poincaré's inequality,  $\|\tilde{c}^{(m)}\|_{H_0^1(\Omega)} \leq C(\Omega) \frac{k_s c_0}{K_s}$ . By Sobolev injections, we can extract a sequence converging weakly in  $H_0^1$ , strongly in  $L^2$  and pointwise to a limit  $\tilde{c}$ . Passing to the limit in the equation,  $c = \tilde{c} + c_0 \geq 0$  is a solution to the original problem.

6. Prove that the solution  $\Phi$  of the equation

$$-\frac{d^2}{dx^2} \Phi(x) = n_D(x) - \int_{\mathbb{R}} \frac{dk}{1 + \exp(\epsilon(k) - \Phi(x))}$$

with  $\int_{\mathbb{R}^2} \frac{dk dx}{1 + \exp(\epsilon(k) - \Phi(x))} = a$  fixed and  $\frac{d\Phi}{dx} \in L^2$  is unique.

Taken from [21]. Assume that there are two solutions  $\Phi_1$  and  $\Phi_2$  satisfying such conditions. Set  $U = \Phi_1 - \Phi_2$ . Then,  $\frac{dU}{dx} \in L^2$  and

$$\frac{d^2 U}{dx^2} = \int_{\mathbb{R}} \frac{dk}{1 + \exp(\epsilon(k) - \Phi_1(x))} - \int_{\mathbb{R}} \frac{dk}{1 + \exp(\epsilon(k) - \Phi_2(x))}.$$

Let us assume first that  $U(x) > 0$  everywhere. Then

$$a = \int_{\mathbb{R}^2} \frac{dkdx}{1 + \exp(\epsilon(k) - \Phi_1(x))} > \int_{\mathbb{R}^2} \frac{dkdx}{1 + \exp(\epsilon(k) - \Phi_2(x))} = a,$$

which is impossible.

Let us assume now that there is a unique point  $x_0$  at which  $U(x_0) = 0$ . We take  $U(x) < 0$  for  $x < x_0$  and  $U(x) > 0$  for  $x > x_0$ . Thus,  $\frac{d^2U}{dx^2} < 0$  if  $x < x_0$  and  $\frac{d^2U}{dx^2} < 0$  if  $x > x_0$ . Then,  $\frac{dU}{dx}$  is decreasing if  $x < x_0$  and  $\frac{dU}{dx}$  is increasing if  $x > x_0$ . On the other hand,

$$\int_{\mathbb{R}} \left( \frac{dU}{dx} \right)^2 dx = \int_{-\infty}^{x^*} \left( \frac{dU}{dx} \right)^2 dx + \int_{x^*}^{\infty} \left( \frac{dU}{dx} \right)^2 dx$$

is finite. If there exists  $x^*$  such that  $\frac{dU(x^*)}{dx} > 0$  and  $x^* < x_0$  then  $\int_{-\infty}^{x^*} \left( \frac{dU}{dx} \right)^2 dx > \left( \frac{dU(x^*)}{dx} \right)^2 \int_{-\infty}^{x^*} dx = \infty$ . This is impossible, so that  $\frac{dU}{dx} \leq 0$  for all  $x$  and  $U$  is decreasing. This contradicts our assumption on  $x_0$ . Therefore, we should have at least two points  $x_0$  and  $x_1$  at which  $U$  vanishes.

Let  $x_0$  and  $x_1$  be such that  $U(x_0) = U(x_1) = 0$ . If  $x_M$  is such that  $U(x_M) = \max \{U(x), x_0 \leq x \leq x_1\} > 0$ , then  $\frac{d^2U(x_M)}{dx^2} \leq 0$  because the maximum is attained at an interior point. However,

$$0 \geq \frac{d^2U(x_M)}{dx^2} = \int_{\mathbb{R}} \frac{dk}{1 + \exp(\epsilon(k) - \Phi_1(x_M))} - \int_{\mathbb{R}} \frac{dk}{1 + \exp(\epsilon(k) - \Phi_2(x_M))} > 0,$$

since  $U(x_M) > 0$ . Hence,  $\max \{U(x), x_0 \leq x \leq x_1\} = 0$ . In an analogous way, we conclude that  $U(x_m) = \min \{U(x), x_0 \leq x \leq x_1\} = 0$ . Therefore,  $U = 0$  on  $[x_0, x_1]$ .

Now we set  $x_0 = \min \{x | U(x) = 0\}$  and  $x_1 = \max \{x | U(x) = 0\}$ . Then,  $U(x) < 0$  for  $x < x_0$  and  $U(x) > 0$  for  $x > x_1$ . Repeating the above arguments, we would obtain  $x' \notin [x_0, x_1]$  such that  $U(x') = 0$ . This contradicts the definition of  $x_0$  and  $x_1$ . Therefore,  $U = 0$  everywhere and  $\Phi_1 = \Phi_2$ .

7. Consider balls  $B_\varepsilon = B(\mathbf{x}, \varepsilon)$  centered at a point  $\mathbf{x}$  of small radius  $\varepsilon$ . Given a smooth function  $u(\mathbf{x})$ , let  $v_\varepsilon$  be the solution of

$$\begin{cases} \Delta v_\varepsilon + k^2 v_\varepsilon = 0, & \text{in } \mathbb{R}^2 \setminus \overline{B_\varepsilon}, \\ v_\varepsilon = -u(\mathbf{x}), & \text{on } \partial B_\varepsilon, \\ \lim_{r \rightarrow \infty} r^{1/2} \left( \frac{\partial v_\varepsilon}{\partial r} - ikv_\varepsilon \right) = 0. \end{cases}$$

What is the behavior of  $\frac{\partial v_\varepsilon}{\partial \mathbf{n}}$  as  $\varepsilon \rightarrow 0$ ?

Taken from [47]. The Dirichlet–to–Neumann provides an expression for the normal derivative of  $v_\varepsilon$  on  $\Gamma_\varepsilon$ :

$$\begin{aligned} & \partial_{\mathbf{n}} v_\varepsilon(\mathbf{x} + \varepsilon(\cos \theta, \sin \theta)) \\ &= \frac{k}{2\pi} \sum_{n=-\infty}^{\infty} \frac{(H_{|n|}^{(1)})'(k\varepsilon)}{H_{|n|}^{(1)}(k\varepsilon)} \int_0^{2\pi} e^{in(\theta-\Theta)} u(\mathbf{x} + \varepsilon(\cos \Theta, \sin \Theta)) d\Theta \end{aligned}$$

in polar coordinates. Here  $H_{|n|}^{(1)}$  denotes the Hankel function of the first kind of order  $|n|$ . We choose the normal vector  $\mathbf{n}$  pointing into  $B_\varepsilon$ . For sufficiently small  $\varepsilon > 0$ ,

$$\frac{\partial v_\varepsilon}{\partial \mathbf{n}}(\mathbf{x} + \varepsilon(\cos \theta, \sin \theta)) = k \frac{(H_0^{(1)})'(k\varepsilon)}{H_0^{(1)}(k\varepsilon)} u(\mathbf{x}) + O(\varepsilon).$$

For small  $\varepsilon > 0$ , the Hankel functions have the following leading parts:

$$H_0^{(1)}(k\varepsilon) \sim \frac{-2 \log(k\varepsilon)}{\pi i}, \quad (H_0^{(1)})'(k\varepsilon) = -H_1^{(1)}(k\varepsilon) \sim \frac{-2}{\pi i k\varepsilon}.$$

Thus,

$$\frac{(H_0^{(1)})'(k\varepsilon)}{H_0^{(1)}(k\varepsilon)} \sim \frac{1}{k\varepsilon \log(k\varepsilon)},$$

and  $\frac{\partial v_\varepsilon}{\partial \mathbf{n}}(\mathbf{x} + \varepsilon(\cos \theta, \sin \theta)) \sim \frac{1}{\varepsilon \log(k\varepsilon)} u(\mathbf{x})$ .

8. Given a bounded open set  $\Omega \subset \mathbb{R}^N$ , we consider the problem: Find  $u > 0$  such that

$$\begin{aligned} -\Delta u &= u^p & \mathbf{x} \in \Omega, \\ u &= 0 & \mathbf{x} \in \partial\Omega, \\ u &> 0 & \mathbf{x} \in \Omega. \end{aligned}$$

Prove that there is a solution when  $1 < p+1 < p^*$ , where  $p^* = \infty$  if  $N \leq 2$  and  $p^* < \frac{2N}{N-2}$  when  $N > 2$ .

Consider the minimization problem

$$I = \text{Min}_{u \in H_0^1(\Omega)} \frac{\int_\Omega |\nabla u|^2 d\mathbf{x}}{\int_\Omega |u|^{p+1} d\mathbf{x}} = \text{Min}_{u \in H_0^1(\Omega)} J(u).$$

The functional  $J(u)$  to be minimized is positive, thus, bounded from below. Consider a minimizing sequence  $u_n \in H_0^1(\Omega)$ , such that  $J(u_n) \rightarrow I$  as  $n \rightarrow \infty$ . The sequence  $v_n = \frac{u_n}{\|u_n\|_{L^{p+1}}}$  is a minimizing sequence satisfying also  $\|v_n\|_{L^{p+1}} = 1$ . Then,  $\int_\Omega |\nabla v_n|^2 d\mathbf{x} \rightarrow I$  implies that  $v_n$  is bounded in  $H_0^1(\Omega)$  and  $v_n$  tends weakly in  $H_0^1$  to a limit  $v \in H_0^1(\Omega)$ . By Sobolev injections,  $v_n$  is compact in  $L^{p+1}$ ,  $p+1 < p^*$ , thus  $v \in L^{p+1}(\Omega)$

and  $\|v_n\|_{L^{p+1}} = 1 \rightarrow \|v\|_{L^{p+1}} = 1$ . By lower semicontinuity of weak convergence, we have  $J(v) \leq \lim_{n \rightarrow \infty} J(v_n) = I$ . Since  $v \in H_0^1(\Omega)$ , we have  $I \leq J(v)$ . Therefore,  $I = J(v)$  and the minimum is attained at  $v$ . Moreover, we can replace  $v$  by  $|v|$  and  $J(|v|) \leq I(v)$ , so that  $w = |v| \geq 0$  is a minimizer too and  $I = J(w)$ .  $w \neq 0$  because  $\|w\|_{L^{p+1}} = 1$ .

Now,  $J(w) \leq J(w + tr)$ ,  $r \in H_0^1(\Omega)$  for real  $t$ . An asymptotic expansion first for  $t > 0$  then for  $t < 0$  leads to

$$\int_{\Omega} \nabla w \nabla r \, d\mathbf{x} = c \int_{\Omega} w^p r \, d\mathbf{x}$$

for all  $r \in H_0^1(\Omega)$  and some  $c > 0$ . This implies  $-\Delta w = c w^p$ . Setting  $u = c^{-1/(p-1)} w$ , we get  $-\Delta u = u^p$  and  $u \geq 0$ ,  $u \neq 0$ . By the strong maximum principle,  $u > 0$ .

If  $p + 1 = p^* = \frac{2N}{N-2}$  and  $N > 2$  existence depends on the geometry of  $\Omega$ , see [1].

9. Prove that the function  $v(\mathbf{x}, t) = |t|^{\frac{p}{p-1}} \phi(\mathbf{x})$ ,  $1 < p < p^* - 1$ , where

$$\begin{aligned} -\Delta \phi &= \left( \frac{p}{p-1} \right)^p |\phi|^{p-1} \phi & \mathbf{x} \in \Omega, \\ \phi &= 0 & \mathbf{x} \in \partial\Omega, \end{aligned}$$

is a solution of the backward parabolic problem

$$\begin{aligned} -\Delta v + |v_t|^{p-1} v_t &= 0 & \mathbf{x} \in \Omega \times (-\infty, 0], \\ v &= 0 & \mathbf{x} \in \partial\Omega \times (-\infty, 0]. \end{aligned}$$

Proof taken from [3, 8]. We see that

$$\begin{aligned} v_t &= -\frac{p}{p-1} |t|^{\frac{1}{p-1}} \phi(\mathbf{x}), \\ |v_t|^{p-1} v_t &= -\left( \frac{p}{p-1} \right)^p |t|^{\frac{p}{p-1}} |\phi(\mathbf{x})|^{p-1} \phi(\mathbf{x}), \\ -\Delta v &= -|t|^{\frac{p}{p-1}} \Delta \phi(\mathbf{x}) = |t|^{\frac{p}{p-1}} \left( \frac{p}{p-1} \right)^p |\phi(\mathbf{x})|^{p-1} \phi(\mathbf{x}), \end{aligned}$$

so that the equation is fulfilled. Existence of  $\phi$  follows from critical point theory.

10. Consider a membrane whose vertical deviation from a flat equilibrium is governed by

$$\rho \frac{\partial^2 w}{\partial t^2} = d\Delta w - \kappa \Delta^2 w + f(x, y, t).$$

where  $\rho, d, \kappa$  are positive constants. Would you expect this system to develop oscillatory patterns with definite wave lengths?

Taken from [58]. The elliptic wave-plate operator with zero Dirichlet boundary conditions in a rectangular admits a sequence of positive eigenvalues  $\lambda_{m,n}$  with eigenfunctions  $\phi_{m,n}$  given by combinations of sinus and cosinus functions whose period is related to the spatial domain and varies with the eigenvalue. Seeking a series solution by separation of variables, we see that the problem admits solutions of the form

$$\sum_{n,m} a_{n,m}(t)\phi_{n,m}(x,y),$$

where  $a_{n,m}(t)$  is solution of

$$a''_{n,m} + \lambda_{n,m}a_{n,m} = f_{n,m},$$

therefore, a combination of  $\sin(\sqrt{\lambda_{n,m}}t)$  and  $\cos(\sqrt{\lambda_{n,m}}t)$ , after expressing  $f(x,y,t) = \sum_{n,m} f_{n,m}(t)\phi_{n,m}(x,y)$  as a series of eigenfunctions. More complex models in which  $w$  is coupled to Navier equations for in-plane motion  $(u,v)$  and  $f$  is given by either spins or functional expressions informed by them are used to explain ripple formation in graphene [59, 58, 54].

11. Given a solution  $u \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^+, H_0^1(\Omega)) \cap W_{\text{loc}}^{2,\infty}(\mathbb{R}^+, L^2(\Omega))$  of

$$u_{tt} - \Delta u + \alpha|u_t|^{p-1}u_t = 0 \quad \text{in } L^\infty(\mathbb{R}^+, H^{-1}(\Omega))$$

with  $\alpha > 0, 1 < p$  and  $p+1 < p^*$ , we set

$$E(t) = \frac{1}{2} \int_{\Omega} |\nabla u(\mathbf{x}, t)|^2 d\mathbf{x} + \frac{1}{2} \int_{\Omega} |u_t(\mathbf{x}, t)|^2 d\mathbf{x}.$$

Then, for some positive constant  $C(E(0))$ , we have

$$E(t) \leq C(E(0))t^{-2/(p-1)}, \quad t > 0.$$

Proof taken from [2]. We set  $\phi(t) = E^{(p-1)/2} \int_{\Omega} uu_t d\mathbf{x}$ . Next, we differentiate with respect to  $t$  to get

$$\begin{aligned} E'(t) &= -\alpha \int_{\Omega} |u_t|^{p+1} d\mathbf{x} \leq 0, \\ \phi'(t) &= E(t)^{(p-1)/2} \left( \int_{\Omega} |u_t|^2 d\mathbf{x} - \int_{\Omega} |\nabla u|^2 d\mathbf{x} - \alpha \int_{\Omega} |u_t|^{p-1} u_t u d\mathbf{x} \right) \\ &\quad + \frac{p-1}{2} E(t)^{(p-3)/2} E'(t) \int_{\Omega} uu_t d\mathbf{x} \end{aligned}$$

First, notice that  $E(t) \leq E(0)$  and  $-\int_{\Omega} |\nabla u|^2 d\mathbf{x} = -2E(t) + \int_{\Omega} |u_t|^2 d\mathbf{x}$ . Moreover,

$$E(t)^{-1} \left| \int_{\Omega} uu_t d\mathbf{x} \right| \leq E(t)^{-1} \left( \frac{1}{2} \int_{\Omega} |u|^2 d\mathbf{x} + \frac{1}{2} \int_{\Omega} |u_t|^2 d\mathbf{x} \right) \leq C(\Omega)$$



for some positive constant  $C(\Omega)$  because Poincaré's inequality implies  $\frac{1}{2} \int_{\Omega} |u|^2 d\mathbf{x} \leq \frac{\lambda(\Omega)}{2} \int_{\Omega} |\nabla u|^2 d\mathbf{x}$ . As a consequence, we get

$$\begin{aligned} \phi'(t) &\leq 2E(t)^{(p-1)/2} \int_{\Omega} |u_t|^2 d\mathbf{x} - \alpha E(t)^{(p-1)/2} \int_{\Omega} |u_t|^{p-1} u_t u d\mathbf{x} \\ &\quad - 2E(t)^{(p+1)/2} - \frac{p-1}{2} C(\Omega) E(0)^{(p-1)/2} E'(t). \end{aligned}$$

Now we set  $\psi_{\varepsilon}(t) = (1 + K_1 \varepsilon)E(t) + \varepsilon \phi(t)$  with  $K_1 = \frac{p-1}{2} C(\Omega) E(0)^{(p-1)/2}$ . We get

$$\begin{aligned} \psi'_{\varepsilon}(t) &\leq 2\varepsilon E(t)^{(p-1)/2} \int_{\Omega} |u_t|^2 d\mathbf{x} - \alpha \varepsilon E(t)^{(p-1)/2} \int_{\Omega} |u_t|^{p+1} d\mathbf{x} \\ &\quad - 2\varepsilon E(t)^{(p+1)/2} - \alpha \int_{\Omega} |u_t|^{p+1} d\mathbf{x} \end{aligned}$$

Notice that  $\|u_t\|_{L^2}^2 \leq \text{meas}(\Omega)^{(p-1)/(p+1)} (\int_{\Omega} |u_t|^{p+1})^{2/(p+1)}$ . By Young's inequality

$$\begin{aligned} 2\varepsilon E(t)^{\frac{(p-1)}{2}} \int_{\Omega} |u_t|^2 d\mathbf{x} &\leq 2\varepsilon \text{meas}(\Omega)^{\frac{p-1}{p+1}} E(t)^{\frac{p-1}{2}} \left( \int_{\Omega} |u_t|^{p+1} \right)^{\frac{2}{p+1}} \\ &\leq \varepsilon E(t)^{\frac{p+1}{2}} + \varepsilon \delta \int_{\Omega} |u_t|^{p+1} \end{aligned}$$

for some positive  $\delta$  depending on  $\Omega$ .

Using Sobolev injections for  $p+1 < p^*$  we find

$$\int_{\Omega} |u_t|^{p-1} u_t u d\mathbf{x} \leq \left( \int_{\Omega} |u_t|^{p+1} d\mathbf{x} \right)^{\frac{p}{p+1}} \|u\|_{L^{p+1}} \leq S(\Omega) \|u_t\|_{L^{p+1}}^p \|\nabla u\|_{L^2}.$$

Notice that  $\|\nabla u\|_{L^2} \leq 2E(t)$ . By Young's inequality again

$$\begin{aligned} \varepsilon \alpha E(t)^{(p-1)/2} \int_{\Omega} |u_t|^{p-1} u_t u d\mathbf{x} &\leq \varepsilon \alpha E(t)^{(p-1)/2} S(\Omega) \|u_t\|_{L^{p+1}}^p \|\nabla u\|_{L^2} \\ &\leq \frac{\alpha}{2} \int_{\Omega} |u_t|^{p+1} + \varepsilon \eta(\varepsilon) E(t)^{(p+1)/2} \end{aligned}$$

where  $\eta > 0$  depends on  $E(0)$ ,  $\Omega$ ,  $\alpha$  and  $\varepsilon$ , and tends to zero as  $\varepsilon$  tends to zero. Adding up, we get

$$\psi'_{\varepsilon}(t) \leq \left(-\frac{\alpha}{2} + \varepsilon \delta\right) \int_{\Omega} |u_t|^{p+1} + \varepsilon(-1 + \eta(\varepsilon)) E(t)^{(p+1)/2}.$$

On the other hand, for  $\varepsilon$  small enough,

$$\frac{1}{\varepsilon} E(t) \leq (1 - K_2 \varepsilon) E(t) \leq \psi_{\varepsilon}(t) \leq (1 + K_2 \varepsilon) \leq 2E(t).$$

Choosing  $\varepsilon$  small enough, we find

$$\psi'_\varepsilon(t) \leq -\frac{\varepsilon}{4}E^{(p+1)/2} \leq -\frac{\varepsilon K_3}{4}\psi_\varepsilon(t)^{(p+1)/2}.$$

Integrating the inequality we find  $E(t) \leq C(E(0))t^{-2/(p-1)}$  for  $t > 0$ .

12. Consider the vorticity equation in two dimensions. Let  $v = \text{curl } \mathbf{u} \in C((0, \infty); W^{1,p}(\mathbb{R}^2))$ ,  $1 \leq p \leq \infty$ , be the solution of

$$\begin{aligned} v_t - \Delta v + \mathbf{u} \cdot \nabla v &= 0, & \mathbf{x} \in \mathbb{R}^2 \times \mathbb{R}^+ \\ v(\mathbf{x}, 0) &= v_0, & \mathbf{x} \in \mathbb{R}^2, \end{aligned}$$

for a divergence free velocity field  $\mathbf{u}$  and an initial datum  $v_0 \in L^1(\mathbb{R}^2)$ .

Prove 1) that the mass  $\int_{\mathbb{R}^2} v_0 \, d\mathbf{x}$  does not change with time and 2) that  $\|v(t)\|_{L^p(\mathbb{R}^2)} \leq Ct^{-1+\frac{1}{p}}$  for  $t > 0$ .

Proof taken from [4, 5]. Notice that  $\mathbf{u} \cdot \nabla v = \text{div}(\mathbf{u}v) = 0$ . Integrating the equation, using the divergence theorem, and the fact that  $v$  vanishes at infinity we get

$$\frac{d}{dt} \int_{\mathbb{R}^2} v_0 \, d\mathbf{x} = 0.$$

The velocity vector is given by

$$\mathbf{u}(\mathbf{x}, t) = K * v(\mathbf{x}, t) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{(-y_2, y_1)}{|\mathbf{y}|^2} v(\mathbf{x} - \mathbf{y}, t) d\mathbf{y}$$

where the kernel  $K \in L^{2,\infty}$  and  $\|K * v\|_{L^r} \leq \|K\|_{L^{2,\infty}} \|v\|_{L^p}$  for  $r > 2$ ,  $1 < p < 2$ ,  $1/r = 1/p - 1/2$ .

Writing down the integral expression for the solution

$$v(t) = G(t) * v_0 + \int_0^t \nabla G(t-s) * [v(s) \mathbf{K} * v(s)] ds,$$

where  $G(t)$  stands for the heat kernel, and taking norms we find

$$\|v(t)\|_{L^p} = \|G(t) * v_0\|_{L^p} + \int_0^t \|\nabla G(t-s) * [v(s) \mathbf{K} * v(s)]\|_{L^p} ds.$$

The integral terms decays faster than the rest, therefore

$$\|v(t)\|_{L^p} \sim \|G(t) * v_0\|_{L^p} \leq Ct^{-1+\frac{1}{p}}.$$

Recall that  $G(t) * v_0$  is a solution of the heat equation with datum  $v_0$  and it belongs to  $L^p$  for all  $1 \leq p \leq \infty$  for any  $t > 0$  if  $v_0 \in L^1$ . Moreover,  $\|G(t) * v_0\|_{L^p} \leq \|G(t)\|_{L^p} \|v_0\|_{L^1}$  and  $\|G(t)\|_{L^p} = Ct^{-1+\frac{1}{p}}$ .

13. Let  $\mathbf{u}$  be a solution of the incompressible Navier-Stokes equations in two dimensions with initial datum  $\mathbf{u}_0 \in L^1 \cap L^2(\mathbb{R}^2)$  such that  $\operatorname{div}(\mathbf{u}_0) = 0$ . Then  $\mathbf{u}(t) \in L^p(\mathbb{R}^2)$  for  $1 \leq p \leq 2$  and  $t > 0$ .

Proof taken from [6, 10]. The theory of classical solutions with  $L^2$  data, that is,  $\mathbf{u}_0 \in L^2(\mathbb{R}^2)$  guarantees that  $\mathbf{u}(t) \in L^\infty([0, \infty); L^2(\mathbb{R}^2))$  and is bounded by  $\|\mathbf{u}_0\|_{L^2}$ . By taking the divergence of Navier-Stokes equations

$$\mathbf{u}_t - \Delta \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = \nabla p, \quad \operatorname{div}(\mathbf{u}) = 0,$$

we get an equation for the pressure

$$-\Delta p = \operatorname{div}(\mathbf{u} \cdot \nabla \mathbf{u}).$$

The pressure is then the convolution  $p = E_2 * \operatorname{div}(\mathbf{u} \cdot \nabla \mathbf{u})$ , where  $E_2$  is the fundamental solution of  $-\Delta$  in  $\mathbb{R}^2$ , up to a function of time. Then  $\mathbf{u}$  satisfies the integral equation

$$\begin{aligned} \mathbf{u}(t) &= G(t) * \mathbf{u}_0 + \int_0^t \partial_i G(t-s) * u_i \mathbf{u}(s) ds \\ &\quad + \int_0^t \partial_i G(t-s) * \partial_j \nabla E_2 * u_i u_j(s) ds, \end{aligned}$$

where  $\partial_i$  denotes partial derivative with respect to  $x_i$ ,  $u_i$  are components of  $\mathbf{u}$  and summation with respect to repeated indices is intended. Since  $u \in L^1$ ,  $G(t) * u_0 \in L^q$  for all  $q > 1$  and  $t > 0$ . On the other hand,  $u(s) \in L^2$  implies that  $u_i u_j(s) \in L^1$ . Moreover,

$$\left\| \int_0^t \partial_i G(t-s) * u_i u_j(s) ds \right\|_{L^q} \leq C \int_0^t (t-s)^{-1+\frac{1}{q}-\frac{1}{2}} \|\mathbf{u}\|_{L^2}^2 ds \leq C t^{\frac{1}{q}-\frac{1}{2}}$$

for  $1 \leq q < 2$ . Thus, the first integral belongs to  $L^q$  for  $1 \leq q < 2$ . Let us consider now the second integral. Since  $\partial_i G(t)$  belongs to the Hardy space  $\mathcal{H}^1(\mathbb{R}^2)$  and  $\partial_j \nabla E_2$  is a Calderon-Zygmund kernel, we conclude that  $\partial_i G(t-s) * \partial_j \nabla E_2 \in L^1$  and

$$\|\partial_i G(t-s) * \partial_j \nabla E_2\|_{L^1} \leq C \|\partial_i G(t-s)\|_{\mathcal{H}^1} < C(t-s)^{-\frac{1}{2}}.$$

Thus,

$$\left\| \int_0^t \partial_i G(t-s) * \partial_j \nabla E_2 * u_i u_j(s) ds \right\|_{L^1} \leq \int_0^t C(t-s)^{-\frac{1}{2}} \|\mathbf{u}(s)\|_{L^2}^2 ds \leq C t^{\frac{1}{2}}.$$

In an analogous way, since  $\partial_j \nabla E_2$  is a Calderon-Zygmund kernel, we conclude that  $\partial_i G(t-s) * \partial_j \nabla E_2 \in L^q$ ,  $1 < q < \infty$  and

$$\|\partial_i G(t-s) * \partial_j \nabla E_2\|_{L^q} \leq C \|\partial_i G(t-s)\|_{L^q} < C(t-s)^{-1+\frac{1}{q}-\frac{1}{2}}.$$

Thus,

$$\begin{aligned} \left\| \int_0^t \partial_i G(t-s) * \partial_j \nabla E_2 * u_i u_j(s) ds \right\|_{L^q} &\leq \int_0^t C(t-s)^{-1+\frac{1}{q}-\frac{1}{2}} \|\mathbf{u}(s)\|_{L^2}^2 ds \\ &\leq Ct^{\frac{1}{q}-\frac{1}{2}} \end{aligned}$$

for  $1 < q \leq 2$ .

14. A line vortex lying along a curve  $\Gamma$  in an incompressible inviscid and irrotational fluid is a solution of the following equations

$$\operatorname{div}(\mathbf{u}) = 0, \quad \operatorname{curl}(\mathbf{u}) = \omega_0 \delta_\Gamma(\mathbf{x}),$$

where  $\mathbf{u}$  is the fluid velocity,  $\omega_0 = 2\pi\gamma$  is the circulation around the vortex and  $\gamma$  is the vortex strength.  $\delta_\Gamma$  is a Dirac function supported at the curve  $\Gamma$ . Express this solution in terms of a vector stream function.

Taken from [11]. We define a vector stream function  $\mathbf{U}$  in  $\mathbb{R}^3$  as the solution of  $\operatorname{div}(\mathbf{U}) = 0$ ,  $\operatorname{curl}(\mathbf{U}) = \mathbf{u}$ . Then  $-\Delta \mathbf{U} = \omega_0 \delta_\Gamma(\mathbf{x})$ . Using the Green function for the Laplacian in  $\mathbb{R}^3$  we get  $\mathbf{U} = \frac{\omega_0}{4\pi} \int_\Gamma \frac{1}{|\mathbf{x}-\mathbf{x}'|} d\mathbf{x}'$ .

15. Set  $v^+(x, t) = u(x, t) + q^+(x, t)$  in  $(x_i, x_{i+1})$  where  $u$  is a solution of

$$\begin{aligned} \frac{\partial u}{\partial t} - D_c \frac{\partial^2 u}{\partial x^2} + \frac{u}{R} &= f^+, \quad x \in (x_i, x_{i+1}) = (i, i+1), \quad t > 0 \\ u(x_i, t) &= 0, \quad u(x_{i+1}, t) = 0, \\ u(x, 0) &= h^+(x, 0), \end{aligned}$$

with

$$\begin{aligned} q^+(x, t) &= v_i(t) \frac{x - x_{i+1}}{x_i - x_{i+1}} + v_{i+1}(t) \frac{x - x_i}{x_{i+1} - x_i}, \\ f^+(x, t) &= \frac{q^+(x, t)}{R} - \frac{\partial q^+}{\partial t}(x, t), \\ h^+(x, 0) &= v(x, 0) - q^+(x, 0). \end{aligned}$$

Obtain an explicit expression for  $v$ .

Taken from [53]. Let  $\lambda_i = D_c(i\pi)^2 + \frac{1}{R}$  and  $\phi_i(x) = \sin(\sqrt{\lambda_i}x) \left( \int_0^1 \sin(\sqrt{\lambda_i}x)^2 dx \right)^{-1}$  be the eigenvalues and orthonormalized eigenfunctions of the operator  $-D_c \frac{\partial^2 u}{\partial x^2} + \frac{u}{R} = 0$  in  $(0, 1)$  with zero boundary conditions. We expand  $f^+$  and  $h^+$  as a Fourier series of the eigenfunctions

$$\begin{aligned} f^+(x, t) &= \sum_{i=0}^{\infty} f_i^+(t) \phi_i(x), \quad f_i^+(t) = \int_0^1 f^+(z + x_i, t) \phi_i(z) dz, \\ h^+(x, 0) &= \sum_{i=0}^{\infty} h_i^+ \phi_i(x), \quad h_i^+ = \int_0^1 h^+(z + x_i, 0) \phi_i(z) dz. \end{aligned}$$

The explicit expression we seek is then given by

$$v^+(x, t) = q^+(x, t) + \sum_{i=0}^{\infty} e^{-\lambda_i t} h_i^+(t) \phi_i(x - x_i) + \sum_{i=0}^{\infty} e^{-\lambda_i t} \phi_i(x - x_i) \int_0^t e^{\lambda_i s} f_i^+(s) ds,$$

where

$$f_i^+(t) = \left( \frac{v_i}{R} - \frac{dv_i}{dt} \right) \int_0^1 (1-z) \phi_i(z) dz + \left( \frac{v_{i+1}}{R} - \frac{dv_{i+1}}{dt} \right) \int_0^1 z \phi_i(z) dz.$$

16. Consider the convection diffusion equation

$$u_t - \Delta u + \partial_y(|u|^{q-1}u) = 0$$

set in  $\mathbb{R}^{n-1} \times \mathbb{R} \times \mathbb{R}^+$ , with  $\mathbf{x} = (x_1, \dots, x_{n-1}, y)$ . Assume that  $V$  is a solution with initial datum  $V_0 \in (L^1 \cap L^\infty)(\mathbb{R}^n)$  and  $v$  is a solution with initial datum  $v_0 \in (L^1 \cap L^\infty)(\mathbb{R}^n)$ . Assume that

$$v, V \in C^1([0, T]; L^2(\mathbb{R}^2)) \cap L^\infty([0, T]; H^2(\mathbb{R}^2)) \cap L^\infty((0, T) \times \mathbb{R}^2)$$

for every  $T > 0$ . Then,  $v \leq V$ .

Proof taken from [7, 9]. The function  $w = v - V$  satisfies

$$w_t - \Delta w + \partial_y(|v|^{q-1}v) - \partial_y(|V|^{q-1}V) \leq 0$$

and  $w(0) \leq 0$ . Multiplying the inequality by  $w^+$  and integrating by parts, we obtain

$$\frac{d}{dt} \int \frac{|w^+(t)|^2}{2} d\mathbf{x} + \int |\nabla w^+(t)|^2 d\mathbf{x} \leq \int a w^+(t) \partial_y w^+(t) d\mathbf{x}$$

where  $a(\mathbf{x}, t) = \frac{|v|^{q-1}v - |V|^{q-1}V}{v - V}$  is a bounded function. Integrating in  $t$  and applying Young's inequality we get

$$\frac{\|w^+(t)\|_2^2}{2} + \int_0^t \|\nabla w^+(s)\|_2^2 ds \leq K_1 \int_0^t \|w^+(s)\|_2^2 ds + \varepsilon \int_0^t \|\nabla w^+(s)\|_2^2 ds$$

for  $\varepsilon$  as small as needed. Notice that  $w^+(0) = 0$ . Gronwall's inequality for

$$\|w^+(t)\|_2^2 \leq 2K_1 \int_0^t \|w^+(s)\|_2^2 ds$$

implies  $w^+(t) = 0$ .

17. Prove that the solution of

$$z_t - \Delta z = \mathbf{d} \cdot \nabla(G^q), \quad z(0) = 0$$

can be calculated in terms of heat kernels.

Taken from [19]. Set  $z = \mathbf{d} \cdot \nabla g$  where  $g_t - \Delta g = G^q$ ,  $g(0) = 0$ , that is,

$$g(t) = \int_0^t G(t-s) * G^q(s) ds.$$

18. Express the solution of the transmission heat problem

$$\begin{cases} U_t - \kappa_e \Delta U = 0, & \text{in } \mathbb{R}^N \setminus \overline{\Omega}_i \times (0, \infty), \\ U_t - \alpha_i \kappa_i \Delta U = 0, & \text{in } \Omega_i \times (0, \infty), \\ U^- - U^+ = U_{\text{inc}}, & \text{on } \partial\Omega_i \times (0, \infty), \\ \alpha_i \frac{\partial}{\partial \mathbf{n}} U^- - \frac{\partial}{\partial \mathbf{n}} U^+ = \frac{\partial}{\partial \mathbf{n}} U_{\text{inc}}, & \text{on } \partial\Omega_i \times (0, \infty), \\ U(\cdot, 0) = 0, & \text{in } \mathbb{R}^N, \end{cases}$$

in terms of Helmholtz problems using Laplace transforms.

Taken from [41]. We define  $u_{\text{inc}}$  and  $u$  as the Laplace transforms in time of  $U_{\text{inc}}$  and  $U$ :

$$u_{\text{inc}}(\mathbf{x}, s) = \int_0^\infty e^{-st} U_{\text{inc}}(\mathbf{x}, t) dt, \quad u(\mathbf{x}, s) = \int_0^\infty e^{-st} U(\mathbf{x}, t) dt, \quad \mathbf{x} \in \mathbb{R}^N.$$

For each value of  $s$ , the function  $u_s(\mathbf{x}) := u(\mathbf{x}, s)$  solves

$$\begin{cases} \Delta u_s + \lambda_{s,e}^2 u_s = 0, & \text{in } \mathbb{R}^N \setminus \overline{\Omega}_i, \\ \alpha_i \Delta u_s + \lambda_{s,i}^2 u_s = 0, & \text{in } \Omega_i, \\ u_s^- - u_s^+ = u_{\text{inc},s}, & \text{on } \Gamma, \\ \alpha_i \partial_{\mathbf{n}} u_s^- - \partial_{\mathbf{n}} u_s^+ = \partial_{\mathbf{n}} u_{\text{inc},s}, & \text{on } \Gamma, \end{cases}$$

where  $\lambda_{s,e}^2 := -s/\kappa_e$ ,  $\lambda_{s,i}^2 := -s/\kappa_i$  and  $u_{\text{inc},s}(\mathbf{x}) := u_{\text{inc}}(\mathbf{x}, s)$ . We set  $\Gamma = \partial\Omega_i$ . This problem has a unique solution satisfying the Sommerfeld radiation condition at infinity,

$$\lim_{r \rightarrow \infty} r^{(N-1)/2} (\partial_r u_s - i \lambda_{s,e} u_s) = 0, \quad r = |\mathbf{x}|,$$

for all  $s \in \mathbb{C} \setminus (-\infty, 0]$ . This characterization of  $u_s(\mathbf{x})$  can be used to define and compute  $u(\cdot, s)$  for all  $s \in \mathbb{C} \setminus (-\infty, 0]$ .

The solution of the time-dependent problem is recovered by inverting the Laplace transform:

$$U(\mathbf{x}, t) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{st} u(\mathbf{x}, s) ds.$$

Since  $u(\cdot, s)$  exists for all  $s \in \mathbb{C} \setminus (-\infty, 0]$  and depends holomorphically on  $s$ , many different choices for the inversion path  $\mathcal{C}$  are possible.

19. When the bounded coefficient  $a \geq 0$ , any positive solution  $p$  of the initial value problem

$$\begin{aligned} \frac{\partial}{\partial t} p(t, \mathbf{x}, \mathbf{v}) - \sigma \Delta_{\mathbf{xv}} p(t, \mathbf{x}, \mathbf{v}) + a(t, \mathbf{x}, \mathbf{v}) p(t, \mathbf{x}, \mathbf{v}) &= f(t, \mathbf{x}, \mathbf{v}), \\ p(0, \mathbf{x}, \mathbf{v}) &= p_0(\mathbf{x}, \mathbf{v}), \end{aligned}$$

when  $(\mathbf{x}, \mathbf{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$ ,  $t \in [0, \infty)$ , with  $a \in L^\infty([0, \infty) \times \mathbb{R}^2 \times \mathbb{R}^2)$ ,  $\sigma \in \mathbb{R}^+$ ,  $f \in L^\infty(0, \infty; L^\infty \cap L^1(\mathbb{R}^2 \times \mathbb{R}^2))$  and  $p_0 \in L^\infty \cap L^1(\mathbb{R}^2 \times \mathbb{R}^2)$ , is bounded from above by a solution of a heat equation with the same initial and source data. Moreover, the following estimates hold for any  $q \in [1, \infty)$ :

$$\begin{aligned} \|p\|_q &\leq \|p_0\|_q + t \max_{s \in [0, t]} \|f(s)\|_q, \\ \|p\|_r &\leq C_1 t^{-(\frac{1}{q} - \frac{1}{r}) \frac{n}{2}} \|p_0\|_q + C_2 t^{-(\frac{1}{q} - \frac{1}{r}) \frac{n}{2} + 1} \max_{s \in [0, t]} \|f(s)\|_q, \end{aligned}$$

provided  $r \geq q$ ,  $(\frac{1}{q} - \frac{1}{r}) \frac{n}{2} < 1$ ,  $n = 2$  being the dimension.

Taken from [70]. Notice that  $p$  is the solution of the heat equation with source  $g = f - ap \leq f$ . Let  $u$  be the solution of:

$$\frac{\partial}{\partial t} u(t, \mathbf{x}, \mathbf{v}) - \sigma \Delta_{\mathbf{xv}} u(t, \mathbf{x}, \mathbf{v}) = f(t, \mathbf{x}, \mathbf{v}), \quad u(0, \mathbf{x}, \mathbf{v}) = p_0(\mathbf{x}, \mathbf{v}).$$

This solution admits integral expressions in terms of the heat kernel  $G(t, \mathbf{x}, \mathbf{v})$ . It is then straightforward that:

$$\begin{aligned} p(t) &= G(t) * p_0 + \int_0^t G(t - \tau) * [f(\tau) - a(\tau)p(\tau)] d\tau \\ &\leq u(t) = G(t) * p_0 + \int_0^t G(t - \tau) * f(\tau) d\tau, \end{aligned}$$

where  $*$  denotes convolution in the  $\mathbf{x}, \mathbf{v}$  variables. Setting  $f = 0$ , the well known  $L^r - L^q$  estimates for heat operators  $\|u\|_q = \|G(t) * p_0\|_q$  follow

$$\begin{aligned} \|u\|_q &\leq \|G(t)\|_1 \|p_0\|_q \leq \|p_0\|_q, \\ \|u\|_r &\leq \|G(t)\|_{q'} \|p_0\|_q \leq C_q t^{-(\frac{1}{q} - \frac{1}{r}) \frac{n}{2}} \|p_0\|_q, \quad 1/r = 1/q + 1/q' - 1, \end{aligned}$$

for  $r \geq q$ . When  $f \neq 0$  we find similar estimates for  $u$ . They extend to  $p$  since  $p \leq u$ .

20. We consider a diffusion problem of the form

$$\begin{aligned} \frac{\partial}{\partial t} c(\mathbf{x}, t) &= d \Delta_{\mathbf{x}} c(\mathbf{x}, t) - \eta c(\mathbf{x}, t) j(\mathbf{x}, t) + h(\mathbf{x}, t), \quad \mathbf{x} \in \Omega, t > 0, \\ \frac{\partial c}{\partial r}(\mathbf{x}, t) &= c_{r_0}(\mathbf{x}, t), \quad \mathbf{x} \in S_{r_0}, \quad \frac{\partial c}{\partial r}(\mathbf{x}, t) = 0, \quad \mathbf{x} \in S_{r_1}, \quad t > 0, \\ c(\mathbf{x}, 0) &= c_0(\mathbf{x}), \quad \mathbf{x} \in \Omega, \end{aligned}$$

where  $d, \eta > 0$ ,  $c_{r_0} < 0$  and  $j(\mathbf{x}, t)$  a bounded positive function. The domain  $\Omega = \{\mathbf{x} \in \mathbb{R}^N \mid r_0 < r = |\mathbf{x}| < r_1\}$ , with boundaries  $S_{r_0} = \{\mathbf{x} \in \mathbb{R}^N \mid |\mathbf{x}| = r_0\}$  and  $S_{r_1} = \{\mathbf{x} \in \mathbb{R}^N \mid |\mathbf{x}| = r_1\}$ . Let  $c \in C([0, T]; L^2(\Omega))$  be a solution with initial datum  $c_0 \in L^2(\Omega)$  and boundary condition  $c_{r_0} \in C([0, T]; L^2(\partial\Omega))$ . If  $c_0 \geq 0$ ,  $h \geq 0$  and  $c_{r_0} \leq 0$ , then  $c \geq 0$ .

Taken from [72]. Multiplying the equation

$$\frac{\partial}{\partial t} c(\mathbf{x}, t) = d\Delta_{\mathbf{x}} c(\mathbf{x}, t) - \eta c(\mathbf{x}, t)j(\mathbf{x}, t) + h,$$

by  $c^- = \text{Max}(-c, 0)$  and integrating, we get

$$\begin{aligned} \frac{1}{2} \|c^-(t)\|_2^2 + \int_0^t \int_{\Omega} [|\nabla c^-|^2 + \eta j |c^-|^2] &= \\ \frac{1}{2} \|c^-(0)\|_2^2 - \int_0^t \int_{\partial\Omega} \frac{\partial c}{\partial \mathbf{n}} c^- - \int_0^t \int_{\Omega} h c^- &\leq 0, \end{aligned}$$

since, in our case,

$$- \int_{\partial\Omega} \frac{\partial c}{\partial \mathbf{n}} c^- = - \int_{r=r_1} \frac{\partial c}{\partial r} (r_1) c^- + \int_{r=r_0} \frac{\partial c}{\partial r} (r_0) c^- = \int_{r=r_0} \frac{\partial c}{\partial r} (r_0) c^- \leq 0.$$

This implies that  $c^- = 0$  and  $c \geq 0$ .

21. Construct solutions of the scalar conservation law  $w_t + (c(x)w)_x = x$  with  $w(0) = w_0$ .

Taken from [13, 17]. We set  $v = cw$ . Then,  $v_t + cv_x = 0$ . Thus,  $v$  is constant along the characteristic curves  $x(t)$  solution of  $x'(t) = c(x(t))$ ,  $x(0) = x_0$ , because

$$\frac{d}{dt} v(x(t), t) = v_x(x(t), t)x'(t) + v_t(x(t), t) = 0.$$

Given  $(x, t)$  we may be able to calculate  $x_0(x, t)$  such that the characteristic curve with initial value  $x_0(x, t)$  satisfies  $x(t) = x$ . Then  $v(x, t) = v(x(t), t) = v_0(x_0(x, t))$  and  $w(x, t) = \frac{v_0(x_0(x, t))}{c(x_0(x, t))}$ . The feasibility of this procedure will depend on the function  $c$ .

22. Solve the problem

$$\begin{aligned} \frac{\partial r}{\partial s} + \frac{\partial}{\partial k} (k^{1/3} r) &= 0, \\ \int_0^\infty k r(s, k) dk &= t, \\ \lim_{k \rightarrow 0} k^{1/3} r(s, k) &= 2c. \end{aligned}$$



Taken from [34]. Integrating the equation over  $k > 0$  we find

$$\frac{d}{ds} \int_0^\infty r(s, k) dk = \lim_{k \rightarrow 0} k^{1/3} r(s, k) = 2c(s).$$

Arguing as in the previous exercise, the method of characteristics yields

$$k^{1/3} r(s, k) = 2c(s - a(k))H(s - a(k)),$$

$$a(k) = \frac{3}{2}k^{2/3},$$

in which  $H(x)$  is the Heaviside function (1 for positive  $x$ , 0 otherwise).

23. Obtain an equation for the upper moving boundary  $x_3 = h(x_1, x_2, t)$  of a three dimensional region with lower boundary  $x_3 = 0$  in such a way that the field  $\mathbf{v}$  satisfies  $\text{div } \mathbf{v} = 0$  in it.

Taken from [76]. We integrate  $\text{div } \mathbf{v} = 0$  in the vertical direction to get

$$\int_0^h \frac{\partial(\mathbf{v} \cdot \hat{\mathbf{x}}_1)}{\partial x_1} dx_3 + \int_0^h \frac{\partial(\mathbf{v} \cdot \hat{\mathbf{x}}_2)}{\partial x_2} dx_3 + \int_0^h \frac{\partial(\mathbf{v} \cdot \hat{\mathbf{x}}_3)}{\partial x_3} dx_3 = 0,$$

$\hat{\mathbf{x}}_1$ ,  $\hat{\mathbf{x}}_2$  and  $\hat{\mathbf{x}}_3$  being the unit vectors in the coordinate directions. By Leibniz's rule:

$$\int_0^h \frac{\partial(\mathbf{v} \cdot \hat{\mathbf{x}}_i)}{\partial x_i} dx_3 = \frac{\partial}{\partial x_i} \left[ \int_0^h (\mathbf{v} \cdot \hat{\mathbf{x}}_i) dx_3 \right] - \mathbf{v} \cdot \hat{\mathbf{x}}_i \Big|_h \frac{\partial h}{\partial x_i}, \quad i = 1, 2.$$

Therefore

$$\frac{\partial}{\partial x_1} \left[ \int_0^h (\mathbf{v} \cdot \hat{\mathbf{x}}_1) dx_3 \right] + \frac{\partial}{\partial x_2} \left[ \int_0^h (\mathbf{v} \cdot \hat{\mathbf{x}}_2) dx_3 \right] - \mathbf{v} \cdot \hat{\mathbf{x}}_1 \Big|_h \frac{\partial h}{\partial x_1} - \mathbf{v} \cdot \hat{\mathbf{x}}_2 \Big|_h \frac{\partial h}{\partial x_2} + \mathbf{v} \cdot \hat{\mathbf{x}}_3 \Big|_h = \mathbf{v} \cdot \hat{\mathbf{x}}_3 \Big|_0.$$

Notice that  $\mathbf{v} \cdot \hat{\mathbf{x}}_i = \frac{dx_i}{dt}$ ,  $i = 1, 2, 3$ . Differentiating  $x_3(t) = h(x_1(t), x_2(t), t)$  with respect to time we find

$$\begin{aligned} \mathbf{v} \cdot \hat{\mathbf{x}}_3 \Big|_h &= \frac{dx_3}{dt} = \frac{d}{dt} h(x_1(t), x_2(t), t) = \frac{\partial h}{\partial t} + \frac{\partial h}{\partial x_1} \frac{dx_1}{dt} + \frac{\partial h}{\partial x_2} \frac{dx_2}{dt} \\ &= \frac{\partial h}{\partial t} + \mathbf{v} \cdot \hat{\mathbf{x}}_1 \Big|_h \frac{\partial h}{\partial x_1} + \mathbf{v} \cdot \hat{\mathbf{x}}_2 \Big|_h \frac{\partial h}{\partial x_2}. \end{aligned}$$

Inserting this identity we obtain the equation

$$\frac{\partial h}{\partial t} + \frac{\partial}{\partial x_1} \left[ \int_0^h (\mathbf{v} \cdot \hat{\mathbf{x}}_1) dx_3 \right] + \frac{\partial}{\partial x_2} \left[ \int_0^h (\mathbf{v} \cdot \hat{\mathbf{x}}_2) dx_3 \right] = \mathbf{v} \cdot \hat{\mathbf{x}}_3 \Big|_0.$$

24. Find self-similar solutions  $h(r, t)$  for

$$h_t - K \left(1 + \frac{3}{2}\right) Re^{3t} \frac{1}{r} (rh_r h^3)_r = 0, \quad K = \frac{9\mu_f}{3\xi_\infty^2 \mu_s (1 - \phi_\infty)^2 R_0} h_0^3.$$

Taken from [78]. We have solutions of the form

$$h = R^{-2}e^t f(r) = R^{-2}e^t(1 - \frac{3}{2}r^2)^{\frac{1}{3}}, \quad R = \left(\frac{7}{3}K(1 + \frac{3}{2})(e^{3t} - 1) + 1\right)^{\frac{1}{7}}.$$

25. We work in variable domains  $\Omega^t$ , whose boundaries  $\Gamma^t$  are generated from a smooth curve  $\Gamma^0 \in C^2$  (twice differentiable) following a family of deformations  $\Gamma^t = \{\mathbf{x} + t\mathbf{V}(\mathbf{x}) \mid \mathbf{x} \in \Gamma^0\}$ , along a smooth vector field  $\mathbf{V} \in C^2(\Gamma^0)$ . For  $t > 0$ , we denote by  $u^t \in H^1(B_R)$  the solutions of

$$\begin{aligned} b^t(\Omega^t; u^t, w) &= \ell(w), \quad \forall w \in H^1(B_R), \\ b^t(\Omega^t; u, w) &= \int_{B_R \setminus \bar{\Omega}^t} (\nabla_{\mathbf{x}^t} u \nabla_{\mathbf{x}^t} \bar{w} - \kappa_e^2 u \bar{w}) d\mathbf{x}^t - \int_{\Gamma_R} Lu \bar{w} dS_{\mathbf{x}} \\ &\quad + \int_{\Omega^t} (\beta \nabla_{\mathbf{x}^t} u \nabla_{\mathbf{x}^t} \bar{w} - \beta \kappa_i^2 u \bar{w}) d\mathbf{x}^t, \quad \forall u, w \in H^1(B_R). \end{aligned}$$

Change variables to reformulate the problems on  $\Omega^0$ .

Taken from [79]. For small  $t > 0$ ,  $\Gamma^t \in C^2$  is a perturbation of  $\Gamma^0$ . The deformation  $\mathbf{x}^t = \phi^t(\mathbf{x}) = \mathbf{x} + t\mathbf{V}(\mathbf{x})$  maps  $\Omega^0$  to  $\Omega^t$ . For small  $t$ ,  $\phi^t$  is a diffeomorphism and its inverse  $\eta^t$  maps  $\Omega^t$  to  $\Omega^0$ . The deformation gradient is the jacobian of the change of variables

$$\mathbf{J}^t(\mathbf{x}) = \nabla_{\mathbf{x}} \phi^t(\mathbf{x}) = \left(\frac{\partial x_i^t}{\partial x_j}(\mathbf{x})\right) = \mathbf{I} + t \nabla \mathbf{V}(\mathbf{x}),$$

and its inverse  $(\mathbf{J}^t)^{-1} = \left(\frac{\partial x_i}{\partial x_j^t}\right)$  is the jacobian of the inverse change of variables. Then, volume and surface elements are related by

$$d\mathbf{x}^t = \det \mathbf{J}^t(\mathbf{x}) d\mathbf{x}, \quad dS_{\mathbf{x}^t} = \det \mathbf{J}^t(\mathbf{x}) \|(\mathbf{J}^t(\mathbf{x}))^{-T} \mathbf{n}\| dS_{\mathbf{x}},$$

and the chain rule for derivatives reads  $\nabla_{\mathbf{x}} u(\mathbf{x}^t(\mathbf{x})) = (J^t(\mathbf{x}))^T \nabla_{\mathbf{x}^t} u(\mathbf{x}^t(\mathbf{x}))$ , that is,  $\nabla_{\mathbf{x}^t} u = (\mathbf{J}^t)^{-T} \nabla_{\mathbf{x}} u$ . For each component we have

$$\frac{\partial u}{\partial x_\alpha^t}(\mathbf{x}^t(\mathbf{x})) = \frac{\partial u}{\partial x_k}(\mathbf{x}^t(\mathbf{x})) (J^t)^{-1}_{k\alpha}(\mathbf{x}).$$

We define  $\hat{u}(\mathbf{x}) = u^t \circ \phi^t(\mathbf{x}) = u^t(\mathbf{x}^t(\mathbf{x}))$ . Changing variables we have:

$$\begin{aligned} b_i^t(\Omega^t; u^t, w) &= \int_{\Omega^t} \left[ \beta \frac{\partial u^t}{\partial x_\alpha^t}(\mathbf{x}^t) \frac{\partial \bar{w}}{\partial x_\alpha^t}(\mathbf{x}^t) - \beta \kappa_i^2 u^t(\mathbf{x}^t) \bar{w}(\mathbf{x}^t) \right] d\mathbf{x}^t = \\ &= \int_{\Omega^0} \beta \left[ \frac{\partial \hat{u}}{\partial x_p}(\mathbf{x}) (J^t)^{-1}_{p\alpha}(\mathbf{x}) \frac{\partial \hat{w}}{\partial x_q}(\mathbf{x}) (J^t)^{-1}_{q\alpha}(\mathbf{x}) - \beta \kappa_i^2 \hat{u}(\mathbf{x}) \hat{w}(\mathbf{x}) \right] \det \mathbf{J}^t(\mathbf{x}) d\mathbf{x} \\ &= \hat{b}_i^t(\Omega^0; \hat{u}, \hat{w}). \end{aligned}$$

A similar relation holds on  $B_R \setminus \bar{\Omega}^t$  defining  $b_e^t(B_R \setminus \bar{\Omega}^t; u^t, w) = \hat{b}_e^t(B_R \setminus \bar{\Omega}^0; \hat{u}, \hat{w})$ . Therefore, we obtain the equivalent variational formulation: Find  $\hat{u} \in H^1(B_R)$  such that

$$\hat{b}^t(\Omega^0; \hat{u}, w) = \hat{b}_i^t(\Omega^0; \hat{u}, w) + \hat{b}_e^t(B_R \setminus \bar{\Omega}^0; \hat{u}, w) - \int_{\Gamma_R} L \hat{u} \bar{w} dS_{\mathbf{x}} = \ell(w),$$

for  $w \in H^1(B_R)$ .

26. The plane  $2 \times 2$  strain  $\varepsilon$  and stress  $\sigma$  tensors for a circular plate are given by

$$\sigma_{xx} = \frac{E}{1-\sigma^2}(\varepsilon_{xx} + \sigma\varepsilon_{yy}), \quad \sigma_{yy} = \frac{E}{1-\sigma^2}(\varepsilon_{yy} + \sigma\varepsilon_{xx}), \quad \sigma_{xy} = \frac{E}{1+\sigma}\varepsilon_{xy},$$

$$\varepsilon_{\alpha\beta} = \frac{1}{2} \left( \frac{\partial u_\alpha}{\partial x_\beta} + \frac{\partial u_\beta}{\partial x_\alpha} + \frac{\partial \xi}{\partial x_\alpha} \frac{\partial \xi}{\partial x_\beta} \right), \quad \alpha = x, y,$$

where  $\mathbf{u} = (u_x, u_y)$  are the in-plane displacements in the directions  $x$  and  $y$ , while  $\xi$  is the out-of-plane displacement in the direction  $z$ . The Föppl-Von Karman equations for the equilibrium of a plate of thickness  $h$  yield

$$D\Delta^2\xi - h \frac{\partial}{\partial x_\beta} \left( \sigma_{\alpha\beta} \frac{\partial \xi}{\partial x_\alpha} \right) = 0,$$

$$\frac{\partial \sigma_{\alpha\beta}}{\partial x_\beta} = 0, \quad D = \frac{Eh^3}{12(1-\sigma^2)}.$$

Characterize radial solutions with radial and angular displacements of the form  $u_r = ar + \frac{b}{r}$ ,  $u_\theta = 0$ , where  $r, \theta$  are the standard polar coordinates. The boundary conditions are  $u_r = -\beta$  at  $r = 1/2$  and  $\sigma_{rr} = 0$  at  $r = 1$ . The equations are set in the corona  $1/2 < r < 1$ .

Taken from [65]. We find

$$\sigma_{rr} = -\alpha \left( 1 - \frac{1}{r^2} \right), \quad \sigma_{\theta\theta} = -\alpha \left( 1 + \frac{1}{r^2} \right).$$

The equilibrium equations become

$$\Delta^2\xi + \alpha\Delta\xi + \frac{\alpha}{r} \left( -\frac{\partial^2\xi}{\partial r^2} + \frac{3}{r} \frac{\partial\xi}{\partial r} + \frac{1}{r^2} \frac{\partial^2\xi}{\partial\theta^2} \right) = 0,$$

$$\Delta^2\xi = \frac{\partial^2\xi}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2\xi}{\partial\theta^2} + \frac{1}{r} \frac{\partial\xi}{\partial r},$$

with boundary conditions  $\xi = 0, \frac{\partial\xi}{\partial r} = 0$  at the fixed edge  $r = 1/2$  and

$$-\frac{\partial r \Delta\xi}{\partial r} + (1-\sigma) \frac{1}{r^3} \left( \frac{\partial^2\xi}{\partial\theta^2} - r \frac{\partial^3\xi}{\partial r \partial\theta^2} \right) = 0,$$

$$\Delta\xi + (\sigma-1) \frac{1}{r^2} \left( \frac{\partial^2\xi}{\partial\theta^2} + r \frac{\partial\xi}{\partial r} \right) = 0,$$

at the free end  $r = 1$ . We have solutions of the form  $\xi(r, \theta) = \zeta(r) \cos(m\theta)$  with integer  $m$ . To find them we realize that all possible  $\zeta$  are combinations of two basis solutions  $\zeta$  of a linear differential equation satisfying that  $(\zeta(1/2), \zeta'(1/2), \zeta''(1/2), \zeta'''(1/2))$  is equal to  $(0, 0, 1, 0)$  and  $(0, 0, 0, 1)$ . To select  $\zeta$  fulfilling the conditions at  $r = 1$  we need to choose  $\alpha(m, 1/2)$  numerically, and then choose  $m$ . These patterns provide an example of corona instability in flat plates. For helical instabilities in filaments see [68].

27. Calculate a solution  $\mathbf{u} = (u_1, u_2)$  of  $\Delta \mathbf{u} = (-b_2, b_1)\delta(x)\delta(y)$  for arbitrary  $b_1, b_2 \in \mathbb{R}$ ,  $\delta$  being the standard Dirac mass supported at zero.

Taken from [67]. The function

$$\mathbf{u} = (b_1, b_2) \frac{1}{2\pi} \arctan\left(\frac{y}{x}\right) + (-b_2, b_1) \frac{1}{2\pi} \ln((x^2 + y^2)^{1/2}).$$

This function also satisfies  $\operatorname{div}(\mathbf{u}) = 0$  and  $\int_C [\frac{\partial u_i}{\partial x} dx + \frac{\partial u_i}{\partial y} dy] = b_i, i = 1, 2$ , for contours  $C$  encircling  $(0, 0)$ . These singular solutions represent defects in elastic materials.

### 3 Integrodifferential Equations

1. We know that the problem

$$\begin{aligned} g_t - \Delta_v g + \mathbf{v} \cdot \nabla_x g + \mathbf{E}(\mathbf{x}, t) \cdot \nabla_v g &= 0, & \mathbf{x} \in \mathbb{R}^3, \mathbf{v} \in \mathbb{R}^3, t \in \mathbb{R}^+, \\ g(\mathbf{x}, \mathbf{v}, 0) &= g_0(\mathbf{x}, \mathbf{v}), & \mathbf{x} \in \mathbb{R}^3, \mathbf{v} \in \mathbb{R}^3, \end{aligned}$$

with  $g_0 \in L^1(\mathbb{R}^3 \times \mathbb{R}^3)$  and bounded and Lipschitz  $\mathbf{E}$  admits fundamental solutions  $\Gamma_{\mathbf{E}}$ . The solution of the initial value problem can be expressed as

$$g(\mathbf{x}, \mathbf{v}, t) = \int \Gamma_{\mathbf{E}}(\mathbf{x}, \mathbf{v}, t; \mathbf{x}', \mathbf{v}', 0) d\mathbf{x}' d\mathbf{v}'$$

and  $\Gamma_{\mathbf{E}}$  satisfies the estimates

$$\begin{aligned} |\Gamma_{\mathbf{E}}(\mathbf{x}, \mathbf{v}, t; \mathbf{x}', \mathbf{v}', t')| &\leq C(\|\mathbf{E}\|_{L_{\mathbf{x},t}^\infty}, T) G(\mathbf{x}/2, \mathbf{v}/2, t; \mathbf{x}'/2, \mathbf{v}'/2, t'), \\ |\partial_{v_i} \Gamma_{\mathbf{E}}(\mathbf{x}, \mathbf{v}, t; \mathbf{x}', \mathbf{v}', t')| &\leq C(\|\mathbf{E}\|_{L_{\mathbf{x},t}^\infty}, T) \frac{G(\mathbf{x}/2, \mathbf{v}/2, t; \mathbf{x}'/2, \mathbf{v}'/2, t')}{(t - t')^{1/2}}, \end{aligned}$$

where  $G$  is the fundamental solution for the problem with  $\mathbf{E} = 0$ . Extend these results to problems for which  $\mathbf{E}$  is just bounded.

Taken from [12]. We regularize  $\mathbf{E}$  by convolution and consider  $\mathbf{E}_\delta = \mathbf{E} * \eta_\delta$  where  $\eta_\delta$  is a mollifying family of functions. Then  $\mathbf{E}_\delta$  are bounded and Lipschitz, so for each of them we can construct solutions  $g_\delta$  of the initial value problem and have estimates on the fundamental solutions  $\Gamma_\delta$ . Moreover,  $\|\mathbf{E}_\delta\|_{L_{x,t}^\infty} \leq \|\mathbf{E}\|_{L_{x,t}^\infty}$  and  $\mathbf{E}_\delta \rightarrow \mathbf{E}$  as  $\delta \rightarrow 0$ .

Since  $\Gamma_\delta$  is bounded (locally in  $t$ ) in any  $L_{xvt}^p$  space, a subsequence converges weakly (locally in  $t$ ) in any  $L_{xvt}^p$  (weakly \* if  $p = \infty$ ) to a function  $\Gamma_{\mathbf{E}}$  and we can pass to the limit in the right-hand side of the integral expressions for the solutions  $g_\delta$  in terms of  $\Gamma_\delta$ .

Moreover, the integral expressions imply that  $g_\delta$  are uniformly bounded in any space  $L_{xvt}^p$  with respect to  $\delta$  and locally in  $t$ . Therefore,  $g_\delta$  converges weakly (locally in  $t$ ) in any  $L_{xvt}^p$  space to a function  $g$  and their derivatives also converge in the sense of distributions.

In the distribution sense, the derivatives of  $\Gamma_\delta$  with respect to  $\mathbf{v}$  converge weakly to the derivatives of  $\Gamma_{\mathbf{E}}$ . We can also pass to the limit in the inequalities satisfied by  $\Gamma_\delta$  and establish similar inequalities for  $\Gamma_{\mathbf{E}}$  because  $\|\mathbf{E}_\delta\|_{L_{x,t}^\infty} \leq \|\mathbf{E}\|_{L_{x,t}^\infty}$ .

Now, multiplying the differential equation satisfied by  $g_\delta$  by  $g_\delta$  we get a uniform  $L_{xvt}^2$  bound on  $\nabla_v g_\delta$ . If we multiply the equation by  $|\mathbf{v}|^2$  we get a uniform  $L_{xvt}^1$  bound on  $|\mathbf{v}|^2 g_\delta$ .

Multiplying the differential equations satisfied by  $g_\delta$  by test functions, we can pass to the limit in all the terms of the weak formulation of the equation except in  $\mathbf{E}_\delta \nabla_v g_\delta$  with the convergences already established. The passage to the limit in this term is technical, see details in [12]. Finally,  $g$  is a solution for the initial value problem with bounded  $\mathbf{E}$  and  $\Gamma_{\mathbf{E}}$  an associated fundamental solution.

2. Calculate the equilibrium solution of the Liouville-master equation

$$\begin{aligned} \partial_t \mathcal{P}(x, p, \boldsymbol{\sigma}, t) + \frac{p}{m} \partial_x \mathcal{P}(x, p, \boldsymbol{\sigma}, t) + \left( -m\omega_0^2 x + \mu \sum_{i=1}^n \sigma_i \sigma_{i+1} \right) \partial_p \mathcal{P}(x, p, \boldsymbol{\sigma}, t) \\ = \sum_{i=1}^N [W_i(R_i \boldsymbol{\sigma} | x, p) \mathcal{P}(x, p, R_i \boldsymbol{\sigma}, t) - W_i(\boldsymbol{\sigma} | x, p) \mathcal{P}(x, p, \boldsymbol{\sigma}, t)]. \end{aligned}$$

Taken from [49]. The equilibrium solution of this equation is the canonical distribution

$$\mathcal{P}_{\text{eq}}(x, p, \boldsymbol{\sigma}) = \frac{1}{Z} e^{-\beta \mathcal{H}(x, p, \boldsymbol{\sigma})},$$

where  $Z$  is the partition function

$$Z = \int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dp \sum_{\boldsymbol{\sigma}} e^{-\beta \mathcal{H}(x, p, \boldsymbol{\sigma})},$$

and  $\beta = (k_B T)^{-1}$ . For a study of nonequilibrium behavior see [50].

3. Consider the Fokker-Planck equation for  $p(\boldsymbol{\eta}, t)$

$$\frac{\partial}{\partial t} p = \sum_{j=1}^N \frac{1}{\gamma_j} \frac{\partial}{\partial \eta_j} \left( \frac{\partial G}{\partial \eta_j} p \right) + T \sum_{j=1}^N \frac{1}{\gamma_j} \frac{\partial^2}{\partial \eta_j^2}$$

with  $\boldsymbol{\eta} \in \mathbb{R}^N$  and  $t \geq 0$ , and  $G(\boldsymbol{\eta}) = A(\boldsymbol{\eta}) - FL$ . Prove that if  $F$  and  $L$  are constants, there are explicit stationary solutions.

Taken from [66]. Check that distributions of the form  $p(\boldsymbol{\eta}) \sim e^{-G(\boldsymbol{\eta})/T}$  solve the equation.

4. The hazard rate  $h(t)$ , aging acceleration  $q(t)$  and survival probability  $p(t)$  of an organism according to the DEB (dynamic energy budget) theory are governed by the system

$$h' = q - ah, \quad q' = bq + c, \quad p' = -ph.$$

Find an explicit solution given initial data at  $t = 0$ .

Taken from [73]. Integrating in cascade we find

$$\begin{aligned} q(t) &= q(0)e^{bt} + \int_0^t e^{b(t-s)}c(s)ds, \\ h(t) &= h(0)e^{-at} + \int_0^t e^{-a(t-s)}q(s)ds, \\ p(t) &= p(0)e^{-\int_0^t h(s)ds}. \end{aligned}$$

5. Given a bounded field  $\mathbf{F}(t, \mathbf{x})$  and  $\sigma > 0$ ,  $k > 0$  the initial value problem

$$\begin{aligned} \frac{\partial}{\partial t}p(t, \mathbf{x}, \mathbf{v}) + \mathbf{v} \cdot \nabla_{\mathbf{x}}p(t, \mathbf{x}, \mathbf{v}) + \nabla_{\mathbf{v}} \cdot [(\mathbf{F}(t, \mathbf{x}) - k\mathbf{v})p(t, \mathbf{x}, \mathbf{v})] - \sigma \Delta_{\mathbf{v}}p(t, \mathbf{x}, \mathbf{v}) \\ = f(t, \mathbf{x}, \mathbf{v}), \\ p(0, \mathbf{x}, \mathbf{v}) = p_0(\mathbf{x}, \mathbf{v}), \end{aligned}$$

admits a positive fundamental solution  $\Gamma_{\mathbf{F}}(t, \mathbf{x}, \mathbf{v}; \tau, \boldsymbol{\xi}, \boldsymbol{\nu})$  satisfying

$$\begin{aligned} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Gamma_{\mathbf{F}}(t, \mathbf{x}, \mathbf{v}; \tau, \boldsymbol{\xi}, \boldsymbol{\nu}) d\boldsymbol{\xi} d\boldsymbol{\nu} &= e^{Nk(t-\tau)}, \\ \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Gamma_{\mathbf{F}}(t, \mathbf{x}, \mathbf{v}; \tau, \boldsymbol{\xi}, \boldsymbol{\nu}) d\mathbf{x} d\mathbf{v} &= 1. \end{aligned}$$

Prove that for  $t \in [0, T]$

$$\begin{aligned} \|p(t)\|_1 &\leq \|p_0\|_1 + \int_0^t \|f(\tau)\|_1 d\tau, \\ \|p(t)\|_{\infty} &\leq e^{Nkt} \|p_0\|_{\infty} + \int_0^t e^{Nk(t-\tau)} \|f(\tau)\|_{\infty} d\tau. \end{aligned}$$

Taken from [71]. The solution of the initial value problem is

$$\begin{aligned} p(t, \mathbf{x}, \mathbf{v}) &= \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Gamma_{\mathbf{F}}(t, \mathbf{x}, \mathbf{v}; 0, \boldsymbol{\xi}, \boldsymbol{\nu}) p_0(\boldsymbol{\xi}, \boldsymbol{\nu}) d\boldsymbol{\xi} d\boldsymbol{\nu} + \\ &\int_0^t \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Gamma_{\mathbf{F}}(t, \mathbf{x}, \mathbf{v}; \tau, \boldsymbol{\xi}, \boldsymbol{\nu}) f(\tau, \boldsymbol{\xi}, \boldsymbol{\nu}) d\boldsymbol{\xi} d\boldsymbol{\nu} d\tau. \end{aligned}$$

The estimates on  $p$  follow from the estimates on  $\Gamma_{\mathbf{F}}$ .

## 4 Numerical methods

1. Given a profile  $c_e > 0$ , functions  $\rho(x) > 0$ ,  $n(x) > 0$ ,  $u(x)$  and constants  $a, R > 0$ , we consider the following free boundary problem. We must find  $x^*$  such that

$$\begin{aligned} c''(x) + au(x)c'(x) &= R\rho(x)n(x)^{1/3}(c(x) - c_e(x)), & 0 < x < x_*, \\ c''(x) + au(x)c'(x) &= 0, & x > x_*, \\ c(x_*) = c_e(x_*) = c_*, & c'(x_*^-) = c'(x_*^+), & c(\infty) = 1, c(0) = c_e(0). \end{aligned}$$

Taken from [42]. We write  $c(x) = 1 + \frac{c_* - 1}{\phi(x_*)} \phi(x)$  where

$$\begin{aligned} \phi''(x) + au(x)\phi'(x) &= 0, & x \geq 0, \\ \phi(0) &= 1, \phi(\infty) = 0, \end{aligned}$$

that is,

$$\phi(x) = \int_x^\infty e^{-a \int_0^y u(x') dx'} dy \left( \int_0^\infty e^{-a \int_0^y u(x') dx'} dy \right)^{-1}.$$

To calculate  $x_*$ , we start from a trial value  $x_*$ . Next, we define  $c(x)$  for  $x > x_*$  as explained above for a trial value of  $x_*$ . Then, we solve  $c''(x) + au(x)c'(x) = R\rho(x)n(x)^{1/3}(c(x) - c_e(x))$ ,  $0 < x < x_*$  with  $c(x_*) = c_*$  and  $c'(x_*) = (c_* - 1) \frac{\phi'(x_*)}{\phi(x_*)}$ . Finally, we compare  $c(0)$  with  $c_e(0)$ . Depending on whether it is larger or smaller we increase or decrease  $x_*$  until the difference is small enough.

2. Consider the scheme

$$\begin{aligned} C_{n_1, n_2}^{\ell+1} &= \left[ 1 - 4 \frac{\delta t}{\delta x^2} \kappa \right] C_{n_1, n_2}^\ell + \\ &\frac{\delta t}{\delta x^2} \kappa [C_{n_1+1, n_2}^\ell + C_{n_1-1, n_2}^\ell + C_{n_1, n_2+1}^\ell + C_{n_1, n_2-1}^\ell] \end{aligned}$$

with  $[1 - 4 \frac{\delta t}{\delta x^2} \kappa] \geq 0$ , in a finite lattice of steps  $\delta x$  and  $\delta t$ . If the initial and boundary data are positive, so is  $C_{n_1, n_2}^\ell$  everywhere. Moreover,  $|C_{n_1, n_2}^\ell|$  is bounded from above by the maximum absolute value of the initial data if the boundary data is zero.

Taken from [75]. First, we proceed by induction. If  $C_{n_1, n_2-1}^0 \geq 0$  everywhere, and the data at the  $n_1, n_2$  lattices borders too, then the recurrence implies that  $C_{n_1, n_2-1}^1 \geq 0$  everywhere. In the same way, if  $C_{n_1, n_2-1}^\ell \geq 0$  everywhere, and the data at the  $n_1, n_2$  lattices borders too,  $C_{n_1, n_2-1}^{\ell+1} \geq 0$  everywhere.

Now, set  $V^\ell = \max_{n_1, n_2} |C_{n_1, n_2-1}^\ell|$ . The recurrence implies that

$$V^{\ell+1} \leq \left[ 1 - 4 \frac{\delta t}{\delta x^2} \kappa \right] V^\ell + 4 \frac{\delta t}{\delta x^2} \kappa V^\ell = V^\ell \leq V^0.$$

3. Consider the hyperbolic problem

$$\begin{aligned} \frac{\partial^2 E}{\partial x \partial t} + A \frac{\partial E}{\partial t} + B \frac{\partial E}{\partial x} + C \frac{\partial J}{\partial t} + D &= 0, & x \in (0, L), t > 0, \\ E(x, 0) &= 0, & x \in (0, L), \\ E(0, t) &= \rho J(t), & t \geq 0, \\ \int_0^L E(x, t) dx &= \phi, & t \geq 0, \end{aligned}$$

where  $\rho, \phi, L$  are positive and  $A, B, C, D$  are bounded functions,  $A$  and  $B$  positive, while  $C$  is negative. What would be an adequate numerical scheme to solve this problem?

Hyperbolic problems are typically discretized in explicit ways. However, in this case i) we have an integral constraint which couples all the values at each time level, ii) the hyperbolic operator is given in non characteristic form. We use forward finite differences of first order for first order time derivatives of  $E$  and  $J$ . We use a second order backward approximation scheme for the space derivative of  $E$  because the use of central differences leads to instabilities. The second order derivative  $E_{xt}$  is approximated combining the space and time derivative approximation just described. At the left end we use for the first order spatial derivative of  $E$  a first order backward difference formula. The integral constraint is discretized by means of a composite trapezoidal rule. For a proof of the convergence and stability properties of the scheme see [16].

4. Consider the Navier equations for crystals with cubic symmetry in two dimensional situations, defined by three positive constants  $c_{11}, c_{22}, c_{44}$ :

$$\begin{aligned} Mu_1'' &= C_{11} \frac{\partial^2 u_1}{\partial x_1^2} + C_{12} \frac{\partial^2 u_2}{\partial x_1 \partial x_2} + C_{44} \frac{\partial^2 u_1}{\partial x_2^2} + C_{44} \frac{\partial^2 u_2}{\partial x_1 \partial x_2}, \\ Mu_2'' &= C_{11} \frac{\partial^2 u_2}{\partial x_2^2} + C_{12} \frac{\partial^2 u_1}{\partial x_1 \partial x_2} + C_{44} \frac{\partial^2 u_2}{\partial x_1^2} + C_{44} \frac{\partial^2 u_1}{\partial x_1 \partial x_2}, \end{aligned}$$

where  $M > 0$ . Propose a stable finite difference discretization.

Taken from [31]. Let us construct a rectangular mesh. We denote by  $D_i^+$  and  $D_i^-$  the first order progressive and regressive finite difference equations in the direction  $i$ , that is,

$$\begin{aligned} D_1^+ u_j(\ell, m) &= \frac{u_j(\ell + \delta x_1, m) - u_j(\ell, m)}{\delta x_1}, \\ D_1^- u_j(\ell, m) &= \frac{u_j(\ell, m) - u_j(\ell - \delta x_1, m)}{\delta x_1}, \end{aligned}$$

for  $i = 1$  and analogous expressions for  $i = 2$ . In view of the presence of



cross terms, we choose

$$\begin{aligned} Mu_1'' &= C_{11} \frac{D_1^- D_1^+ u_1}{\delta x_1^2} + C_{12} \frac{D_1^- D_2^+ u_2}{\delta x_1 \delta x_2} + C_{44} \frac{D_2^- D_2^+ u_1}{\delta x_2^2} + C_{44} \frac{D_2^- D_1^+ u_2}{\delta x_1 \delta x_2}, \\ Mu_2'' &= C_{11} \frac{D_2^- D_2^+ u_2}{\delta x_2^2} + C_{12} \frac{D_2^- D_1^+ u_1}{\delta x_1 \delta x_2} + C_{44} \frac{D_1^- D_1^+ u_2}{\delta x_1^2} + C_{44} \frac{D_1^- D_2^+ u_1}{\delta x_1 \delta x_2}. \end{aligned}$$

See [35] for extensions to three dimensional crystals and lattices with two bases.

5. Consider a planar hexagonal graphene lattice and ignore possible vertical deflections. In the continuum limit, in-plane deformations are described by the Navier equations of linear elasticity for the two-dimensional (2D) displacement vector  $(u, v)$ ,

$$\begin{aligned} \rho_2 \frac{\partial^2 u}{\partial t^2} &= (\lambda + 2\mu) \frac{\partial^2 u}{\partial x^2} + \mu \frac{\partial^2 u}{\partial y^2} + (\lambda + \mu) \frac{\partial^2 v}{\partial x \partial y}, \\ \rho_2 \frac{\partial^2 v}{\partial t^2} &= \mu \frac{\partial^2 v}{\partial x^2} + (\lambda + 2\mu) \frac{\partial^2 v}{\partial y^2} + (\lambda + \mu) \frac{\partial^2 u}{\partial x \partial y}, \end{aligned}$$

where  $\rho_2$  is the 2D mass density and  $\lambda$  and  $\mu$  are the 2D Lamé coefficients ( $\lambda = C_{12}$ ,  $\mu = C_{66}$ ,  $\lambda + 2\mu = C_{11}$ ). Propose a finite difference discretization in a hexagonal lattice of constant  $a$ .

Taken from [40]. Consider a point  $A$  in the hexagonal lattice with coordinates  $(x, y)$ . Its 9 (3+6) closest neighbours have coordinates

$$\begin{aligned} n_1 &= \left( x - \frac{a}{2}, y - \frac{a}{2\sqrt{3}} \right), n_2 = \left( x + \frac{a}{2}, y - \frac{a}{2\sqrt{3}} \right), n_3 = \left( x, y + \frac{a}{\sqrt{3}} \right), \\ n_4 &= \left( x - \frac{a}{2}, y + \frac{a\sqrt{3}}{2} \right), n_5 = \left( x + \frac{a}{2}, y + \frac{a\sqrt{3}}{2} \right), n_6 = (x - a, y), \\ n_7 &= (x + a, y), n_8 = \left( x - \frac{a}{2}, y + \frac{a\sqrt{3}}{2} \right), n_9 = \left( x + \frac{a}{2}, y + \frac{a\sqrt{3}}{2} \right). \end{aligned}$$

Let us define the following operators acting on functions of the coordinates  $(x, y)$  of node  $A$ :

$$\begin{aligned} Tu &= [u(n_1) - u(A)] + [u(n_2) - u(A)] + [u(n_3) - u(A)], \\ Hu &= [u(n_6) - u(A)] + [u(n_7) - u(A)], \\ D_1 u &= [u(n_4) - u(A)] + [u(n_9) - u(A)], \\ D_2 u &= [u(n_5) - u(A)] + [u(n_8) - u(A)], \end{aligned}$$

Taylor expansions of these finite difference combinations about  $(x, y)$  yield

$$\begin{aligned} Tu &\sim (\partial_x^2 u + \partial_y^2 u) \frac{a^2}{4}, \\ Hu &\sim (\partial_x^2 u) a^2, \\ D_1 u &\sim \left( \frac{1}{4} \partial_x^2 u + \frac{\sqrt{3}}{2} \partial_x \partial_y u + \frac{3}{4} \partial_y^2 u \right) a^2, \\ D_2 u &\sim \left( \frac{1}{4} \partial_x^2 u - \frac{\sqrt{3}}{2} \partial_x \partial_y u + \frac{3}{4} \partial_y^2 u \right) a^2, \end{aligned}$$

as  $a \rightarrow 0$ . Now we replace in the motion equations  $Hu/a^2$ ,  $(4T - H)u/a^2$  and  $(D_1 - D_2)u/(\sqrt{3}a^2)$  instead of  $\partial_x^2 u$ ,  $\partial_y^2 u$  and  $\partial_x \partial_y u$ , respectively, with similar substitutions for the derivatives of  $v$ , thereby obtaining the following equations at each point of the lattice:

$$\begin{aligned} \rho_2 a^2 \frac{\partial^2 u}{\partial t^2} &= 4\mu Tu + (\lambda + \mu) Hu + \frac{\lambda + \mu}{\sqrt{3}} (D_1 - D_2)v, \\ \rho_2 a^2 \frac{\partial^2 v}{\partial t^2} &= 4(\lambda + 2\mu) Tv - (\lambda + \mu) Hv + \frac{\lambda + \mu}{\sqrt{3}} (D_1 - D_2)u. \end{aligned}$$

6. Consider a planar hexagonal lattice of lattice constant  $a$ . The isotropic Navier equations have singular solutions such as

$$\begin{aligned} u &= \frac{a}{2\pi} \left[ \tan^{-1} \left( \frac{y}{x} \right) + \frac{xy}{2(1-\nu)(x^2 + y^2)} \right], \\ v &= \frac{a}{2\pi} \left[ -\frac{1-2\nu}{4(1-\nu)} \ln \left( \frac{x^2 + y^2}{b^2} \right) + \frac{y^2}{2(1-\nu)(x^2 + y^2)} \right], \end{aligned}$$

where  $\nu = \lambda/[2(\lambda + \mu)]$  for any  $a$ . We choose  $(x_0, y_0)$  different from a lattice point and solve a damped version of the discrete Navier equations formulated in the previous exercise. How would you expect the system to evolve starting from  $(u(x - x_0, y - y_0), v(x - x_0, y - y_0))$ ?

Taken from [38]. The damped equations take the form

$$\begin{aligned} \rho_2 a^2 \frac{\partial^2 u}{\partial t^2} + \gamma \frac{\partial u}{\partial t} &= 4\mu Tu + (\lambda + \mu) Hu + \frac{\lambda + \mu}{\sqrt{3}} (D_1 - D_2)v, \\ \rho_2 a^2 \frac{\partial^2 v}{\partial t^2} + \gamma \frac{\partial v}{\partial t} &= 4(\lambda + 2\mu) Tv - (\lambda + \mu) Hv + \frac{\lambda + \mu}{\sqrt{3}} (D_1 - D_2)u, \end{aligned}$$

with  $\gamma > 0$ . We expect the system to relax to a stationary configuration behaving like  $(u(x - x_0, y - y_0), v(x - x_0, y - y_0))$  at a distance of  $(x_0, y_0)$ . Such solutions represent lattice defects with the chosen elastic far fields. A wide variety of defects is studied in [52, 55].

7. Write the Helmholtz equation set in the whole space

$$\begin{aligned} \Delta u + k^2 u &= 0, & \mathbf{x} \in \mathbb{R}^N, \\ \lim_{r=|\mathbf{x}| \rightarrow \infty} |\mathbf{x}|^{\frac{N-1}{2}} \left( \frac{\partial}{\partial r} (u - u_{\text{inc}}) - ik(u - u_{\text{inc}}) \right) &= 0, \end{aligned}$$

in an equivalent variational form set in a bounded domain by means of the Dirichlet-to-Neumann operator.

Taken from [37]. Let  $B_R$  be a sphere of radius  $R$  and  $\Gamma_R$  its boundary. The Dirichlet-to-Neumann (also called Steklov-Poincaré) operator associates to any Dirichlet data on  $\Gamma_R$  the normal derivative of the solution of the exterior Dirichlet problem:

$$\begin{aligned} L : H^{1/2}(\Gamma_R) &\longrightarrow H^{-1/2}(\Gamma_R) \\ f &\longmapsto \frac{\partial w}{\partial \mathbf{n}} \end{aligned}$$

where  $w \in H_{loc}^1(\mathbb{R}^N \setminus \overline{B_R})$ ,  $B_R := B(\mathbf{0}, R)$ , is the unique solution of

$$\begin{cases} \Delta w + k^2 w = 0, & \text{in } \mathbb{R}^N \setminus \overline{B_R}, \\ w = f, & \text{on } \Gamma_R, \\ \lim_{r \rightarrow \infty} r^{N-1/2} \left( \frac{\partial w}{\partial r} - ikw \right) = 0. \end{cases}$$

$H^{1/2}(\Gamma_R)$  and  $H^{-1/2}(\Gamma_R)$  are standard trace spaces. One can study an equivalent boundary value problem in  $B_R$  with a non-reflecting boundary condition on its boundary  $\Gamma_R$ :

$$\begin{cases} \Delta u + k^2 u = 0, & \text{in } B_R, \\ \frac{\partial}{\partial \mathbf{n}} (u - u_{\text{inc}}) = L(u - u_{\text{inc}}), & \text{on } \Gamma_R. \end{cases}$$

The solution  $u$  also solves the variational equation

$$\begin{cases} u \in H^1(B_R), \\ b(u, v) = \ell(v), \quad \forall v \in H^1(B_R), \end{cases}$$

where

$$\begin{aligned} b(u, v) &= \int_{B_R} (\nabla u \nabla \bar{v} - k^2 u \bar{v}) d\mathbf{x} - \int_{\Gamma_R} L u \bar{v} dl, \quad \forall u, v \in H^1(B_R), \\ \ell(v) &= \int_{\Gamma_R} \left( \frac{\partial u_{\text{inc}}}{\partial \mathbf{n}} - L u_{\text{inc}} \right) \bar{v} dl, \quad \forall v \in H^1(B_R). \end{aligned}$$

8. Write the transmission Helmholtz problem

$$\left\{ \begin{array}{ll} \nabla \cdot (\alpha_e \nabla u) + \lambda_e^2 u = 0, & \text{in } \mathbb{R}^2 \setminus \overline{\Omega}_i, \\ \nabla \cdot (\alpha_i \nabla u) + \lambda_i(k)^2 u = 0, & \text{in } \Omega_i, \\ u^- - u^+ = 0, & \text{on } \partial\Omega_i, \\ \alpha_i \frac{\partial u^-}{\partial \mathbf{n}} - \alpha_e \frac{\partial u^+}{\partial \mathbf{n}} = 0, & \text{on } \partial\Omega_i, \\ \lim_{r \rightarrow \infty} r^{1/2} \left( \frac{\partial}{\partial r} (u - u_{\text{inc}}) - i \lambda_e (u - u_{\text{inc}}) \right) = 0, & r = |\mathbf{x}|, \end{array} \right.$$

in variational form and calculate the derivative of  $J(k) = \int_{\Gamma} |u(k) - d|^2 dl$  with respect to  $k$ .

Taken from [39]. Arguing as in the previous exercise we have

$$\left\{ \begin{array}{l} u \in H^1(B_R), \\ S(\Omega_i; u, v) = \ell(v), \quad \forall v \in H^1(B_R), \end{array} \right.$$

where

$$\begin{aligned} S(\Omega_i; u, v) &:= \int_{B_R \setminus \overline{\Omega}_i} (\alpha_e \nabla u \nabla \bar{v} - \lambda_e^2 u \bar{v}) d\mathbf{x} + \int_{\Omega_i} (\alpha_i \nabla u \nabla \bar{v} - \lambda_i^2 u \bar{v}) d\mathbf{x} \\ &\quad - \int_{\Gamma_R} \alpha_e L u \bar{v} dl, \quad \forall u, v \in H^1(B_R), \\ \ell(v) &:= \int_{\Gamma_R} \alpha_e \left( \frac{\partial u_{\text{inc}}}{\partial \mathbf{n}} - L u_{\text{inc}} \right) \bar{v} dl, \quad \forall v \in H^1(B_R). \end{aligned}$$

where  $L$  denotes the Dirichlet-to-Neumann operator defined by

$$\left\{ \begin{array}{ll} \nabla \cdot (\alpha_e \nabla w) + \lambda_e^2 w = 0, & \text{in } \mathbb{R}^2 \setminus \overline{B}_R, \\ w = f, & \text{on } \Gamma_R, \\ \lim_{r \rightarrow \infty} r^{1/2} \left( \frac{\partial w}{\partial r} - i \lambda_e w \right) = 0. \end{array} \right.$$

Differentiating  $J$  with respect to  $k$  we see that

$$\frac{dJ}{dk} = 2 \int_{\Gamma} \overline{(u(k) - d)} u_k(k) dl,$$

where the derivative  $u_k(k) = \frac{du(k)}{dk} \in H^1(B_R)$  is a solution of

$$\begin{aligned} \int_{B_R \setminus \overline{\Omega}_i} (\alpha_e \nabla u_k(k) \nabla \bar{v} - \lambda_e^2 u_k(k) \bar{v}) d\mathbf{x} + \int_{\Omega_i} (\alpha_i \nabla u_k(k) \nabla \bar{v} - \lambda_i(k)^2 u_k(k) \bar{v}) d\mathbf{x} \\ - \int_{\Gamma_R} \alpha_e L u_k(k) \bar{v} dl = 2 \int_{\Omega_i} \lambda_i(k) \lambda_i'(k) u(k) \bar{v} d\mathbf{x}, \end{aligned}$$

for all  $v \in H^1(B_R)$  and  $u(k)$  the solution of the Helmholtz problem for  $\lambda_i(k)$ .

9. Consider the cost  $J(a, k) = \sum_{m=1}^M \int_{\Gamma} |u_m - d_m|^2$ , where  $u_m$  solves

$$\begin{cases} \operatorname{div}(a_e \nabla u) + k_e^2 u = 0, & \text{in } \mathbb{R}^N \setminus \overline{\Omega}_i, & \operatorname{div}(a \nabla u) + k^2 u = 0, & \text{in } \Omega_i, \\ u^- = u^+, & a \frac{\partial u^-}{\partial \mathbf{n}} = a_e \frac{\partial u^+}{\partial \mathbf{n}}, & \text{on } \partial \Omega_i, \\ r^{(N-1)/2} \left( \frac{\partial(u - u_{\text{inc}}^m)}{\partial r} - \imath k_e (u - u_{\text{inc}}^m) \right) \rightarrow 0, & \text{as } r := |\mathbf{x}| \rightarrow \infty. \end{cases}$$

Given  $a_j, k_j$ , find descent directions for

$$J(\delta) := J(a_j + \delta \phi, k_j + \delta \psi),$$

where  $\delta > 0$ , in order to implement an optimization procedure.

Taken from [46]. We seek  $\delta, \phi$  and  $\psi$  such that  $\frac{dJ(\delta)}{d\delta} < 0$ . Differentiating we find

$$\left. \frac{dJ}{d\delta} \right|_{\delta=0} = - \sum_{m=1}^M \operatorname{Re} \left[ \int_{\Omega_j} [\phi \nabla u_m \nabla \bar{w}_m - 2\psi k_j u_m \bar{w}_m] dz \right],$$

where  $u_m$  solves the forward problem with  $a = a_j$ , and  $k = k_j$ . The adjoint fields  $w_m$  solve

$$\begin{cases} \operatorname{div}(a_e \nabla w_m) + k_e^2 w_m = (d_m - u_m) \delta_{\Gamma_{meas}}, & \text{in } \mathbb{R}^N \setminus \overline{\Omega}_i, \\ \operatorname{div}(a_j \nabla w_m) + k_j^2 w_m = 0, & \text{in } \Omega_i, \\ w_m^- = w_m^+, & a_i \frac{\partial w_m^-}{\partial \mathbf{n}} = a_e \frac{\partial w_m^+}{\partial \mathbf{n}}, & \text{on } \partial \Omega_i, \\ r^{(N-1)/2} \left( \frac{\partial w_m}{\partial r} + \imath \kappa_e w_m \right) \rightarrow 0, & \text{as } r \rightarrow \infty. \end{cases}$$

Setting

$$\phi(\mathbf{x}) = \sum_{m=1}^M \operatorname{Re} (\nabla u_m(\mathbf{x}) \nabla \bar{w}_m(\mathbf{x})), \quad \psi(\mathbf{x}) = - \sum_{m=1}^M \operatorname{Re} (u_m(\mathbf{x}) \bar{w}_m(\mathbf{x})), \quad \mathbf{x} \in \Omega_j,$$

and

$$a_{j+1} = a_j + \delta \phi, \quad k_{j+1} = k_j + \delta \psi,$$

we guarantee  $J(a_{j+1}, k_{j+1}) < J(a_j, k_j)$  for  $\delta$  small.

10. Let  $B_R$  be a sphere centered at  $(0, 0, 0)$  with radius  $R$  and  $k_e > 0, k_i > 0$  two constants. Calculate solutions of

$$\begin{cases} \Delta u + k_e^2 u = 0, & \text{in } \mathbb{R}^3 \setminus \overline{B_R}, \\ \Delta u + k_i^2 u = 0, & \text{in } B_R, \\ u^- = u^+ + U, & \text{on } \partial B_R, \\ \beta \partial_{\mathbf{n}} u^- = \partial_{\mathbf{n}} u^+ + \partial_{\mathbf{n}} U, & \text{on } \partial B_R, \\ \lim_{r \rightarrow \infty} r(\partial_r u - \imath k_e u) = 0, & \end{cases}$$

for given smooth functions  $U$  as a series expansion.

Taken from [74]. We have

$$\begin{aligned} u(\mathbf{x}) &= \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{nm} h_n^{(1)}(k_e |\mathbf{x}|) Y_n^m(\hat{\mathbf{x}}), \quad |\mathbf{x}| \geq R, \\ u(\mathbf{x}) &= \sum_{n=0}^{\infty} \sum_{m=-n}^n b_{nm} j_n(k_i |\mathbf{x}|) Y_n^m(\hat{\mathbf{x}}), \quad |\mathbf{x}| \leq R, \end{aligned}$$

where  $\mathbf{x} = |\mathbf{x}|\hat{\mathbf{x}}$ ,  $j_n$  are the spherical Bessel functions of the first kind,  $h_n^{(1)}$  are the spherical Hankel functions and  $Y_n^m$  are the standard spherical harmonics,

$$Y_n^m(\theta, \phi) = \sqrt{\frac{2n+1}{4\pi} \frac{(n-|m|)!}{(n+|m|)!}} P_n^{|m|}(\cos(\theta)) e^{im\phi},$$

for associated Legendre polynomials  $P_n^{|m|}$ . More precisely, if  $U$  can be expanded as

$$U(\mathbf{x}) = \sum_{n=0}^{\infty} \sum_{m=-n}^n u_{nm} j_n(k_e |\mathbf{x}|) Y_n^m(\hat{\mathbf{x}})$$

in a ball containing  $B_R$ , the coefficients are computed as follows.

On the boundary of the sphere  $|\mathbf{x}| = R$ , the transmission conditions hold. We impose these relations on the inner and outer series expansions and equate the coefficients of  $Y_n^m(\hat{\mathbf{x}})$  since the spherical harmonics form a basis in  $L^2(\partial B_1)$ . This yields the relations:

$$\begin{aligned} u_{nm} j_n(k_e R) + a_{nm} h_n^{(1)}(k_e R) - b_{nm} j_n(k_i R) &= 0, \\ u_{nm} k_e j_n'(k_e R) + a_{nm} k_e h_n^{(1)'}(k_e R) - \beta b_{nm} k_i j_n'(k_i R) &= 0. \end{aligned}$$

Solving the system we obtain the value of the coefficients:

$$\begin{aligned} a_{nm} &= u_{nm} a_n(R) = u_{nm} \frac{k_e j_n(k_i R) j_n'(k_e R) - \beta k_i j_n'(k_i R) j_n(k_e R)}{\beta k_i j_n'(k_i R) h_n^{(1)}(k_e R) - k_e j_n(k_i R) h_n^{(1)'}(k_e R)}, \\ b_{n,m} &= u_{nm} b_n(R) = u_{nm} \frac{k_e j_n'(k_e R) h_n^{(1)}(k_e R) - k_e j_n(k_e R) (h_n^{(1)})'(k_e R)}{\beta k_i j_n'(k_i R) h_n^{(1)}(k_e R) - k_e j_n(k_i R) h_n^{(1)'}(k_e R)}. \end{aligned}$$

To calculate these coefficients, notice that the spherical Bessel function is related to the Bessel functions of the first kind by  $j_n(s) = \sqrt{\frac{\pi}{2s}} J_{n+1/2}(s)$ . The spherical Hankel function is related to the Hankel functions of the first kind by  $h_n^{(1)}(s) = \sqrt{\frac{\pi}{2s}} H_{n+1/2}(s)$ . Their derivatives are evaluated using the formula  $f_n'(s) = \frac{n}{s} f_n(s) - f_{n+1}(s)$ , which holds for both  $j_n$  and  $h_n^{(1)}$ .

11. Explain how to solve the following equations using the deterministic particle method:

$$\partial_t f + \frac{\Delta l}{2\hbar v_M} \sin(k) \partial_x f + \frac{\tau_e}{\eta} F \partial_k f = \frac{1}{\eta} \left[ f^{FDa}(k; \mu(n)) - \left( 1 + \frac{\nu_{imp}}{2\nu_{en}} \right) f + \frac{\nu_{imp}}{2\nu_{en}} f(x, -k, t) \right],$$

$$\partial_x^2 V = \partial_x F = n - 1$$

$$n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x, k, t) dk = \frac{1}{2\pi} \int_{-\pi}^{\pi} f^{FDa}(k; \mu(n)) dk$$

$$f^{FDa}(k; \mu) = \alpha \ln [1 + \exp(\mu - \delta + \delta \cos(k))]$$

$$\eta = \frac{v_M}{\nu_{en} x_0} \quad \delta = \frac{\Delta}{2k_B T}.$$

The boundary conditions are, for  $x = 0$ :

$$f^+ = \beta F - \frac{f^{(0)}}{\int_0^\pi \sin(k) f^{(0)} dk} \int_{-\pi}^0 \sin(k) f^- dk$$

with

$$\beta = \frac{2\pi \hbar \sigma F_M}{e \Delta N_D}$$

and for  $x = L/x_0$ :

$$f^- = \frac{f^{(0)}}{(1/(2\pi)) \int_{-\pi}^0 f^{(0)} dk} \left( 1 - \frac{1}{2\pi} \int_0^\pi f^+ dk \right)$$

The boundary conditions for the electric potential  $V$  are

$$V(0, t) = 0, \quad V(L, t) = \phi_L \sim \frac{\phi}{F_M} \frac{L}{x_0}.$$

The initial condition is

$$f^{(0)}(k; n) = \sum_{j=-\infty}^{\infty} \exp(\nu j k) \frac{1 - \nu j F / \tau_e}{1 + j^2 (F)^2} f_j^{FD}(n)$$

$$f_j^{FD}(n) = \frac{1}{\pi} \int_0^\pi f^{FD}(k; \mu(n)) \cos(jk) dk$$

with  $x \in [0, L = L/x_0]$  and  $f$  periodic in  $k$  with period  $2\pi$ . The average energy  $E$  is defined as

$$E = \frac{E}{k_B T} = \frac{\int_{-\pi/l}^{\pi/l} \varepsilon(k) f(x, k, t) dk}{k_B T \int_{-\pi/l}^{\pi/l} f(x, k, t) dk} = \delta \frac{\int_{-\pi}^{\pi} (1 - \cos k) f(x, k, t) dk}{\int_{-\pi}^{\pi} f(x, k, t) dk}.$$

Taken from [43]. We rely on particle description of the distribution function, which means that  $f(x, k, t)$  is written as a sum of delta functions

$$f(x, k, t) \approx \sum_{i=1}^N \omega_i f_i(t) \delta(x - x_i(t)) \otimes \delta(k - k_i(t))$$

where  $\omega_i$ ,  $f_i(t)$ ,  $x_i(t)$  and  $k_i(t)$  are, respectively, the (constant) control volume, the weight, the position and the wave vector of the  $i$ th particle.  $N$  is the number of numerical particles. The motion of particles is governed by collisionless dynamics, whereas the collisions are accounted for by the variation of weights. Large gradients in the solution profile arise from appropriate particles acquiring large weights, not by accumulating many particles in the large gradient regions. The evolution of the particles is determined by their positions and wave vectors which are the characteristic curves of the convective part of the equation. Their equations are:

$$\frac{d}{dt}k = \frac{\tau_e}{\eta} F, \quad \frac{d}{dt}x = \frac{\Delta l}{2\hbar v_M} \sin(k).$$

The evolution of the distribution function over these characteristic curves is given by the ordinary differential equation:

$$\frac{d}{dt}f = \frac{1}{\eta} \left[ - \left( 1 + \frac{\nu_{imp}}{2\nu_{en}} \right) f + \frac{\nu_{imp}}{2\nu_{en}} f(-k) + f^{FD} \right].$$

The system of ordinary differential equations is now discretized by using a modified Euler method:

$$f_i^n = f_i^{n-1} + dt \frac{1}{\eta} \left[ - \left( 1 + \frac{\nu_{imp}}{2\nu_{en}} \right) f_i^{n-1} + \frac{\nu_{imp}}{2\nu_{en}} f_i^{(-k)} + f_i^{FD, n-1} \right]$$

with  $f_i^{(-k)} = f(x_i^{n-1}, -k_i^{n-1}, t^{n-1})$ ,

$$k_i^n = k_i^{n-1} + dt \frac{\tau_e}{\eta} F_i^{n-1},$$

$$x_i^n = x_i^{n-1} + dt \frac{\Delta l}{2\hbar v_M} \sin(k_i^n).$$

For stability reasons, we use  $k_i^n$  to update  $x_i^n$ . We have also used multi-step methods but they yield worse results.

The boundary conditions are taken into account as follows:

- If  $k_i^n > \pi$ , we set  $k_i^n = k_i^n - 2\pi$ . If  $k_i^n < -\pi$ , we set  $k_i^n = k_i^n + 2\pi$ .
- If  $x_i^n > L$ , we set  $x_i^n = x_i^n - L$  and  $f_i^{n-1} = f_i^+$ . If  $x_i^n < 0$ , we set  $x_i^n = x_i^n + L$  and  $f_i^{n-1} = f_i^-$ . Here  $f_i^+$  and  $f_i^-$  are calculated by discretization of the integrals using Simpson's rule on an equally spaced mesh  $K_{m'}$  with step  $\Delta k$ .



To calculate  $x_i$ ,  $k_i$  and  $f_i$  at the next time step  $t^{n+1}$ , we need to update the electric field and the Fermi-Dirac distribution in the equations for the particles. This updating requires an interpolation procedure to generate an approximation of the distribution function on a regular mesh  $X_m, K_{m'}$  which is then used to approximate the electric field and the chemical potential. To approximate the values of the distribution function over the mesh,  $f_{m,m'}^n$ , we use its values for the particles,  $f_i^n$ . The idea is obtain a weighted mean by:

$$f_{m,m'}^n = \frac{\sum_{i=1}^N f_i^n W_{m,m'}^i}{\sum_{i=1}^N W_{m,m'}^i}$$

where

$$W_{m,m'}^i = \max \left\{ 0, 1 - \frac{|X_m - x_i^n|}{\Delta x} \right\} \cdot \max \left\{ 0, 1 - \frac{|K_{m'} - k_i^n|}{\Delta k} \right\}$$

and  $\Delta x$  and  $\Delta k$  are the spatial and wave vector steps.

An approximation for the density and average energy at the mesh points,  $n(X_m, t^n) \approx n_m^n$  and  $(k_B T)^{-1} E(X_m, t^n) \approx (k_B T)^{-1} E_m^n$ , are obtained using Simpson's rule and the interpolated values of the distribution function on the mesh.

We calculate the nondimensional chemical potential  $\mu$  by using a Newton-Raphson iterative scheme to solve the equations. The extended Simpson's rule is employed to approximate the integrals for  $n(\mu)$  and  $dn(\mu)/d\mu$ . Once we know the chemical potential  $\mu$ , we find the Fermi-Dirac distribution function at mesh points,  $f^{FD}(K_{m'}; n_m^n)$ , which is then interpolated to get the Fermi-Dirac distribution function for the particles.

To compute the electric field at time  $t^n$ , we use finite differences to discretize the Poisson equation on the grid  $X_m$ :

$$\begin{aligned} V_{m+1}^n - 2V_m^n + V_{m-1}^n &= n_m^n - 1, \\ F_m^n &= \frac{V_{m+1}^n - V_{m-1}^n}{2\Delta x}. \end{aligned}$$

Here  $V(0, t^n) = 0$  and  $V(L, t^n) = \phi L$ . Let  $V_m^n$  and  $F_m^n$  denote our approximations of  $V(X_m, t^n)$  and  $F(X_m, t^n)$  on the equally spaced mesh  $X_m$ . Finally, the electric field is interpolated at the location of the particle  $i$

$$F_i^n = \left( \frac{X_{m+1} - x_i^n}{\Delta x} \right) F_m^n + \left( \frac{x_i^n - X_m}{\Delta x} \right) F_{m+1}^n.$$

The total current density  $J$  is given by

$$J(t) = \frac{\zeta}{L} \int_0^L \left[ \int_{-\pi}^{\pi} \sin(k) f(x, k, t) dk \right] dx,$$

in which

$$\varsigma = \frac{l\Delta}{4\pi\hbar v_M}.$$

We use the Simpson rule to approximate  $J(t^n)$ .

12. Consider a set of particles at points  $\mathbf{r}_i$  in a rectangular box. We use them as seeds to generate a Voronoi tessellation of the rectangular region, with vertices  $\mathbf{r}_\mu$ . Sketch equations for the motion of the vertices due to self interaction.

Taken from [80]. Each configuration of the mesh has the following associated energy

$$E_{\text{VM}} = \sum_{i=1}^N \left[ \frac{K_i}{2} (A_i - A_i^0)^2 + \frac{\Gamma_i}{2} P_i^2 \right] + \sum_{\langle \mu, \nu \rangle} \Lambda_{\mu\nu} l_{\mu\nu}.$$

Here  $N$  is the total number of polygon,  $A_i$  is the area of polygon  $i$ ,  $A_i^0$  is its reference area, and  $K_i$  is the area modulus, i.e., a constant with units of energy per area squared measuring how hard it is to change the area of the polygon.  $P_i$  is the polygon perimeter and  $\Gamma_i$  (with units of energy per length squared) is the perimeter modulus that determines how hard it is to change perimeter  $P_i$ .  $l_{\mu\nu}$  is the length of the junction between vertices  $\mu$  and  $\nu$ , and  $\Lambda_{\mu\nu}$  is the tension of that junction (with units of energy per length). The sum in the last term is over all pairs of vertices that share a junction. The area  $A_i$  of polygon  $\Omega_i$ , given by the following discrete version of Green's formula:

$$A_i = \frac{1}{2} \sum_{\mu \in \Omega_i} (\mathbf{r}_\mu \times \mathbf{r}_{\mu+1}) \cdot \mathbf{N}_i,$$

where  $\mathbf{r}_\mu$  is the position of vertex  $\mu$ , and  $\mathbf{N}_i$  is a unit vector perpendicular to the surface of the polygon. The sum is over all vertices of the Voronoi polygon and we close the loop with  $\mu + 1 = 1$  when  $\mu$  equals the total number of vertices in the polygon,  $N_{\Omega_i}$ . The polygon perimeter is

$$P_i = \frac{1}{2} \sum_{\mu \in \Omega_i} |\mathbf{r}_\mu - \mathbf{r}_{\mu+1}|.$$

The relation between the vertices  $\mathbf{r}_\mu$  of the Voronoi polygons and the vertices  $\mathbf{r}_i$  of the Delaunay triangles (seeds of the Voronoi polygons, i.e., centers) is

$$\mathbf{r}_\mu = \frac{\lambda_1 \mathbf{r}_i + \lambda_2 \mathbf{r}_j + \lambda_3 \mathbf{r}_k}{\lambda_1 + \lambda_2 + \lambda_3}.$$

The usual dynamics for the polygon centers is a gradient flow of the energy with forces  $\mathbf{F}_i$

$$\gamma \mathbf{r}'_i = \mathbf{F}_i$$

with

$$\begin{aligned}
\mathbf{F}_i = & - \sum_{k=1}^N \frac{K_k}{2} (A_k - A_k^0) \sum_{\nu \in \Omega_k} [\mathbf{r}_{\nu+1, \nu-1} \times \mathbf{N}_k]^T \left[ \frac{\partial \mathbf{r}_\nu}{\partial \mathbf{r}_i} \right] \\
& - \sum_{k=1}^N \Gamma_k P_k \sum_{\nu \in \Omega_k} (\hat{\mathbf{r}}_{\nu, \nu-1} - \hat{\mathbf{r}}_{\nu+1, \nu})^T \left[ \frac{\partial \mathbf{r}_\nu}{\partial \mathbf{r}_i} \right] \\
& - \sum_{k=1}^N \sum_{\nu \in \Omega_k} [\Lambda_{\nu-1, \nu} \hat{\mathbf{r}}_{\nu, \nu-1} - \Lambda_{\nu, \nu+1} \hat{\mathbf{r}}_{\nu+1, \nu}]^T \left[ \frac{\partial \mathbf{r}_\nu}{\partial \mathbf{r}_i} \right] \\
& + k \sum_{\langle j, i \rangle} (2a - |\mathbf{r}_i - \mathbf{r}_j|) \frac{\mathbf{r}_i - \mathbf{r}_j}{|\mathbf{r}_i - \mathbf{r}_j|} \Theta(2a - |\mathbf{r}_i - \mathbf{r}_j|).
\end{aligned}$$

Here  $\left[ \frac{\partial \mathbf{r}_\nu}{\partial \mathbf{r}_i} \right]$  is the  $3 \times 3$  Jacobian matrix connecting coordinates of cell centres with coordinates of the dual Voronoi tessellation, and the non-commutative row-matrix product  $[\cdot]^T [\cdot]$  is a  $3 \times 1$  column vector.  $\Theta(x) = 1$  if  $x > 0$ , else  $\Theta(x) = 0$ , is the Heaviside unit step function. We have included a range repulsive force of short range  $a$  that avoids self intersections of the triangulation for very obtuse triangles.

## 5 Differential-Difference Equations

1. Consider the equation

$$x'' + \frac{1}{2\alpha\theta} \frac{1 + \tanh^2(x/\theta)}{1 - \tanh^2(x/\theta)} x' + x - H - \tanh\left(\frac{x}{\theta}\right) = 0.$$

Study the equilibria and the behavior of the trajectories in terms of the control parameters  $\theta$  and  $H$ .

Taken from [56]. We introduce the potential  $V(x; H, \theta) = \frac{x^2}{2} - Hx - \theta \ln \cosh\left(\frac{x}{\theta}\right)$ . Typically,  $\theta \in (0, 1)$ . The equation becomes

$$x'' + \frac{1}{2\alpha\theta} R(x, \theta) x' - V'(x; H, \theta) = 0$$

with  $R(x, \theta) = \frac{1 + \tanh^2(x/\theta)}{1 - \tanh^2(x/\theta)} > 0$ . For  $H = 0$  and  $\theta < 1$ , the potential has two equally deep minima at symmetric positions. In view of the presence of a damping term, trajectories wrap around these points (spiral attractors). For  $|H| < H_c$ , there are two minima  $x_+ > 0$  and  $x_- < 0$ , each of them with a basin of attraction.

2. Consider a system with energy  $A(\eta, Y) = \sum_{j=1}^N a(\eta_j; Y)$ ,  $\eta = (\eta_1, \dots, \eta_N)$  under the constraint  $\sum_{j=1}^N \eta_j = L$ . Given  $F$ , study the minima of  $A(\eta, Y) - FL$ , where  $F = F(L)$  is a Lagrange multiplier to be calculated in such a way that the constraint holds.

Taken from [61]. The curve  $F(L)$  has  $N+1$  branches, that we can compute imposing  $\frac{\partial a}{\partial \eta_j} = F$  for all  $j$ .

3. Consider the differential difference equation  $u'_n(t) = u_{n+1} - 2u_n + u_{n-1} - A \sin(u_n)$ , where  $A$  is a positive parameter. Prove that there is a monotone solution such that  $u_{-\infty} = 0$  and  $u_{\infty} = 2\pi$  with  $u_0 = \pi$  and  $u_n - \pi = \pi - u_{-n}$  for all  $n$ .

Taken from [14]. We set  $u_0 = \pi$  and vary  $u_1$  in the interval  $(\pi, 2\pi)$  to find the desired solution. The condition  $u_0 = \pi$  ensures that  $u_n - \pi$  is an odd function of  $n$ . We first choose  $\epsilon > 0$  so that  $-A \sin(u) > \epsilon(u - \pi)$  for  $\pi < u \leq \frac{3}{2}\pi$ . Then, we choose  $N$  large so that  $\epsilon(N-1) > 1$ . Next, we choose  $u_1 - \pi$  small so that  $u_j \leq \frac{3}{2}\pi$  for  $1 \leq j \leq N$ . We wish to show that under these conditions, the finite sequence  $\{u_1, \dots, u_N\}$  is not monotone increasing. It is convenient to let  $U_n = u_n - \pi$ . If  $\{U_1, \dots, U_N\}$  is monotone increasing, then  $2 \leq j \leq N$  and  $U_j \leq (2 - \epsilon)U_{j-1} - U_{j-2}$ . Adding these inequalities results in  $U_N - U_{N-1} \leq \epsilon \sum_{i=2}^{N-1} U_i + (1 - \epsilon)U_1$ . Since we assumed that  $U_i \geq U_1$  for  $2 \leq i \leq N$ , our lower bound on  $N$  then shows that  $U_N < U_{N-1}$ , a contradiction. Therefore, we have shown that for sufficiently small  $U_1$ , the sequence starts to decrease before crossing  $\pi$ . On the other hand, we have simply to choose  $U_1 > \pi$  to have the sequence cross  $\pi$  before decreasing. Note that if the sequence increases until some first  $N$  such that  $U_N = \pi$ , then  $U_{N+1} > \pi$ . If, finally, there is an  $N$  such that the sequence increases up to  $n = N$ , with  $U_N < \pi$ , and  $U_N = U_{N+1}$ , then  $U_{N+2} < U_{N+1}$  so that the sequence decreases before reaching  $\pi$ .

4. Let  $U_i(t)$  and  $L_i(t)$ ,  $i \in \mathbb{Z}$  be differentiable sequences such that

$$\begin{aligned} U'_i(t) - d_1(U_i)(U_{i+1} - U_i) - d_2(U_i)(U_{i-1} - U_i) - f(U_i) &\geq \\ L'_i(t) - d_1(L_i)(L_{i+1} - L_i) - d_2(L_i)(L_{i-1} - L_i) - f(L_i) & \end{aligned}$$

and  $U_i(0) < L_i(0)$  for all  $i$ , where  $f$ ,  $d_1 > 0$  and  $d_2 > 0$  are Lipschitz continuous functions. Then,  $U_i(t) > L_i(t)$  for all  $t > 0$  and  $i \in \mathbb{Z}$ .

Taken from [15]. By contradiction, set  $W_i(t) = U_i(t) - L_i(t)$ . At  $t = 0$ ,  $W_i(0) > 0$  for all  $i$ . Let us assume that  $W_i$  changes sign after a certain minimum time  $t_1 > 0$ , at some value of  $i$ ,  $i = k$ . Thus  $W_k(t_1) = 0$  and  $W'_k(t) \leq 0$ , as  $t \rightarrow t_1$ . We shall show that this is contradictory. At  $t = t_1$ , there must be an index  $m$  (equal or different from  $k$ ) such that  $W_m(t_1) = 0$ , while its next neighbor  $W_{m+j}(t_1) > 0$  ( $j$  is either 1 or  $-1$ ), and  $W_i(t_1) = 0$  for all indices between  $k$  and  $m$ . For otherwise  $W_k$  should be identically 0 for all  $k$ . The differential inequality implies

$$W'_m(t_1) \geq d_1(U_m(t_1))W_{m+1}(t_1) + d_2(U_m(t_1))W_{m-1}(t_1) > 0.$$

This contradicts the fact that  $W'_m(t)$  should have been nonpositive as  $t \rightarrow t_1$ , for  $W_m(t_1)$  to have become zero in the first place.

5. Consider the equation

$$U'(t) = z_1(F/A) + z_3(F/A) - 2U(t) - A \sin(U(t)) + F,$$

for  $|F| < A$ ,  $A \gg 1$  where  $z_1(F/A) < z_2(F/A) < z_3(F/A)$  are three consecutive solutions of the equation  $\sin(z) = F/A$  in one period. Prove that there is a critical value  $F_c$  such that this equation has three stable constant solutions if  $0 \leq F < F_c$  but one if  $F > F_c$ . Characterize  $F_c$ .

Taken from [18]. When  $F = 0$ ,  $z_1(0) = 0$ ,  $z_2(0) = \pi$  and  $z_3(0) = 2\pi$ . We need to solve

$$2z + A \sin(z) = F + 2\arcsin(F/A) + 2\pi.$$

As we increase  $F$  from 0, we keep on finding three solutions  $z_1(F/A) < z_2(F/A) < z_3(F/A)$  continuing these branches until  $F + 2\arcsin(F/A) + 2\pi$  hits the first local maximum of  $2z + A \sin(z)$  (remember that  $A$  is large). The value  $F_c$  at which this happens is characterized by the existence of a double zero, a value  $u_0$  such that  $2 + A \cos(u_0) = 0$  and  $2u_0 + A \sin(u_0) = F_c + 2\arcsin(F_c/A) + 2\pi$ . Then,  $u_0 = \arccos(-2/A)$  and  $F_c$  is the solution of  $2u_0(A) + A \sin(u_0(A)) = F_c + 2\arcsin(F_c/A) + 2\pi$ . Below  $F_c$  we have three zeroes, at  $F_c$  two collapse, above  $F_c$  the collapsing ones,  $z_1(F/A)$  and  $z_2(F/A)$  are lost.

$z_1(F/A)$  and  $z_3(F/A)$  are stable while they exist. This picture corresponds to a saddle node bifurcation in the system, see [18]. These bifurcations are essential to understand a variety of biological phenomena, see [64].

6. The system of equations

$$\frac{dE_i}{dt} + \frac{v(E_i)}{\nu}(E_i - E_{i-1}) - \frac{D(E_i)}{\nu}(E_{i+1} - 2E_i + E_{i-1}) = J - v(E_i),$$

for  $i \in \mathbb{Z}$  admits traveling wave solutions of the form  $E_i(t) = E(i - ct)$  propagating at constant velocity  $c$  when the parameter  $J$  is large enough. Here,  $v, D$  are positive functions and  $\nu > 0$  is large.  $v$  is a cubic, it grows from 0 to a local maximum, decreases to a positive minimum, and increases to infinity later. Justify that the wavefront velocity scales as  $(J - J_c)^{1/2}$  where  $J_c$  is the threshold for existence of travelling waves.

Taken from [20]. For  $\nu$  large, we can construct stationary solutions, which can be approximated by

$$E_i \sim z_1(J) \quad i < 0, \quad E_i \sim z_3(J) \quad i > 0,$$

for  $|J| < J_c$ , while  $E_0$  solves

$$J - v(E_0) - \frac{v(E_0)}{\nu}(E_0 - z_1(J)) + \frac{D(E_0)}{\nu}(z_3(J) - 2E_0 + z_1(J)) = 0,$$

where  $z_1(J) < z_2(J) < z_3(J)$  are solutions of  $J = v(z)$ . At a value  $J_c$ ,  $z_1(J_c) = z_2(J_c)$  and these roots are lost for  $J > J_c$ , only  $z_3(J)$  remains. The reduced equation

$$\frac{dE_0}{dt} = J - v(E_0) - \frac{v(E_0)}{\nu}(E_0 - z_1(J)) + \frac{D(E_0)}{\nu}(z_3(J) - 2E_0 + z_1(J)),$$

for the middle point undergoes a saddle node bifurcation at  $J_c$  with normal form

$$\phi' = \alpha(J_c)(J - J_c) + \beta(J_c)\phi^2,$$

which has solutions of the form  $\sqrt{\frac{\alpha}{\beta}}(J - J_c) \tan(\sqrt{\alpha\beta}(J - J_c)(t - t_0))$ , blowing up when the argument of the tangent approaches  $\pm\pi/2$ , over a time  $t - t_0 \sim \pi/\sqrt{\alpha\beta}(J - J_c)$ . This value  $J_c$  separates the regime for which we have stationary (pinned) wave front solutions and travelling wave front solutions. It marks the depinning transition.

Now, for  $J > J_c$  but close to  $J_c$ , simulations show staircase like wave profiles, in which a point stays near the vanished equilibrium  $E_0(J_c)$  until it moves following the tangent path given by the normal form and is replaced at position  $E_0(J_c)$  by a neighbouring one, once and again. The wave velocity is the reciprocal of the time this transition takes  $c(J, \nu) \sim \frac{\sqrt{\alpha\beta}(J - J_c)}{\pi}$ , see [20] for details.

7. We consider a problem with noise

$$\frac{du_i}{dt} = u_{i+1} - 2u_i + u_{i-1} + F - A \sin(u_i) + \gamma\xi_i,$$

where  $A > 0$  is large and  $\gamma > 0$  characterizes the disorder strength and  $\xi_i$  is a zero mean random variable taking values on an interval  $(-1, 1)$  with equal probability. Show that the speed of the wavefronts for  $F$  larger than the critical value  $F_c^*$  scales as  $(F - F_c^*)^{3/2}$ .

Taken from [22]. Setting  $\gamma = 0$ , we can repeat with this equation the study done in the previous exercise and obtain a velocity that scales like  $(F - F_c)^{1/2}$ . However, when we add noise, for each realization of the noise, the threshold  $F_c$  is shifted slightly up or down by the noise. The observed velocity will be the average of the velocities observed for a large number of realizations. If

$$|c_R| \sim \frac{1}{\pi} \sqrt{\alpha(F_c)\beta(F_c)(F - F_c) + \gamma\beta(F_c)\xi_0}$$

the average

$$\bar{c} = \frac{1}{N} \sum_{R=1}^N |c_R| = \frac{1}{2\pi} \int_{-1}^1 (\alpha\beta(F - F_c) + \gamma\beta\xi)^{1/2} d\xi \sim (F - F_c^*)^{3/2}$$

where the new critical field is  $F_c^* = F_c - \frac{\gamma}{\alpha}$ .

8. Consider the problem

$$\frac{du_i}{dt} = u_{i+1} - 2u_i + u_{i-1} + F - A \sin(u_i),$$

with  $A$  large. Let  $z_1(F/A) < z_2(F/A) < z_3(F/A)$  be the three consecutive branches of zeros of  $F - A \sin(z) = 0$  which start at  $z_1(0) = 0$ ,  $z_2(0) = \pi$ ,  $z_3(0) = 2\pi$ . We know that for  $|F| < F_c(A)$  the problem admits stationary solutions increasing from  $z_1(F/A)$  at  $-\infty$  to  $z_3(F/A)$  at  $\infty$ . When  $F$  surpasses that threshold, we have travelling wave solutions. Write the equation for such travelling wave solutions and find a formula for the velocity.

Taken from [24]. Travelling wave solutions have the form  $u_i(t) = u(i - ct)$ , where  $c$  is a constant wave speed and  $u(z)$ ,  $z = i - ct$  is a wave profile, which solve

$$-cu_z(z) = u(z+1) - 2u(z) + u(z-1) + F - A \sin(u(z)), \quad z \in \mathbb{R}$$

with  $u(-\infty) = z_1(F/A)$  and  $u(\infty) = z_3(F/A)$ . These type of travelling wave solutions are called fronts. Multiplying the equation by  $u_z$  and integrating, we find

$$-c \int_{-\infty}^{\infty} u_z^2 dz = F [z_3(F/A) - z_1(F/A)].$$

9. The discrete Fitz Hugh-Nagumo system is a typical model for pulse propagation

$$\begin{aligned} \epsilon u'_i &= d(u_{i+1} - 2u_i + u_{i-1}) + u_i(2 - u_i)(u_i - a) - v_i, \\ v'_i &= u_i - Bv_i. \end{aligned}$$

when the parameter values  $\epsilon, d > 0$  and  $a$  are such that  $(0, 0)$  is the only constant solution.  $\epsilon$  is small and  $a$  is such that  $z(2 - z)(z - a)$  has three roots  $z_1(a) < z_2(a) < z_3(a)$ . Explain how to describe the evolution of pulse solutions in terms of front solutions for Nagumo type equations

$$\epsilon u'_i = d(u_{i+1} - 2u_i + u_{i-1}) + u_i(2 - u_i)(u_i - a) - w.$$

Taken from [25]. Pulse-like solutions take the form  $u_i(t) = u(z)$ ,  $v_i(t) = v(z)$ ,  $z = i - ct \in \mathbb{R}$ , with

$$\begin{aligned} -\epsilon u_z(z) &= d(u(z+1) - 2u(z) + u(z-1)) + u(z)(2 - u(z))(a - u(z)) - v, \\ -\epsilon v_z(z) &= 0, \end{aligned}$$

for  $z \in \mathbb{R}$ . For small enough  $v$ , we denote by  $z_1(a, v) < z_2(a, v) < z_3(a, v)$  the three roots of  $u(z)(2 - u(z))(a - u(z)) - v = 0$ . Since  $\epsilon$  is small,  $u_i$  and  $v_i$  evolve in different time scales. We distinguish 5 regions in a pulse like solution

- Pulse front:  $u_i = z_1(a, v_i)$  and  $v'_i = z_1(a, v_i) - Bv_i$ , which evolves to  $(0, 0)$  as  $i$  grows.
- Pulse leading edge: Described by a traveling solution of  $\epsilon u'_i = d(u_{i+1} - 2u_i + u_{i-1}) + u_i(2 - u_i)(u_i - a) - 0$  which decreases from 2 to 0, with  $v_i \sim 0$ . It travels at speed  $c$ .
- Pulse peak:  $u_i = z_2(a, v_i)$  and  $v'_i = z_3(a, v_i) - Bv_i$ .
- Pulse trailing edge: Described by a traveling solution of  $\epsilon u'_i = d(u_{i+1} - 2u_i + u_{i-1}) + u_i(2 - u_i)(u_i - a) - w$  which increases from 0 to 2, with  $v_i \sim w$ ,  $w$  selected in such a way that it travels with speed  $c$  too.
- Pulse tail:  $u_i = z_1(a, v_i)$  and  $v'_i = z_1(a, v_i) - Bv_i$ , which evolves to  $(0, 0)$  as  $i$  decreases.

See [25] for a visualization. See [32] for an application of these ideas to Hodgkin-Huxley models for myelinated nerves. Pulse solutions fail to propagate when the leading pulse cannot move because for the parameters we use the reduced from equation has only stationary front solutions, they are pinned.

10. Consider the system

$$\begin{aligned} v'_j &= d(v_{j+1} - 2v_j + v_{j-1}) + f(v_j, w_j), \\ w'_j &= \lambda g(v_j, w_j), \end{aligned}$$

with  $d, \lambda > 0$  and  $\lambda$  is small, for the two variables to evolve in different scales. For  $w$  fixed,  $f(v, w)$  is a 'bistable cubic', that is, it has three zeros, two of which are stable. When  $f(v, w) = 0 = g(v, w)$  has a unique solution, which is stable, we have pulse like solutions for the differential system, as for Fitz Hugh-Nagumo. When it is unstable, show that oscillating solutions appear.

Taken from [33]. When  $g$  and  $f$  intersect at a stable zero, we have an excitable system displaying pulse like solutions. When they intersect at an unstable zero, limit cycle solutions  $(V(t), W(t))$  with period  $T$ ,  $T > 0$  of

$$v' = f(v, w), \quad w' = \lambda g(v, w),$$

for  $\lambda$  small, play a role. The trajectories of the system behave like  $v_j(t) = V(t + \phi_j)$  and  $w_j(t) = W(t + \phi_j)$ , for a slowly varying phase  $\phi_j$  which may become independent of  $t$  as  $t \rightarrow \infty$ . All the trajectories are then synchronized.

11. Let  $u_{i,j}(t)$  be a solution to

$$\frac{\partial u_{i,j}}{\partial t} = u_{i-1,j} - 2u_{i,j} + u_{i+1,j} + A(\sin(u_{i,j-1} - u_{i,j}) \sin(u_{i,j+1} - u_{i,j}))$$



for  $i, j \in \mathbb{Z}$  and  $u_{i,j}(0) = \alpha_{i,j}$  satisfying  $\alpha_{i+1,j} - 2\alpha_{i,j} + \alpha_{i-1,j} \in l^2$ ,  $\sin(\alpha_{i,j-1} - \alpha_{i,j}) \sin(\alpha_{i,j+1} - \alpha_{i,j}) \in l^2$  and  $\alpha_{i,j} \in l_{\text{loc}}^\infty$ . If  $(u_{i,j+1} - u_{i,j})(t) \in \cap_{n \in \mathbb{Z}} [-\frac{\pi}{2} + 2n\pi, \frac{\pi}{2} + 2n\pi]$  holds for all  $i, j, t$ , then  $u_{i,j}(t)$  tends to a limit  $s_{i,j}$  as  $t \rightarrow 0$  which is a stationary solution of the problem.

Taken from [23]. Define  $w_{i,j}(t) = u_{i,j}(t+\tau) - u_{i,j}(t)$  for some  $\tau > 0$ . Then

$$\begin{aligned} \frac{d}{dt} \left( \frac{1}{2} \sum_{i,j} |w_{i,j}(t)|^2 \right) &= - \sum_{i,j} ((w_{i+1,j} - w_{i,j})(t))^2 - \sum_{i,j} (\sin((u_{i,j+1} - u_{i,j})(t+\tau)) \\ &\quad - \sin((u_{i,j+1} - u_{i,j})(t))) ((u_{i,j+1} - u_{i,j})(t+\tau) - (u_{i,j+1} - u_{i,j})(t)) \leq 0. \end{aligned}$$

This implies  $w_{i,j}(t) \rightarrow 0$  as  $t \rightarrow \infty$  for every  $i, j$ . In conclusion,  $u_{i,j}(t)$  tends to a limit  $s_{i,j}$  which is a stationary solution of the problem.

12. We solve

$$\frac{\partial u_{i,j}}{\partial t} = u_{i-1,j} - 2u_{i,j} + u_{i+1,j} + A(\sin(u_{i,j-1} - u_{i,j}) \sin(u_{i,j+1} - u_{i,j}))$$

with boundary conditions  $s_{i,j} = \theta(i, j/\sqrt{A}) + Fj$  where  $\theta$  is the angle function from 0 to  $2\pi$  and  $F > 0$  is a control parameter. For  $F = 0$ , the previous exercise ensures existence of stationary solutions. Can you expect a change as  $F$  grows?

Taken from [26]. As  $F$  grows, the condition

$$(u_{i,j+1} - u_{i,j})(t) \in \cap_{n \in \mathbb{Z}} \left[ -\frac{\pi}{2} + 2n\pi, \frac{\pi}{2} + 2n\pi \right]$$

will fail. Stationary solutions will disappear and travelling patterns will be observed. Notice that if we linearize the spatial operator about  $s_{i,j}$ , we have a discrete elliptic problem for  $F$  small but it changes type as  $F$  grows.

13. We construct numerically solutions of

$$\begin{aligned} m \frac{\partial^2 u_{i,j}}{\partial t^2} + \alpha \frac{\partial u_{i,j}}{\partial t} &= u_{i-1,j} - 2u_{i,j} + u_{i+1,j} \\ &\quad + A(\sin(u_{i,j-1} - u_{i,j}) \sin(u_{i,j+1} - u_{i,j})) \end{aligned}$$

in a square lattice  $i = 1, \dots, N_x$ ,  $j = 1, \dots, N_y$ , with boundary conditions  $u_{i,j} = F(j - (N_y + 1)/2)$ . This is equivalent to 'shearing' the lattice. As  $F$  grows, we observe that the initial zero solution for  $F = 0$  changes into slowly varying stationary solutions until we reach a point  $F_c$  past which the lattice structure is distorted in two main different ways. Linearizing the problem at  $F = F_c$  we find a zero eigenvalue for the resulting matrix, while all the eigenvalues are negative for  $F < F_c$ . How do you explain this?

Taken from [36]. The branch of stationary solutions  $s_{i,j}(F)$  seems stable. At  $F = F_c$  and two new branches appear. The system undergoes a pitchfork bifurcation.

14. We construct numerically solutions of

$$m \frac{\partial^2 v_{i,j}}{\partial t^2} + \alpha \frac{\partial v_{i,j}}{\partial t} = v_{i-1,j} - 2v_{i,j} + v_{i+1,j} + A(\sin(v_{i,j-1} - v_{i,j}) \sin(v_{i,j+1} - v_{i,j}))$$

in a square lattice  $i = 1, \dots, N_x$ ,  $j = 1, \dots, N_y$ . We set the boundary conditions representing a 'push down' from the central top part:

- Left-hand side:  $v_{1,j} = v_{0,j}$ .
- Right-hand side:  $v_{N_x,j} = v_{N_x+1,j}$ .
- Left-hand-side of the top layer ( $1 \leq i < p_1$ ):  $v_{i,N_y} = v_{i,N_y+1}$ .
- Right-hand-side of the top layer ( $p_2 < i \leq N_x$ ):  $v_{i,N_y} = v_{i,N_y+1}$ .
- Bottom layer of the domain:  $v_{i,0} = 0$ .
- Central atoms ( $p_1 \leq i \leq p_2$ ) are pushed downwards according to:  $v_{i,N_y+1} - v_{i,N_y} = -f(i)$ , where  $f$  has a triangular profile, pointing downwards, with magnitude  $F > 0$ .

As  $F$  grows, we observe that the initial zero solution for  $F = 0$  develops localized lattice distortions that travel downwards. As we decrease  $F$  to zero the distortions travel upwards and may disappear. How do you explain that?

Taken from [45]. The branch of stationary solutions that starts at  $F = 0$  develops bifurcations at specific values of  $F$  at which lattice with different distortions are created. Such new branches are stable for some ranges of  $F$ , while the defects simply travel. The configuration bifurcates at new  $F$  values, new distortions are created, that travel for while, and the process is repeated as  $F$  grows. When we decrease  $F$ , the process is reversed. Created distortions travel upwards, and disappear.

15. Consider the problem

$$u_j'' + \alpha u_j' = u_{j+1} - 2u_j + u_{j-1} + F - Ag(u_j),$$

where  $g(u) = u + 1$  if  $u < 0$  and  $g(u) = u - 1$  if  $u > 0$ . Construct traveling wave front solutions.

Taken from [27]. A traveling wave front solution takes the form  $u_i(t) = u(i - ct)$ ,  $z = i - ct$ . The profile  $v(z) = u(z) + 1$  satisfies

$$\begin{aligned} c^2 v_{zz}(z) - \alpha c v_z(z) - (v(z+1) - 2v(z) + v(z-1)) + Av(z) \\ = F + 2AH(-\text{sign}(cF)z), \quad z \in \mathbb{R}, \end{aligned}$$

with  $v(-\infty) = 0$  and  $v(\infty) = 2$ . We have written  $g(u) = u + 1 - 2H(u)$ , where  $u$  is the Heaviside function. Using the complex contour integral expression for the Heaviside function

$$H(-z) = -\frac{1}{2\pi i} \int_C \frac{e^{ikx}}{k} dk.$$

$C$  is a contour formed by a closed semicircle in the upper complex plane oriented counterclockwise and another one oriented clockwise in the lower half plane, which includes zero inside and forms a small semicircle around it. The profile we seek admits the expression

$$v(z) = \frac{F}{A} - \frac{A}{\pi i} \int_C \frac{\exp(ik \operatorname{sign}(cF)z) dk}{k A + 4 \sin^2(k/2) - k^2 c^2 - ik|c|\alpha \operatorname{sign}(F)}.$$

Imposing  $v(0) = 1$  we obtain a relation between the velocity  $c$  and the applied force  $F$ . Once we know  $c(F)$ , the above expression provides the profiles  $v$ . Unlike previous exercises, such profiles are not monotonic, but display oscillations, see [27].

16. Show that the initial value problem

$$\begin{aligned} u_j'' + \alpha u_j' &= d(u_{j+1} - 2u_j + u_{j-1}) - u_j + F, \\ u_j(0) &= u_j^0, \quad u_j'(0) = u_j^1, \end{aligned}$$

$d > 0$ ,  $\alpha \geq 0$ , admits solutions of the form

$$u_j(t) = \sum_k [G_{j,k}^0(t) u_k^1(0) + G_{j,k}^1(t) u_k^0(0)] + \int_0^t \sum_k G_{j,k}^0(t-s) f_k(s) ds$$

for adequate Green functions  $G_{j,k}^0$  and  $G_{j,k}^1$ .

Taken from [28]. Firstly, we get rid of the difference operator by using the generating functions  $p(\theta, t)$  and  $f(\theta, t)$

$$p(\theta, t) = \sum_j u_j(t) e^{-vj\theta}, \quad f(\theta, t) = \sum_j f_j(t) e^{-vj\theta}.$$

Differentiating  $p$  with respect to  $t$  and using the equation, we see that  $p$  solves the ordinary differential equation

$$p''(\theta, t) + \alpha p'(\theta, t) + \omega(\theta)^2 p(\theta, t) = f(\theta, t)$$

with  $\omega(\theta)^2 = 1 + 4d \sin^2(\theta/2)$  and initial conditions for  $p$  from those for  $u_j$ . Fixed  $\theta$  we know how to calculate explicit solutions of this linear second order equation with constant coefficients to get

$$p(\theta, t) = p(\theta, 0) g^0(\theta, t) + p'(\theta, 0) g^1(\theta, t) + \int_0^t g^1(\theta, t-s) f(\theta, s) ds,$$

for

$$g^0(\theta, t) = \begin{cases} \frac{e^{r_+(\theta)t} - e^{r_-(\theta)t}}{r_+(\theta) - r_-(\theta)}, & \alpha^2/4 > \omega^2(\theta), \\ t e^{-\alpha t/2}, & \alpha^2/4 = \omega^2(\theta), \\ e^{-\alpha t/2} \frac{\sin(I(\theta)t)}{I(\theta)}, & \alpha^2/4 < \omega^2(\theta), \end{cases}$$

$$g^1(\theta, t) = \begin{cases} \frac{e^{r_+(\theta)t} r_+(\theta) - e^{r_-(\theta)t} r_-(\theta)}{r_+(\theta) - r_-(\theta)}, & \alpha^2/4 > \omega^2(\theta), \\ te^{-\alpha t/2} \left(1 + \frac{\alpha}{2}t\right), & \alpha^2/4 = \omega^2(\theta), \\ e^{-\alpha t/2} \left(\cos(I(\theta)t) + \frac{\alpha \sin(I(\theta)t)}{2I(\theta)}\right), & \alpha^2/4 < \omega^2(\theta). \end{cases}$$

We recover  $u_j$  as

$$u_j(t) = \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} e^{ij\theta} p(\theta, t),$$

and find

$$G_{jk}^0(t) = \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} e^{i(j-k)\theta} g^0(\theta, t), \quad G_{jk}^1(t) = \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} e^{i(j-k)\theta} g^1(\theta, t).$$

17. *Use the expression of the solutions of the initial value problem established before to define a nonreflecting boundary condition at  $n = 0$  for truncated problems set in  $n \geq 0$ , so that the solution we obtain is the same we would obtain solving the system for all  $n$ .*

Taken from [48]. We place an artificial boundary at  $n = 0$  and restrict the computational domain to the region  $n \geq 0$ . Thus, we need a boundary condition to compute  $u_0(t)$  and close the system. In principle,

$$\frac{d^2 u_0}{dt^2} = d(u_1 - 2u_0 + u_{-1}) + f_0,$$

but  $u_{-1}(t)$  is unknown unless we solve also for  $n \leq 0$ . The equation at  $n = -1$  can be rewritten as:

$$\frac{d^2 u_{-1}}{dt^2} = d(0 - 2u_{-1} + u_{-2}) + f_{-1} + du_0.$$

Assuming we know  $u_0(t)$ , the problem for  $n \leq 0$  with boundary condition  $u_0(t)$  can be seen as a problem with zero boundary condition at the wall and a modified source term:  $f_n + d\delta_{n,-1}u_0$  for  $n < 0$ . We can extend this problem to the whole space setting:

$$v_n = \begin{cases} u_n & n < 0 \\ 0 & n = 0 \\ -u_{-n} & n > 0 \end{cases}$$

The extension  $v_n$  solves:

$$\frac{d^2 v_n}{dt^2} = d(v_{n+1} - 2v_n + v_{n-1}) + g_n, \\ v_n(0) = v_n^0, \quad \frac{dv_n}{dt}(0) = v_n^1,$$

for all  $n$ , where  $v_n^0$  and  $v_n^1$  are odd extensions of  $u_n^0$  and  $u_n^1$ . The source  $g_n$  is obtained extending  $f_n + \delta_{n,-1}u_0$ . We have included the boundary

condition  $u_0$  as a force acting on  $u_{-1}$  to allow for an odd extension with  $v_0 = 0$ . Using the symmetry of the data:

$$u_n(t) = v_n(t) = \sum_{n' < 0} \left[ \mathcal{G}_{n,n'}^0(t) \frac{du_{n'}}{dt}(0) + \frac{d\mathcal{G}_{n,n'}^0}{dt}(t) u_{n'}(0) \right] \\ + \int_0^t \sum_{n' < 0} \mathcal{G}_{n,n'}^0(t-s) (f_{n'}(s) + d\delta_{n',-1} u_0(s)) ds, \quad n < 0$$

where  $\mathcal{G}_{n,n'}^0 = G_{n,n'}^0 - G_{n,-n'}^0$  is the Green function for the half space  $n < 0$  with zero boundary condition at  $n = 0$ . In this way, we obtain the desired formula for  $u_{-1}$ :

$$u_{-1}(t) = r_{-1}(t) + d \int_0^t \mathcal{G}_{-1,-1}^0(t-s) u_0(s) ds, \\ r_{-1}(t) = \sum_{n' < 0} \left[ \mathcal{G}_{-1,n'}^0(t) \frac{du_{n'}}{dt}(0) + \frac{d\mathcal{G}_{-1,n'}^0}{dt}(t) u_{n'}(0) \right. \\ \left. + \int_0^t \mathcal{G}_{-1,n'}^0(t-s) f_{n'}(s) ds \right].$$

The term  $r_{-1}(t)$  represents the contribution of the data in the outer region. Our boundary condition at  $n = 0$  takes the form:

$$\frac{d^2 u_0}{dt^2} = d \left( u_1 - 2u_0 + d \int_0^t \mathcal{G}_{-1,-1}^0(t-s) u_0(s) ds \right) + dr_{-1} + f_0,$$

where the kernel is:

$$\mathcal{G}_{-1,-1}^0(t) = \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{1 - e^{-2i\theta}}{\omega(\theta)} \sin(\omega(\theta)t).$$

In a similar way, we can set no reflecting boundary conditions in finite intervals  $-N \leq n \leq N$ , see [48].

18. Consider the initial value problem

$$u_j'' = d(u_{j+1} - (2+r)u_j + u_{j-1}) + f(u_j), \quad j = 1, \dots, N \\ u_j(0) = u_j^0, \quad u_j'(0) = u_j^1, \quad j = 1, \dots, N \\ u_0(t) = u_{N+1}(t) = 0,$$

for a continuous function  $f$ . Set  $V(u) = -\int_0^u f(s) ds$ . Assume  $uf(u) + 2(2\sigma + 1)V(u) \geq 0$  for  $\sigma > 0$ . Define the energy

$$E(t) = \frac{1}{2} \sum_{j=-\infty}^{\infty} u_j'^2(t) + \frac{d}{2} \sum_{j=-\infty}^{j=\infty} [(u_{j+1} - u_j)^2(t) + ru_j^2(t)] + \sum_{j=-\infty}^{j=\infty} V(u_j(t)).$$

If  $E(0) < 0$ , then  $\sum_{j=1}^N |u_j(t)|^2 \rightarrow \infty$  as  $t \rightarrow T$  for some finite  $T > 0$ .

Taken from [29]. We define  $H(t) = \sum_{j=1}^N |u_j(t)|^2 + \rho(t + \tau)^2$ ,  $\rho, \sigma > 0$  to be selected so that  $(H^{-\sigma})'' = \sigma H^{-\sigma-2}((\sigma + 1)(H')^2 - HH'') \leq 0$ . When  $H(0) \neq 0$  we have

$$H^\sigma(t) \geq H^{\sigma+1}(0)(H(0) - \sigma t H'(0))^{-1}$$

and  $H(t)$  blows up at some time  $T \leq H(0)/\sigma H'(0)$  provided  $H'(0) > 0$ . Let us explain how to do this. We calculate  $H'$  and  $H''$ , and use the equation to get

$$HH'' - (\sigma + 1)(H')^2 = 4(\sigma + 1)Q + 2HG,$$

$$Q = \left( \sum_{j=1}^N |u_j|^2 + \rho(t + \tau)^2 \right) \left( \sum_{j=1}^N |u'_j|^2 + \rho \right) - \left( \sum_{j=1}^N u_j u'_j + \rho(t + \tau) \right)^2,$$

$$G = \sum_{j=1}^N u_j f(u_j) - \sum_{i,j} u_i a_{i,j} u_j - (2\sigma + 1) \left( \sum_{j=1}^N |u'_j|^2 + \rho \right),$$

where  $\mathbf{A} = (a_{ij})$  is the matrix defining the linear part of the system. We have  $Q \geq 0$ . We estimate  $G'(t)$  to find  $G(t) \geq \sigma(2\sigma + 1) \left( -\frac{\rho}{2} - E(0) \right) \geq 0$  for  $\rho = -2E(0) > 0$ .

We have  $(H^{-\sigma})'' \leq 0$  and  $H(0) \neq 0$ . Moreover,  $H'(0) = 2 \sum_{j=1}^N u_j^0 u_j^1 + 2\rho\tau > 0$  if  $\tau > -\rho^{-1} \sum_{j=1}^N u_j^0 u_j^1$ .

19. Let  $u_n(t)$  be a solution of

$$u'_n = d(u_n)(u_{n+1} - 2u_n + u_{n-1}) + v(u_n)(u_{n-1} - u_n) + f(u_n),$$

with non negative initial data and a strong reactive source  $f$ , such that  $f(u) > Cu^p$ , with  $p > 1$ ,  $C > 0$ , when  $u > 0$  large. We set  $a(u) = -(2d(u) + v(u))u + f(u)$  and assume that  $d(u) > 0$ ,  $d(u) + v(u) > 0$  grow slower than  $u^p$  for  $u$  large. For any component  $k$  such that  $a(u_k(0)) > 0$  and  $a'(u) > 0$  when  $u > u_k(0)$

$$u_k(t) \rightarrow \infty \quad \text{as } t \rightarrow T \leq T_b = \int_{u_k(0)}^{\infty} \frac{ds}{a(s)} < \infty.$$

Taken from [44]. In all cases, a maximum principle ensures the positivity of  $u_n(t)$  everywhere. Using  $u_{k+1}, u_{k-1} \geq 0$ , we obtain the differential inequality  $u'_k(t) \geq a(u_k)$ . By hypothesis,  $a(u) > a(u_k(0)) > 0$  for  $u \geq u_k(0)$ . Then  $u_k(t)$  is increasing and it is bounded from below by the solution  $y(t)$  of  $y'(t) = g(y)$ ,  $y(0) = u_k(0)$ , which is given implicitly by:

$$t = \int_{u_k(0)}^{y(t)} \frac{ds}{a(s)}.$$

The integral  $\int_{u_k(0)}^{\infty} \frac{ds}{a(s)} < \infty$  due to the growth condition  $a(s) \gg s^p$ ,  $p > 1$  for  $s$  large, since  $a(u) > 0$  for  $u \geq u_k(0)$ . When  $t \rightarrow T_b = \int_{u_k(0)}^{\infty} \frac{ds}{a(s)} < \infty$ ,  $y(t) \rightarrow \infty$ .

20. Consider the Becker-Döring equations

$$\begin{aligned}\sum_{k=1}^{\infty} k\rho_k &= \rho > 0, \\ \rho'_k &= j_{k-1} - j_k, \quad k \geq 2, \\ j_k &= d_k(e^{aD+\epsilon_k} \rho_1 \rho_k - \rho_{k+1})\end{aligned}$$

for a given sequence  $\epsilon_k > 0$  with  $D+\epsilon_k = \epsilon_{k+1} - \epsilon_k$ , with  $a$  and  $\rho$  positive constants. Calculate the equilibrium distributions.

Taken from [30]. We set  $j_k = 0$ . Then  $\rho_k = \rho_1^k e^{a\epsilon_k}$ . This system admits traveling wavefront solutions, see [30].

21. Consider the kinetic system

$$\begin{aligned}\frac{dr_k}{ds} &= (k-1)^{1/3}D(k-1)r_{k-1} - k^{1/3}D(k)r_k, \quad k \geq 3, \\ \frac{dr_2}{ds} &= 2cD(1) - 2^{1/3}D(2)r_2, \\ c\frac{dc}{ds} + 4c^2D(1) + cM_{\frac{1}{3}} &= 1, \\ \frac{dt}{ds} &= \frac{1}{c}.\end{aligned}$$

Find an expression for  $r_k$  in terms of the parameter problems.

Taken from [51]. Notice that the equations for  $s$  and  $c$  start from a singularity at  $s = 0$ . Laplace transforming the equations:

$$\begin{aligned}\frac{dr_2}{ds} &= 2cD(1) - 2^{1/3}D(2)r_2, \\ \frac{dr_k}{ds} &= (k-1)^{1/3}D(k-1)r_{k-1} - k^{1/3}D(k)r_k, \quad k \geq 3.\end{aligned}$$

we find:

$$\begin{aligned}\hat{r}_2(\sigma) &= \frac{2D(1)}{\sigma + 2^{\frac{1}{3}}D(2)}\hat{c}, \\ \hat{r}_k(\sigma) &= \frac{(k-1)^{\frac{1}{3}}D(k-1)}{\sigma + k^{\frac{1}{3}}D(k)}\hat{r}_{k-1}, \quad k \geq 3.\end{aligned}$$

Therefore,

$$\begin{aligned}2^{\frac{1}{3}}D(2)\hat{r}_2(\sigma) &= \frac{2D(1)}{1 + \sigma 2^{\frac{-1}{3}}D(2)^{-1}}\hat{c}, \\ k^{\frac{1}{3}}D(k)\hat{r}_k(\sigma) &= \frac{(k-1)^{\frac{1}{3}}D(k-1)}{1 + \sigma k^{\frac{-1}{3}}D(k)^{-1}}\hat{r}_{k-1}, \quad k \geq 3.\end{aligned}$$

By iteration,

$$k^{\frac{1}{3}}D(k)\hat{r}_k = 2\hat{c}D(1)\hat{R}_k,$$

where

$$\hat{R}_k(\sigma) = \prod_{j=2}^k \frac{1}{1 + \sigma j^{\frac{-1}{3}} D(j)^{-1}}.$$

Using the inversion formula

$$f(t) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{st} \hat{f}(s) ds = \frac{1}{2\pi i} \int_{s_1+i\infty}^{s_1-i\infty} e^{st} \hat{f}(s) ds,$$

we find  $r_k$  as a function of the inverse transforms  $R_k$  of  $\hat{R}_k$ :

$$r_k(s) = \frac{2D(1)}{k^{\frac{1}{3}}D(k)} \int_0^s R_k(s-s') c(s') ds', \quad k \geq 2,$$

with

$$R_k(t) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{st} \hat{R}_k(s) ds = \frac{1}{2\pi i} \int_{s_1+i\infty}^{s_1-i\infty} e^{st} \hat{f}(s) ds = \lim_{L \rightarrow \infty} \frac{1}{2\pi} \int_{-L}^L e^{its} \hat{R}_k(is) ds,$$

where  $\mathcal{C}$  is an inversion contour. A classical choice for inversion paths are Bromwich contours  $s_1 - is$ , parallel to the imaginary axis and located to the right of the singularities of  $\hat{R}_k(s)$ . In this case, we may select the imaginary axis  $s_1 = 0$ . For numerical purposes, the best choices of the inversion contour are those along which this inversion formula can be approximated by a quadrature formula involving a few points. We may resort instead to deformations of Bromwich contours, such as Talbot paths or hyperbolic paths.

## References

- [1] A Carpio Rodriguez, M Comte, R Lewandoski, A nonexistence result for a nonlinear equation involving critical Sobolev exponent, *Annales de l'Institut Henri Poincaré - Analyse Non linéaire*, 9(3), 243-261, 1992
- [2] A. Carpio, Sharp estimates of the energy for the solutions of some dissipative second order evolution equations, *Potential Analysis* 1(3), 265-289, 1992
- [3] A. Carpio, Existence de solutions globales rétrogrades pour des équations des ondes non linéaires dissipatives, *Comptes rendus de l'Académie des sciences. Série 1, Mathématique*, 316(8), 803-808, 1993
- [4] A. Carpio, Comportement asymptotique des solutions des équations du tourbillon en dimensions 2 et 3, *Comptes rendus de l'Académie des sciences. Série 1, Mathématique*, 316(12), 1289-1294, 1993
- [5] A. Carpio, Asymptotic behavior for the vorticity equations in dimensions two and three, *Communications in partial differential equations* 19 (5-6), 827-872, 1994



- [6] A. Carpio, Unicité et comportement asymptotique pour des équations de convection-diffusion scalaires, *Comptes rendus de l'Académie des sciences. Série 1, Mathématique* 319 (1), 51-56, 1994
- [7] A. Carpio, Comportement asymptotique dans les équations de Navier-Stokes, *Comptes rendus de l'Académie des sciences. Série 1, Mathématique* 319 (3), 223-228, 1994
- [8] A. Carpio, Existence of global-solutions to some nonlinear dissipative wave-equations, *Journal de mathématiques pures et appliquées* 73 (5), 471-488, 1994
- [9] A. Carpio, Large time behaviour in convection-diffusion equations, *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze* 23 (3), 551-574, 1996
- [10] A. Carpio, Large-time behavior in incompressible Navier-Stokes equations, *SIAM Journal on Mathematical Analysis* 27 (2), 449-475, 1996
- [11] A Carpio, SJ Chapman, SD Howison, JR Ockendon, Dynamics of line singularities, *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 355(1731), 2013-2024, 1997
- [12] A. Carpio, Long-time behaviour for solutions of the Vlasov-Poisson-Fokker-Planck equation, *Mathematical methods in the applied sciences* 21 (11), 985-1014, 1998
- [13] A Carpio, SJ Chapman, On the modelling of instabilities in dislocation interactions, *Philosophical Magazine B* 78 (2), 155-157, 1998
- [14] A Carpio, SJ Chapman, S Hastings, JB McLeod, Wave solutions for a discrete reaction-diffusion equation, *European Journal of Applied Mathematics* 11 (4), 399-412, 2000
- [15] A Carpio, LL Bonilla, A Wacker, E Schöll, Wave fronts may move upstream in semiconductor superlattices, *Physical Review E* 61 (5), 4866, 2000
- [16] A Carpio, P Hernando, M Kindelan, Numerical study of hyperbolic equations with integral constraints arising in semiconductor theory, *SIAM Journal on Numerical Analysis* 39 (1), 168-191, 2001
- [17] A Carpio, SJ Chapman, JLL Velázquez, Pile-up solutions for some systems of conservation laws modelling dislocation interaction in crystals, *SIAM Journal on Applied Mathematics* 61 (6), 2168-2199, 2001
- [18] A Carpio, LL Bonilla, Wave front depinning transition in discrete one-dimensional reaction-diffusion systems, *Physical Review Letters* 86 (26), 6034, 2001

- [19] G Duro, A Carpio, Asymptotic profiles for convection–diffusion equations with variable diffusion, *Nonlinear Analysis: Theory, Methods & Applications* 45 (4), 407-433, 2001
- [20] A Carpio, LL Bonilla, G Dell’Acqua, Motion of wave fronts in semiconductor superlattices, *Physical Review E* 64 (3), 036204, 2001
- [21] A Carpio, E Cebrian, FJ Mustieles, Long time asymptotics for the semiconductor Vlasov-Poisson-Boltzmann equations, *Mathematical Models and Methods in Applied Sciences* 11 (09), 1631-1655, 2001
- [22] A Carpio, LL Bonilla, A Luzón, Effects of disorder on the wave front depinning transition in spatially discrete systems, *Physical Review E* 65 (3), 035207, 2002
- [23] A Carpio, Wavefronts for discrete two-dimensional nonlinear diffusion equations, *Applied Mathematics Letters* 15 (4), 415-421, 2002
- [24] A Carpio, LL Bonilla, Depinning transitions in discrete reaction-diffusion equations, *SIAM Journal on Applied Mathematics* 63 (3), 1056-1082, 2003
- [25] A Carpio, LL Bonilla, Pulse propagation in discrete systems of coupled excitable cells, *SIAM Journal on Applied Mathematics* 63 (2), 619-635, 2003
- [26] A Carpio, LL Bonilla, Edge dislocations in crystal structures considered as traveling waves in discrete models, *Physical Review Letters* 90 (13), 135502, 2003
- [27] A Carpio, LL Bonilla, Oscillatory wave fronts in chains of coupled nonlinear oscillators, *Physical Review E* 67 (5), 056621, 2003
- [28] A Carpio, Nonlinear stability of oscillatory wave fronts in chains of coupled oscillators, *Physical Review E* 69 (4), 046601, 2004
- [29] A Carpio, G Duro, Instability and collapse in discrete wave equations, *Computational Methods in Applied Mathematics* 5 (3), 223-241, 2005
- [30] JC Neu, LL Bonilla, A Carpio, Igniting homogeneous nucleation, *Physical Review E* 71 (2), 021601, 2005
- [31] A Carpio, LL Bonilla, Discrete models of dislocations and their motion in cubic crystals, *Physical Review B* 71 (13), 134105, 2005
- [32] A Carpio, Asymptotic construction of pulses in the discrete Hodgkin-Huxley model for myelinated nerves, *Physical Review E* 72 (1), 011905, 2005
- [33] A Carpio, Wave trains, self-oscillations and synchronization in discrete media, *Physica D: Nonlinear Phenomena* 207 (1-2), 117-136, 2005
- [34] LL Bonilla, A Carpio, JC Neu, WG Wolfer, Kinetics of helium bubble formation in nuclear materials, *Physica D: Nonlinear Phenomena* 222 (1-2), 131-140, 2006

- [35] LL Bonilla, A Carpio, I Plans, Dislocations in cubic crystals described by discrete models, *Physica A: Statistical Mechanics and its Applications* 376, 361-377, 2007
- [36] I Plans, A Carpio, LL Bonilla, Homogeneous nucleation of dislocations as bifurcations in a periodized discrete elasticity model, *EPL (Europhysics Letters)* 81 (3), 36001, 2008
- [37] A Carpio, ML Rapún, Topological derivatives for shape reconstruction, In: Bonilla, L.L. (eds) *Inverse Problems and Imaging. Lecture Notes in Mathematics*, vol 1943. Springer, Berlin, Heidelberg, 2008
- [38] A Carpio, LL Bonilla, F de Juan, MAH Vozmediano, Dislocations in graphene, *New Journal of Physics* 10 (5), 053021, 2008
- [39] A Carpio, ML Rapún, Solving inhomogeneous inverse problems by topological derivative methods, *Inverse Problems* 24 (4), 045014, 2008
- [40] A Carpio, LL Bonilla, Periodized discrete elasticity models for defects in graphene, *Physical Review B* 78 (8), 085406, 2008
- [41] A Carpio, ML Rapún, Domain reconstruction using photothermal techniques, *Journal of Computational Physics* 227 (17), 8083-8106, 2008
- [42] JC Neu, A Carpio, LL Bonilla, Theory of surface deposition from boundary layers containing condensable vapour and particles, *Journal of fluid mechanics* 626, 183-210, 2009
- [43] E Cebrián, LL Bonilla, A Carpio, Self-sustained current oscillations in the kinetic theory of semiconductor superlattices, *Journal of Computational Physics* 228 (20), 7689-7705, 2009
- [44] A Carpio, G Duro, Explosive behavior in spatially discrete reaction-diffusion systems, *Discrete Contin. Dyn. Syst.* 12, 693-711, 2009
- [45] I Plans, A Carpio, LL Bonilla, Toy nanoindentation model and incipient plasticity, *Chaos, Solitons & Fractals* 42 (3), 1623-1630, 2009
- [46] A Carpio, ML Rapún, An iterative method for parameter identification and shape reconstruction, *Inverse Problems in Science and Engineering* 18 (1), 35-50, 2010
- [47] A Carpio, BT Johansson, ML Rapún, Determining planar multiple sound-soft obstacles from scattered acoustic fields, *Journal of Mathematical Imaging and Vision* 36 (2), 185-199, 2010
- [48] A Carpio, B Tapiador, Nonreflecting boundary conditions for discrete waves, *Journal of Computational Physics* 229 (5), 1879-1896, 2010

- [49] A Prados, LL Bonilla, A Carpio, Phase transitions in a mechanical system coupled to Glauber spins, *Journal of Statistical Mechanics: Theory and Experiment* P06016, 2010
- [50] LL Bonilla, A Prados, A Carpio, Nonequilibrium dynamics of a fast oscillator coupled to Glauber spins, *Journal of Statistical Mechanics: Theory and Experiment* P09019, 2010
- [51] A Carpio, B Tapiador, Analysis of helium bubble growth in radioactive waste, *Nonlinear Analysis: Real World Applications* 11 (5), 4174-4184, 2010
- [52] LL Bonilla, A Carpio, Theory of defect dynamics in graphene: defect groupings and their stability, *Continuum Mechanics and Thermodynamics* 23 (4), 337-346, 2011
- [53] A Carpio, I Peral, Propagation failure along myelinated nerves, *Journal of nonlinear science* 21 (4), 499-520, 2011
- [54] LL Bonilla, A Carpio, A Prados, RR Rosales, Ripples in a string coupled to Glauber spins, *Physical Review E* 85 (3), 031125, 2012
- [55] LL Bonilla, A Carpio, Driving dislocations in graphene, *Science* 337 (6091), 161-162, 2012
- [56] A Prados, A Carpio, LL Bonilla, Spin-oscillator model for the unzipping of biomolecules by mechanical force, *Physical Review E* 86 (2), 021919, 2012
- [57] A Carpio, ML Rapún, Hybrid topological derivative and gradient-based methods for electrical impedance tomography, *Inverse Problems* 28 (9), 095010, 2012
- [58] LL Bonilla, A Carpio, Ripples in a graphene membrane coupled to Glauber spins *Journal of Statistical Mechanics: Theory and Experiment* P09015, 2012
- [59] LL Bonilla, A Carpio, Model of ripples in graphene, *Physical Review B* 86 (19), 195402, 2012
- [60] D Rodriguez, B Einarsson, A Carpio, Biofilm growth on rugose surfaces, *Physical Review E* 86 (6), 061914, 2012
- [61] A Prados, A Carpio, LL Bonilla, Sawtooth patterns in force-extension curves of biomolecules: An equilibrium-statistical-mechanics theory, *Physical Review E* 88 (1), 012704, 2013
- [62] A Carpio, ML Rapún, Hybrid topological derivative-gradient based methods for nondestructive testing, *Abstract and applied analysis* 2013, 816134
- [63] A Carpio, ML Rapún, Parameter identification in photothermal imaging, *Journal of mathematical imaging and vision* 49 (2), 273-288, 2014

- [64] LL Bonilla, A Carpio, A Prados, Protein unfolding and refolding as transitions through virtual states, *EPL (Europhysics Letters)* 108 (2), 28002, 2014
- [65] DR Espeso, A Carpio, B Einarsson, Differential growth of wrinkled biofilms, *Physical Review E* 91 (2), 022710, 2015
- [66] LL Bonilla, A Carpio, A Prados, Theory of force-extension curves for modular proteins and DNA hairpins, *Physical Review E* 91 (5), 052712, 2015
- [67] LL Bonilla, A Carpio, C Gong, JH Warner, Measuring strain and rotation fields at the dislocation core in graphene, *Physical Review B* 92 (15), 155417, 2015
- [68] DR Espeso, A Carpio, E Martínez-García, V De Lorenzo, Stenosis triggers spread of helical *Pseudomonas* biofilms in cylindrical flow systems, *Scientific reports* 6, 27170, 2016
- [69] A Carpio, TG Dimiduk, ML Rapún, V Selgás, Noninvasive imaging of three-dimensional micro and nanostructures by topological methods, *SIAM Journal on Imaging Sciences* 9 (3), 1324-1354, 2016
- [70] A Carpio, G Duro, Well posedness of an angiogenesis related integrodifferential diffusion model, *Applied Mathematical Modelling* 40 (9-10), 5560-5575, 2016
- [71] A Carpio, G Duro, Well posedness of an integrodifferential kinetic model of Fokker–Planck type for angiogenesis, *Nonlinear Analysis: Real World Applications* 30, 184-212, 2016
- [72] A Carpio, G Duro, M Negreanu, Constructing solutions for a kinetic model of angiogenesis in annular domains, *Applied Mathematical Modelling* 45, 303-322, 2017
- [73] B Birnir, A Carpio, E Cebrián, P Vidal, Dynamic energy budget approach to evaluate antibiotic effects on biofilms, *Communications in Nonlinear Science and Numerical Simulation* 54, 70-83, 2018
- [74] A Carpio, TG Dimiduk, V Selgás, P Vidal, Optimization methods for inline holography, *SIAM Journal on Imaging Sciences* 11, 923-956, 2018
- [75] LL Bonilla, A Carpio, M Carretero, G Duro, M Negreanu, F Terragni, A convergent numerical scheme for integrodifferential kinetic models of angiogenesis, *Journal of Computational Physics* 375, 1270-1294, 2018
- [76] A Carpio, E Cebrián, P Vidal, Biofilms as poroelastic materials, *International Journal of Non-Linear Mechanics* 109, 1-8, 2019
- [77] A Carpio, TG Dimiduk, F Le Louër, ML Rapún, When topological derivatives met regularized Gauss-Newton iterations in holographic 3D imaging, *Journal of Computational Physics* 388, 224-251, 2019

- [78] A Carpio, E Cebrián, Incorporating Cellular Stochasticity in Solid-Fluid Mixture Biofilm Models, *Entropy* 22 (2), 188, 2020
- [79] A Carpio, S Iakunin, G Stadler, Bayesian approach to inverse scattering with topological priors, *Inverse Problems* 36 (10), 105001, 2020
- [80] LL Bonilla, A Carpio, C Trenado, Tracking collective cell motion by topological data analysis, *PLOS Computational Biology* 16 (12), e1008407, 2020