



On the principle of pseudo-linearized stability: Applications to some delayed nonlinear parabolic equations[☆]

A.C. Casal^{a,*}, J.I. Díaz^b

^a*Depto. de Matemática Aplicada, ETS Arquitectura, Univ. Politécnica de Madrid, 28040 Madrid, Spain*

^b*Depto. de Matemática Aplicada, Fac. de Matemáticas, Univ. Complutense de Madrid, 28040 Madrid, Spain*

1. Introduction

We study the stabilization, as $t \rightarrow \infty$, of the solutions of the nonlinear abstract functional differential equation

$$\begin{cases} \frac{du}{dt}(t) + Au(t) + Bu(t) \ni F(u_t(\cdot)) & \text{in } X, \\ u(s) = u_0(s) & s \in [-\tau, 0], \end{cases} \quad (1)$$

on a Banach space X , where

$$u_t(\theta) = u(t + \theta), \theta \in [-\tau, 0],$$

to the associated equilibria: $w \in D(A) \subset D(B) \subset X$ such that

$$Aw + Bw \ni F(\widehat{w}(\cdot)),$$

where $\widehat{w} \in C := C([-\tau, 0] : X)$ is the function which takes constant values equal to w . Our main goal is to extend, to a broad class of nonlinear operators A , the usual linearized stability principle saying, roughly speaking, that for the special case of A linear (single valued) and B and F are differentiable, the asymptotic stability of the zero solution of the

[☆] The research of both authors was partially supported by project REN2003-0223-C03 of the DGISGPI (Spain). J.I. Díaz is member of the team of the RTN HPRN-CT-2002-00274 of the EC.

* Corresponding author.

E-mail address: acasal@aq.upm.es (A.C. Casal).

linearized equation,

$$\begin{cases} \frac{dv}{dt}(t) + Av(t) + DB(w)v(t) = DF(\widehat{w})v_t(\cdot) & \text{in } X, \\ v(s) = u_0(s) & s \in [-\tau, 0], \end{cases}$$

implies that $u(t : u_0) \rightarrow w$ as $t \rightarrow \infty$, at least if $u_0(\cdot)$ is close enough to \widehat{w} . We point out that our results seem to be new even without the delayed and nonlocal term (i.e. for $F \equiv 0$).

Our main motivation comes from some previous works by the authors and collaborators [11,12,15] dealing with the stabilization of the uniform oscillations for the *complex Ginzburg–Landau equation*. This stabilization takes place by means of some global delayed feedback. If, for instance, we consider the case in which the domain is $\Omega = (0, L_1) \times (0, L_2)$ with periodic boundary conditions, and define the faces of the boundary

$$\Gamma_j = \partial\Omega \cap \{x_j = 0\}, \Gamma_{j+2} = \partial\Omega \cap \{x_j = L_j\}, \quad j = 1, 2,$$

this problem can be stated as follows:

$$(P_1) \quad \begin{cases} \frac{\partial \mathbf{u}}{\partial t} - (1 + i\varepsilon)\Delta \mathbf{u} = (1 - i\omega)\mathbf{u} \\ \quad - (1 + i\beta)|\mathbf{u}|^2\mathbf{u} + \mu e^{i\chi_0}\mathbf{F}(\mathbf{u}, t, \tau) & \Omega \times (0, +\infty), \\ \mathbf{u}|_{\Gamma_j} = \mathbf{u}|_{\Gamma_{j+2}}, \left(-\frac{\partial \mathbf{u}}{\partial \mathbf{n}} \Big|_{\Gamma_j} = \right) \frac{\partial \mathbf{u}}{\partial x_j} \Big|_{\Gamma_j} \\ \quad = \frac{\partial \mathbf{u}}{\partial x_j} \Big|_{\Gamma_{j+2}} \left(= \frac{\partial \mathbf{u}}{\partial \mathbf{n}} \Big|_{\Gamma_{j+2}} \right) & \partial\Omega \times (0, +\infty), \\ \mathbf{u}(x, s) = \mathbf{u}_0(x, s) & \Omega \times [-\tau, 0], \end{cases}$$

where \mathbf{n} is the outpointing normal unit vector and

$$\mathbf{F}(\mathbf{u}, t, \tau) = [m_1\mathbf{u}(t) + m_2\bar{\mathbf{u}}(t) + m_3\mathbf{u}(t - \tau, x) + m_4\bar{\mathbf{u}}(t - \tau)]$$

$$\text{with } \bar{\mathbf{u}}(s) = (1/|\Omega|) \int_{\Omega} \mathbf{u}(s, x) dx.$$

Here the parameters $\varepsilon, \beta, \omega, \mu, \chi_0, m_i$ and τ are real numbers, in contrast with the solution $\mathbf{u}(x, t) = u_1(x, t) + iu_2(x, t)$.

This type of equations (called as of Stuart–Landau in absence of the diffusion term) arise in the study of the stability of reaction diffusion equations such as $(\partial \mathbf{x} / \partial t) - \mathbf{D}\Delta \mathbf{X} = \mathbf{f}(\mathbf{X} : \eta)$ where $\mathbf{X} : \Omega \times (0, +\infty) \rightarrow \mathbb{R}^n$ and η is a real scalar parameter when the deviation \mathbf{v} from the uniform state solution \mathbf{X}_∞ is developed asymptotically in terms of some multiple scales (see [17]). Coefficient ε measures the degree to which the diffusion matrix \mathbf{D} deviates from a scalar. With the basis of a sound experimental work, many recent studies of a more descriptive nature, but of a great originality and interest, have been written. In those studies the delay term $\mathbf{F}(\mathbf{u}, t, \tau)$ has been taken corresponding to $m_4 = 1, m_i = 0$ for $i = 1, 2, 3$ and introduced as a control mechanism (see [6,18]).

If we focus our attention on the so-called *slowly varying complex amplitudes* defined by $\mathbf{u}(x, t) = \mathbf{v}(x, t)e^{-i\omega t}$, thus, \mathbf{v} satisfies

$$(P_2) \begin{cases} \frac{\partial \mathbf{v}}{\partial t} - (1 + i\varepsilon)\Delta \mathbf{v} = \mathbf{v} - (1 + i\beta)|\mathbf{v}|^2 \mathbf{v} \\ \quad + \mu e^{i\chi_0} [m_1 \mathbf{v} + m_2 \bar{\mathbf{v}} \\ \quad + e^{i\omega\tau} (m_3 \mathbf{v}(t - \tau, x) + m_4 \bar{\mathbf{v}}(t - \tau))] \quad \text{in } \Omega \times (0, +\infty), \\ \mathbf{v}|_{\Gamma_j} = \mathbf{v}|_{\Gamma_{j+2}}, \left(-\frac{\partial \mathbf{v}}{\partial \mathbf{n}} \Big|_{\Gamma_j} = \right) \frac{\partial \mathbf{v}}{\partial x_j} \Big|_{\Gamma_j} \\ = \frac{\partial \mathbf{v}}{\partial x_j} \Big|_{\Gamma_{j+2}} \left(= \frac{\partial \mathbf{v}}{\partial \mathbf{n}} \Big|_{\Gamma_{j+2}} \right) \quad \text{on } \partial\Omega \times (0, +\infty), \\ \mathbf{v}(x, s) = \mathbf{u}_0(x, s)e^{i\omega s} \quad \text{on } \Omega \times [-\tau, 0]. \end{cases}$$

The existence and uniqueness of a solution of (P₁) can be proven once we assume, for instance, that $\mathbf{u}_0 \in C([-\tau, 0] : \mathbf{L}^2(\Omega))$ (see [15]). In the mentioned references we were interested in the stability analysis of the time-periodical function $\mathbf{v}_{uosc}(x, t) = \rho_0 e^{-i\theta t}$. We can reduce the study to the stability of stationary solutions of some auxiliary problem by introducing the change of unknown $\mathbf{z}(x, t) = \mathbf{v}(x, t)e^{i\theta t}$ where $\mathbf{v}(x, t)$ is a solution of (P₂). Thus $\mathbf{z}(x, t)$ satisfies

$$(P_3) \begin{cases} \frac{\partial \mathbf{z}}{\partial t} - (1 + i\varepsilon)\Delta \mathbf{z} = (1 + i\theta)\mathbf{z} - (1 + i\beta)|\mathbf{z}|^2 \mathbf{z} \\ \quad + \mu e^{i\chi_0} [m_1 \mathbf{z} + m_2 \bar{\mathbf{z}} \\ \quad + e^{i(\omega+\theta)\tau} (m_3 \mathbf{z}(t - \tau, x) + m_4 \bar{\mathbf{z}}(t - \tau))] \quad \text{in } \Omega \times (0, +\infty), \\ \mathbf{z}|_{\Gamma_j} = \mathbf{z}|_{\Gamma_{j+2}}, \left(-\frac{\partial \mathbf{z}}{\partial \vec{n}} \Big|_{\Gamma_j} = \right) \frac{\partial \mathbf{z}}{\partial x_j} \Big|_{\Gamma_j} \\ = \frac{\partial \mathbf{z}}{\partial x_j} \Big|_{\Gamma_{j+2}} \left(= \frac{\partial \mathbf{z}}{\partial \vec{n}} \Big|_{\Gamma_{j+2}} \right) \quad \text{on } \partial\Omega \times (0, +\infty), \\ \mathbf{z}(x, s) = \mathbf{u}_0(x, s)e^{i(\omega-\theta)s} \quad \text{on } \Omega \times [-\tau, 0]. \end{cases}$$

Notice that now, $\mathbf{v}_{uosc}(x, t) = \rho_0 e^{-i\theta t}$ is an uniform oscillation if and only if $\mathbf{z}(x, t) = \mathbf{v}_{uosc}(x, t)e^{i\theta t} = \mathbf{y} = \rho_0$ is a stationary solution of (P₃): i.e. $\mathbf{0} = (1 + i\theta)\mathbf{y} - (1 + i\beta)|\mathbf{y}|^2 \mathbf{y} + \mu e^{i\chi_0} [m_1 + m_2 + e^{i(\omega+\theta)\tau} (m_3 + m_4)]\mathbf{y}$.

The motivation to keep A nonlinear after the process of linearization (reason why we used the term of *pseudo-linearization principle*) comes from the fact that if we use the representation for the unknown of the delayed nonlinear equation (P₃) as $\mathbf{z}(x, t) = \rho(x, t)e^{i\phi(x, t)}$ then we arrive to a coupled nonlinear system of delayed equations for ρ and ϕ which can be described in terms of the representation operator given by $\mathbf{P} : \mathbb{R}^2 \rightarrow \mathbb{C}$, $\mathbf{P}(\rho, \phi) = \rho e^{i\phi}$. Indeed, notice that \mathbf{P} is nonlinear and that if $\mathbf{q} = (\rho, \phi)$ then $\mathbf{z}(x, t) = \mathbf{P}(\mathbf{q}(x, t))$ and the (P₃) can be formulated as $d\mathbf{P}(\mathbf{q}(\cdot, t))/dt + A\mathbf{P}(\mathbf{q}(\cdot, t)) + B\mathbf{P}(\mathbf{q}(\cdot, t)) = F(\mathbf{P}(\mathbf{q}(\cdot, t)))$. By using that the matrix $\mathbf{C}(\mathbf{q}(\cdot, t)) = \text{grad } \mathbf{P}(\mathbf{q}(\cdot, t))$ is not singular, we can arrive to the simpler

formulation

$$\frac{d\mathbf{q}}{dt}(\cdot, t) + \mathbf{C}(\mathbf{q}(\cdot, t))^{-1}[\mathbf{A}\mathbf{P}(\mathbf{q}(\cdot, t)) + \mathbf{B}\mathbf{P}(q(\cdot, t))] = \mathbf{C}(\mathbf{q}(\cdot, t))^{-1}F(\mathbf{P}(\mathbf{q}(\cdot, t))). \tag{2}$$

Notice that, although this delayed system can be also (formally) linearized (this is the procedure followed in [6,18]) the above diffusion operator $\mathbf{C}(\mathbf{q}(\cdot, t))^{-1}\mathbf{A}\mathbf{P}(\mathbf{q}(\cdot, t))$ becomes now quasilinear on \mathbf{q} and thus the mathematical justification is much more delicate.

Other examples, given in Section 3, justify also the philosophy of keeping A nonlinear after linearizing the rest of the terms of the equation. For instance, this is the case when A is multivalued, or nondifferentiable or a degenerate quasilinear operator. We point out that some relevant examples of nonlinear functional equations arise in the most different contexts (see, for instance, [14] for one example in Climatology, [13] for a family of examples dealing with the wealth of nations and the general exposition made in [16]).

Coming back to the abstract formulation, the structural assumptions we shall assume in this paper are the following

(H1): $A \in \mathcal{A}(\omega : X)$, for some $\omega \in \mathbb{C}$, with

$$\mathcal{A}(\omega : X) = \{A : D_X(A) \subset X \rightarrow \mathcal{P}(X) \text{ such that } A + \omega I \text{ is a } m\text{-accretive operator}\},$$

(see [9] for the case of $X = H$ a Hilbert space and the works by Benilan, Crandall, Pazy and others for the case of a general Banach space: see the monographs [8,21]),

(H2): The operators semigroup $T(t) : \overline{D_X(A)}^X \rightarrow X, t \geq 0$, generated by A , is compact (see [21]),

(H3): $B \in \mathcal{A}(0 : X)$, B is single valued, Fréchet differentiable, and B is dominated by A ; i.e.

$$D_X(A) \subset D_X(B) \quad \text{and} \quad |Bu| \leq k|A^0u| + \sigma(|u|) \tag{3}$$

for any $u \in D_X(A)$ and for some $k < 1$ and some continuous function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$, where, here and in what follows, $|\cdot|$ denotes the norm in the space X (in contrast with the norm in space C which will be denoted by $\|\cdot\|$ if there is no ambiguity, when handling two spaces X and Y the corresponding norms will be indicated), $|A^0u| := \inf\{|\xi| : \xi \in Au\}$ for $u \in D_X(A)$,

(H4): $F : C \rightarrow X$ satisfies a local Lipschitz condition, i.e., for any $R > 0$ there exists $L(R) > 0$ such that

$$|F(\phi) - F(\psi)| \leq L(R)\|\phi - \psi\| \quad \text{for any } \phi, \psi \in C \text{ and } \|\phi\|, \|\psi\| \leq R. \tag{4}$$

(H5): There exists $\delta^F > 0$ such that $F : B_{\delta^F}^X(\widehat{w}) \rightarrow X$ is Fréchet differentiable with the Fréchet derivative $DF(\widehat{w})$ given by $D(F(\widehat{w}))\phi = \int_{-\tau}^0 d\eta(\theta)\phi(\theta), \phi \in C$, for $\eta : [-\tau, 0] \rightarrow B(X, X)$ of bounded variation and the Fréchet derivative is locally Lipschitz continuous, where $B_{\delta^F}^X(\widehat{w}) = \{\phi \in C; \|\phi - \widehat{x}\| < \delta^F\}$,

We further assume the main condition of our arguments:

(H6): The operator $y \rightarrow Ay + By - DF(\widehat{w})(e^{\omega \cdot} y)$ belongs to $\mathcal{A}(\omega : X)$, for some $\omega \in \mathbb{C}$ with $\text{Re } \omega = \gamma < 0$ where $e^{\omega \cdot} v \in C$ is defined by

$$(e^{\omega \cdot} v)(s) = e^{\omega s} \widehat{v}(s), \quad \text{with } \widehat{v}(s) = v \quad \text{for any } s \in [-\tau, 0] \text{ for } v \in X. \tag{5}$$

In order to treat the case in which B is differentiable we introduce the conditions

(H7): There exists a Banach space Y and there exists $\delta^B > 0$ such that B is Fréchet differentiable as function from $B_{\delta^B}(w) = \{z \in D(B); |w - z| < \delta^B\}$ into Y , with the Fréchet derivative $DB(w)$ locally Lipschitz continuous,

(H8): The operator $y \rightarrow Ay + DB(w)y - DF(\widehat{w})(e^{\omega^* \cdot} y)$ belongs to $\mathcal{A}(\omega^* : Y)$, for some $\omega^* \in \mathbb{C}$ with $\text{Re } \omega^* = \gamma^* < 0$.

2. The abstract results

Theorem 1. Assume (H1)–(H6). Then there exists $\alpha > 0$, $\varepsilon > 0$ and $M \geq 1$ such that if $u_0 \in B_\varepsilon^X(\widehat{w})$, $u_0(s) \in D_X(B)$ for any $s \in [-\tau, 0]$ then the solution $u(\cdot : u_0)$ of (1) exists on $[-\tau, +\infty)$ and

$$|u(t : u_0) - w| \leq M e^{-\alpha t} \|u_0 - \widehat{w}\| \quad \text{for any } t > 0. \tag{6}$$

Moreover, if we also assume (H7), that (H1)–(H5) holds on the space Y and (H8) then there exists $\alpha^* > 0$, $\varepsilon^* \in (0, \varepsilon)$ and $M^* \geq 1$ such that if $u_0 \in B_{\varepsilon^*}^{X \cap Y}(\widehat{w})$, $u_0(s) \in D_X(B) \cap D_Y(B)$ for any $s \in [-\tau, 0]$ then

$$\begin{aligned} &|u(t : u_0) - w|_X + |u(t : u_0) - w|_Y \\ &\leq M^* e^{-\alpha^* t} (\|u_0 - \widehat{w}\|_X + \|u_0 - \widehat{w}\|_Y) \quad \text{for any } t > 0. \end{aligned} \tag{7}$$

Proof. From assumptions (H4) and (H5)

$$F(\phi) = F(\widehat{w}) + DF(\widehat{w})(\phi - \widehat{w}) + G^F(\widehat{w}, \phi) \quad \text{for any } \phi \in B_{\delta^F}^X(\widehat{w}).$$

Moreover since $DF(\widehat{w})$ is locally Lipschitz continuous, there exists a continuous increasing functions b_X^F such that

$$|G^F(\widehat{w}, \phi)| \leq b_X^F(\|\phi - \widehat{w}\|) \|\phi - \widehat{w}\| \quad \text{for any } \phi \in B_{\delta^F}^X(\widehat{w}). \tag{8}$$

Then

$$\frac{du}{dt}(t) - \frac{dw}{dt} + Au(t) - Aw + Bu(t) - Bw - DF(\widehat{w})(u_t - \widehat{w}) \ni -G^F(\widehat{w}, u_t). \tag{9}$$

We now use assumption (H6). We claim that we can find a constant $K \geq 1$ and such that

$$\|u_t - \widehat{w}\| \leq K e^{\gamma t} \|u_0 - \widehat{w}\| + \int_0^t K e^{\gamma(t-s)} |G^F(\widehat{w}, u_s)| ds. \tag{10}$$

Indeed, as $u(t)$ and w are “integral solutions” in the sense of Benilan (see, e.g. [8]), then, by (H6), if we multiply (9) by $u(t) - w$ (by using the usual semiinner braket $[\cdot, \cdot]$; see, for instance [8] or [21, Section 1.4]) we get that

$$|u(t) - w| \leq K e^{\gamma(t-t_0)} |u(t_0) - w| + \int_{t_0}^t K e^{\gamma(t-s)} |G^F(\widehat{w}, u_s)| ds \tag{11}$$

for any $t \geq t_0 \geq 0$ (see, for instance, [8] or [21, Theorem 1.7.5]). Then,

$$|u(t) - w| \leq Ke^{\gamma t} \|u_0 - \widehat{w}\| + \int_0^t Ke^{\gamma(t-s)} |G^F(\widehat{w}, u_s)| ds \tag{12}$$

for any $t \geq 0$. Finally, since (12) holds trivially for $t \in [-\tau, 0]$ we get (10) by taking the maximum, in (11), on intervals of the form $[t - \tau, t]$ for any $t \geq 0$.

Now, let $R \in (0, \delta^F)$ be chosen so that

$$b_X^F(R) < (-\gamma)/(4K). \tag{13}$$

Define $\varepsilon = \min\{R/(2K), \delta_X^F\}$. Let us show that if $u_0 \in B_\varepsilon^X(\widehat{w})$ then the associated solution u of (1) exists and $\|u_t - \widehat{w}\| < R$ for all $t \geq 0$. Thanks to assumption (H2) we can apply some maximal continuation results (see, for instance, Chapter 3 of [21], or Chapter 2 of [22] when A is linear), it suffices to show that there exists no $t_1 > 0$ so that $\|u_{t_1}\| = R$ and $\|u_t\| < R$ for $t \in [0, t_1)$. By contradiction, if there exists such a t_1 , then on $[0, t_1]$ we have

$$\begin{aligned} \|u_t - \widehat{w}\| &\leq Ke^{\gamma t} \|u_0 - \widehat{w}\| + \int_0^t Ke^{\gamma(t-s)} |G^F(\widehat{w}, u_s)| ds \\ &\leq Ke^{\gamma t} \|u_0 - \widehat{w}\| + 2Kb_X^F(R) \int_0^t e^{\gamma(t-s)} \|u_s - \widehat{w}\| ds. \end{aligned}$$

In particular, at $t = t_1$ we have

$$\|u_{t_1} - \widehat{w}\| \leq K\varepsilon + \frac{2Kb_X^F(R)}{(-\gamma)} R \leq R,$$

a contradiction to the choice of t_1 .

Finally, to end the proof, let $u_0 \in B_\varepsilon^X(\widehat{w})$, $u_0(s) \in D_X(B)$ for any $s \in [-\tau, 0]$ and let u the associated solution of (1). Since we have shown that $\|u_t - \widehat{w}\| \leq R$ for all $t \geq 0$ we get that

$$\|u_t - \widehat{w}\| \leq Ke^{\gamma t} \|u_0 - \widehat{w}\| + Kb_X^F(R) \int_0^t e^{\gamma(t-s)} \|u_s - \widehat{w}\| ds \tag{14}$$

holds for all $t \geq 0$. Thus, by using the Gronwall's inequality, we get

$$\|u_t - \widehat{w}\| \leq Ke^{[\gamma - Kb_X^F(R)]t} \|u_0 - \widehat{w}\| \leq Ke^{(\gamma/2)t} \|u_0 - \widehat{w}\|, u_0 \in B_\varepsilon^X(\widehat{w})$$

which shows (6).

In order to show the decay estimate (7), we repeat the same arguments as before but now on the space Y . Then, from assumptions (H3) on Y and (H7), there exist δ_Y^F and δ_X^B such that

$$\begin{aligned} B(z) &= B(w) + DB(w)(z - w) + G^B(w, z) \quad \text{for any } z \in B_{\delta_X^B}(w), \\ F(\phi) &= F(\widehat{w}) + DF(\widehat{w})(\phi - \widehat{w}) + G^F(\widehat{w}, \phi) \quad \text{for any } \phi \in B_{\delta_Y^F}^Y(\widehat{w}), \end{aligned}$$

where now

$$\begin{aligned} B_{\delta_X^B}(w) &= \{z \in D_X(B) \cap D_Y(B); |w - z| < \delta_X^B\}, \\ B_{\delta_Y^F}^Y(\widehat{w}) &= \{\phi \in C; \|\phi - \widehat{x}\|_Y < \delta_Y^F\} \end{aligned}$$

and, as before, $\|\cdot\|_Y$ denotes the norm on the space $C_Y := C([-τ, 0] : Y)$. Moreover, there exists two continuous increasing functions b_X^B and b_Y^F such that

$$|G^B(w, z)|_Y \leq b_X^B(|w - z|)|w - z| \quad \text{for any } z \in B_{\delta_X^B}(w), \tag{15}$$

$$|G^F(\widehat{w}, \phi)|_Y \leq b_Y^F(\|\phi - \widehat{w}\|_Y)\|\phi - \widehat{w}\|_Y \quad \text{for any } \phi \in B_{\delta_Y^F}(\widehat{w}). \tag{16}$$

Now

$$\begin{aligned} \frac{du}{dt}(t) - \frac{dw}{dt} + Au(t) - Aw + DB(w)(u(t) - w) - DF(\widehat{w})(u_t - \widehat{w}) \\ \ni G^B(w, u(t)) - G^F(\widehat{w}, u_t). \end{aligned} \tag{17}$$

Thus, by using (H8) and arguing as in the first part we get that there exists a constant $K^* \geq 1$ such that

$$\begin{aligned} \|u_t - \widehat{w}\|_Y \leq K^* e^{\gamma^* t} \|u_0 - \widehat{w}\|_Y + \int_0^t K^* e^{\gamma^*(t-s)} (|G^B(w, u(s))|_Y \\ + |G^F(\widehat{w}, u_s)|_Y) ds \end{aligned} \tag{18}$$

and then, by taking $\delta = \min(\delta_X^B, \delta_Y^F)$ and $R^* \in (0, \delta)$ such that

$$\max(b_X^B(R^*), b_Y^F(R^*)) < (-\gamma)/(4K), \tag{19}$$

we obtain that

$$\begin{aligned} \|u_t - \widehat{w}\|_Y \leq K^* e^{\gamma^* t} \|u_0 - \widehat{w}\|_Y \\ + K^* \int_0^t e^{\gamma^*(t-s)} (b_X^B(R^*) \|u_s - \widehat{w}\|_X + b_Y^F(R^*) \|u_s - \widehat{w}\|_Y) ds. \end{aligned} \tag{20}$$

We define $\widetilde{R} = \min(R, R^*)$, $\widetilde{K} = \max(K, K^*)$, $\widetilde{\gamma} = \max(\gamma, \gamma^*) < 0$ and $\varepsilon^* = \min\{\widetilde{R}/(2\widetilde{K}), \delta\}$. Then, if $u_0 \in B_{\varepsilon^*}^{X \cap Y}(\widehat{w})$, $u_0(s) \in D_X(B) \cap D_Y(B)$ for any $s \in [-\tau, 0]$ and we assume, for instance, that $\widetilde{\gamma} = \gamma$, by adding (14) and (20) we deduce that

$$\begin{aligned} \|u_t - \widehat{w}\|_X + \|u_t - \widehat{w}\|_Y \leq \widetilde{K} e^{\widetilde{\gamma} t} (\|u_0 - \widehat{w}\|_X + e^{(\gamma^* - \gamma)t} \|u_0 - \widehat{w}\|_Y) \\ + \widetilde{K} \int_0^t e^{\widetilde{\gamma}(t-s)} [(b_X^F(\widetilde{R}) + b_X^B(\widetilde{R}) e^{(\gamma^* - \gamma)t}) \|u_s - \widehat{w}\|_X \\ + b_Y^F(R^*) e^{(\gamma^* - \gamma)t} \|u_s - \widehat{w}\|_Y] ds. \end{aligned} \tag{21}$$

and estimate (7) follows, again, by Gronwall’s inequality. \square

Remark 2. It is not difficult to show that the assumption (H8) is implied (when A is linear) by the condition: “if $\lambda \in \mathbb{C}$ is given so that there exists $y \in D(B) \setminus \{0\}$ such that $Ay + DB(w)y - \lambda y \ni DF(\widehat{w})(e^{\lambda \cdot} y)$ then $\text{Re } \lambda > 0$ ”. This allow to see Theorem 4.1 of Wu [22] (see also [19] and its references) as an special case of our abstract result with $B = 0$. In that case the “variation of the constants formula” can be used to get a different proof of the theorem since A is linear. Notice that if $B \neq 0$ and $D(B)\overline{A}X$ then the arguments of the proof of Wu [22] do not work (in spite of the claimed in the Example 4.8 given there).

Remark 3. When A is linear, as in the case without delay, assumption (H7) implies that the zero solution of the linearized problem $(dU/dt)(t) + AU(t) + DB(w)U(t) - DF(\widehat{w})U_t(\cdot) = 0$ in X , is locally asymptotically stable [22].

Remark 4. It is possible to prove the existence of global solutions for a general class of initial data (not necessarily near \widehat{w}) by using that $A + B \in \mathcal{A}(\omega : X)$, for some $\omega \in \mathbb{C}$, some truncation of the nonlocal term $F(u_t)$ and passing to the limit by the compactness of the semigroup generated by A (see [21] for some related results).

An easy adaptation of the above proof leads to the following linearization result (now on a possibly smaller neighborhood of w) when A is differentiable.

Theorem 5. *The conclusion of the above result remains true if we assume, additionally, that condition (H7) also holds for A and we replace condition (H8) by*

(H9): The operator $y \rightarrow DA(w)y + DB(w)y - DF(\widehat{w})(e^{\omega \cdot} y)$ belongs to $\mathcal{A}(\omega)$, for some $\omega \in \mathbb{C}$ with $\text{Re } \omega = \gamma < 0$.

Remark 6. We claim that our arguments keeping A nonlinear after linearizing the rest of the terms (and in particular the way in which we apply Gronwall inequality) allow to extend, to the case of quasilinear equations, the so-called “method of quasilinearization” which, introduced by Bellman and Kalaba [7], we used to find solutions of a parabolic semilinear problem through the iteration of solutions of the linearized equation when starting in a super and a subsolution of the original semilinear problem (see, e.g., [10] and its references). This will be the subject of a future work by the authors.

3. Some examples

3.1. Example 1. The complex Ginzburg–Landau equation with a global delayed mechanism

Motivated by the special form of the nonlinear term of the equation in (P_3) we shall take $X = \mathbf{L}^4(\Omega)$ and $Y = \mathbf{L}^{4/3}(\Omega)$ (notice that, in contrast with the case of scalar equations (see [19]) the space $\mathbf{L}^\infty(\Omega)$ is not suitable space to check assumption (H1): see [5]). A detailed analysis of the associated diffusion operator is consequence of some previous results in the literature (see, for instance [3]). Notice that the operator Au can be formulated matrixially as

$$\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \rightarrow \begin{pmatrix} \Delta & -\varepsilon\Delta \\ \varepsilon\Delta & \Delta \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}.$$

So, if $\varepsilon \neq 0$ the diffusion matrix has a nonzero antisymmetric part. In particular, A is the generator of a semigroup of contractions $\{T(t)\}_{t \geq 0}$ on X and the compactness of the semigroup is consequence of the compactness of the inclusion $D(A) \subset X$ (notice that,

since $N = 2$, $\mathbf{W}^{1,4}(\Omega) \subset \mathbf{W}^{1,4/3}(\Omega) \subset \mathbf{C}(\overline{\Omega})$ with compact imbedding) and some regularity results for nonsymmetric systems.

Concerning the rest of the terms of the equation in (P_3) , we define $B\mathbf{u} = (1 + i\beta)|\mathbf{u}|^2\mathbf{u}$ with $D(B) = \mathbf{L}^{12}(\Omega)$. By using the characterization of the semiinner bracket $[\cdot, \cdot]$ for the spaces $L^p(\Omega)$ (see, for instance [8]) it is easy to see that B verifies (H3). Moreover, by the results on the Frechet differentiability of Nemitsky operators (see Theorem 2.6 (with $p = 4$) of Ambrosetti and Prodi [4]) we get that (H7) holds, with $DB(\mathbf{y})\mathbf{v} = 3(1 + i\beta)|\mathbf{y}|^2\mathbf{v}$, if we take $Y = \mathbf{L}^{4/3}(\Omega)$. It can be found in the above-mentioned reference that assumption (H7) does not hold if we take $X = Y = \mathbf{L}^2(\Omega)$.

The nonlocal term is defined, by

$$F(\mathbf{u}_t) = (1 + i\theta)\mathbf{u}(t) + \mu e^{i\lambda_0} [m_1\mathbf{u}(t) + m_2\bar{\mathbf{u}}(t) + e^{i(\omega+\theta)\tau}(m_3\mathbf{u}(t - \tau) + m_4\bar{\mathbf{u}}(t - \tau))],$$

is locally Lipschitz continuous and its Frechet derivative is given by

$$DF(\widehat{\mathbf{y}})\mathbf{v}(t) = - (1 + i\theta)\mathbf{v}(t) - \mu e^{i\lambda_0} [m_1\mathbf{v}(t) + m_2\bar{\mathbf{v}}(t) - e^{i(\omega+\theta)\tau}(m_3\mathbf{v}(t - \tau) - m_4\bar{\mathbf{v}}(t - \tau))], \tag{22}$$

since for any $\phi \in C$, the nonlocal operator $\phi \rightarrow (1/|\Omega|) \int_{\Omega} \phi(s) dx$ is linear and we can write $DF(\widehat{\mathbf{y}})\phi = \int_{-\tau}^0 d\eta(s)\phi(s)$, with

$$d\eta(s)\mathbf{v}(s) = \delta_0(s)(\mathbf{1} + i\theta)\mathbf{v}(s) + \mu e^{i\lambda_0} [\delta_0(s)(m_1\mathbf{v}(s) + m_2\bar{\mathbf{v}}(s)) + e^{i(\omega+\theta)\tau}\delta_{-\tau}(s)(m_3\mathbf{v}(s) + m_4\bar{\mathbf{v}}(s))] \tag{23}$$

for any $\mathbf{v} \in C([-\tau, \infty) : \mathbf{L}^4(\Omega))$ and any $s \in [-\tau, \infty)$, where $\delta_0(s)$, $\delta_{-\tau}(s)$ denote the Dirac delta at the points $s = 0$ and $-\tau$, respectively. By well-known results, we have that $\eta : [-\tau, 0] \rightarrow B(X, X)$ has a bounded variation and so, conditions (H4) and (H5) hold (and analogously replacing X by Y).

Finally, assumption (H6) can be read as a condition on the stationary state \mathbf{y} (a study of the eigenvalue of operator A can be found, for instance, in [20]).

Remark 7. By introducing the representation operator $\mathbf{P} : \mathbb{R}^2 \rightarrow \mathbb{C}$, $\mathbf{P}(\rho, \phi) = \rho e^{i\phi}$ it is clear that the quasilinear operator $A\mathbf{P}(\mathbf{q})$ obtained from the operator $A\mathbf{u} = -(1 + i\varepsilon)\Delta\mathbf{u}$ satisfies also condition $A \in \mathcal{A}(\omega)$ (since \mathbf{P} is merely a change of variables). We point out that,

$$A\mathbf{P}(\mathbf{q}) = -(1 + i\varepsilon)[\Delta\rho - \rho|\nabla\phi|^2 + i(2\nabla\rho \cdot \nabla\phi + \rho\Delta\phi)]e^{i\phi}.$$

Then, the “ formal linearization” of the operator $\mathbf{E}(\mathbf{q}) := A\mathbf{P}(\mathbf{q})$ at $\mathbf{q}^*(x, y) := \mathbf{y} \equiv \rho_0$ becomes

$$D\mathbf{E}(\mathbf{q}^*)(\rho e^{i\phi}) = -(1 + i\varepsilon)[\Delta\rho + i\rho_0\Delta\phi]e^{i\phi}.$$

Notice that the linearization of $\mathbf{C}(\mathbf{q})^{-1}A\mathbf{P}(\mathbf{q})$ needs a slight modification of the above linear expression.

3.2. Example 2. Case in which A is nonlinear and nondifferentiable

It is not difficult to adapt the results of the first example to the case in which the vectorial operator is given by

$$\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \rightarrow \begin{pmatrix} A_1 & -\varepsilon\Delta \\ \varepsilon\Delta & A_2 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \tag{24}$$

with $A_i : D(A_i) \rightarrow \mathcal{P}(L^4(\Omega))$ two (possibly different) m -accretive operators in $L^4(\Omega)$ as for instance,

$$\left\{ \begin{array}{l} A_i u = -\operatorname{div}(|\nabla u|^{p_i-2} \nabla u) + \beta_i(u) \\ D(A_i) = \left\{ u \in W^{1,1}(\Omega) \cap L^4(\Omega), u(x) \in D(\beta) \text{ a.e. } x \in \Omega, A_i u \in L^4(\Omega) \right. \\ \left. \text{and } -\left| \frac{\partial u}{\partial n} \right|^{p_i-2} \frac{\partial u}{\partial n} \in \gamma_i(u) \text{ on } \partial\Omega \right\}, \end{array} \right.$$

where $p_i \in (1, +\infty)$ and β_i, γ_i are maximal monotone graphs of \mathbf{R}^2 (not necessarily associated to differentiable functions). We send the reader to Vrabie [21] (and its references) for the study of the assumptions (H1) and (H2) for each of the nonlinear operators A_i . We point out that the structure of the nonlinear diffusion operator (24) allows to guarantee that the diffusion operator is m -accretive on $\mathbf{L}^{4/3}(\Omega)$. For related works see Aftalion and Pacella [1,2]. The same holds also on $\mathbf{L}^{4/3}(\Omega)$.

References

- [1] A. Aftalion, F. Pacella, Uniqueness and nondegeneracy for some nonlinear elliptic problems in a ball, preprint.
- [2] A. Aftalion, F. Pacella, Morse index and uniqueness for positive radial solutions of p -Laplace equations, preprint.
- [3] H. Amann, Dynamic theory of quasilinear parabolic equations: II. Reaction–diffusion systems, *Differential Integral Equations* 3 (1990) 13–75.
- [4] A. Ambrosetti, G. Prodi, *A Primer of Nonlinear Analysis*, Cambridge University Press, Cambridge, 1993.
- [5] P. Auscher, L. Bathélemy, Ph. Bénilan, E.M. Ouhabaz, Absence de la L^∞ -contractivité pour les semi-groupes associés aux opérateurs elliptiques complexes sous forme divergence, *Potential Anal.* 12 (2000) 169–189.
- [6] D. Battogtokh, A. Mikhailov, Controlling turbulence in the complex Ginzburg–Landau equation, *Physica D* 90 (1996) 84–95.
- [7] R. Bellman, R. Kalaba, *Quasilinearization and Nonlinear Boundary Value Problems*, American Elsevier, New York, 1965.
- [8] Ph. Benilan, M.G. Crandall, A. Pazy, *Nonlinear Evolution Equations in Banach Spaces*, in preparation.
- [9] H. Brezis, *Opérateurs Maximaux Monotones*, North-Holland, Amsterdam, 1973.
- [10] S. Carl, V. Lakshmikantham, Generalized quasilinearization and semilinear parabolic problems, *Nonlinear Anal.* 48 (2002) 947–960.
- [11] A.C. Casal, J.I. Díaz, On the stabilization of periodic solutions for the complex Ginzburg–Landau equation by a delayed feedback, Two-dimensional systems, in: S.J. Álvarez (Ed.), *Jornada matemática en honor de J.M. Fraile*, Universidad Complutense de Madrid, 2004.
- [12] A.C. Casal, J.I. Díaz, J.F. Padial, L. Tello, On the stabilization of uniform oscillations for the complex Ginzburg–Landau equation by means of a global delayed mechanism, in: J.M. Jorner, J.M. López, C. Olivé, R. Ramírez (Eds.), *CD-Rom Actas XVIII CEDYA/VIII CMA*, Servicio de Publicaciones de la Univ. de Tarragona, 2003.

- [13] E.N. Chukwu, *Differential Models and Neutral Systems for Controlling the Wealth of Nations*, World Scientific, Singapore, 2001.
- [14] J.I. Díaz, G. Hetzer, A functional quasilinear reaction–diffusion equation arising in climatology, in: J.-L. Lions (Ed.), *Équations aux dérivées partielles et applications, Articles dédiés à*, Elsevier, Paris, 1998, pp. 461–480.
- [15] J.I. Díaz, J.F. Padial, J.I. Tello, L. Tello, in preparation.
- [16] J.K. Hale, *Theory of Functional Differential Equations*, Springer, New York, 1977.
- [17] Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence*, Springer, Berlin, 1984.
- [18] F. Mertens, R. Imbihl, A. Mikhailov, Turbulence and standing waves in oscillatory chemical reactions with global coupling, *Chem. Phys.* 101 (1994) 9903–9908.
- [19] M.E. Parrot, Linearized stability and irreducibility for a functional differential equation, *SIAM J. Math. Anal.* 23 (1992) 649–661.
- [20] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, Springer, New York, 1988.
- [21] I.I. Vrabie, *Compactness Methods for Nonlinear Evolutions*, second ed., Pitman Monographs, Longman, Essex, 1995.
- [22] J. Wu, *Theory and Applications of Partial Functional Differential Equations*, Springer, New York, 1996.