

Pointwise gradient estimates of solutions to onedimensional nonlinear parabolic equations

PHILIPPE BÉNILAN⁺ and JESÚS ILDEFONSO DÍAZ

Abstract. We present here an improved version of the method introduced by the first author to derive pointwise gradient estimates for the solutions of one-dimensional parabolic problems. After considering a general quasilinear equation in divergence form we apply the method to the case of a nonlinear diffusion-convection equation. The conclusions are stated first for classical solutions and then for generalized and mild solutions. In the case of unbounded initial datum we obtain several regularizing effects for $t > 0$. Some unilateral pointwise gradient estimates are also obtained. The case of the Dirichlet problem is also considered. Finally, we collect, in the last section, several comments showing the connections among these estimates and the study of the free boundaries associated to the solutions of the diffusion-convection equation.

1. Introduction

This paper deals with a very old task which seems to have its origin in the work by S. N. Bernstein the beginning of last century ([14]): find a priori pointwise estimates on the (spatial) gradient of solutions of nonlinear second order equations. To be more precise we shall consider onedimensional parabolic equations.

Our main purpose is to present here an improved version of a general method introduced, originally, by the first author (see Section I.2 of Bénilan [7]). The method has important differences with respect to the way in which D. G. Aronson [3] derived, in 1969, this type of estimates for the porous media equation. His technique was later applied by different authors to many other nonlinear degenerate equations (see references in the surveys [35] and [46]). Such a already classical technique consists of, firstly, introducing the variable $u = f(w)$ with a concave function of the form $f(r) = C_1 r(C_2 - r)$, for some suitable constants $C_i > 0$ and, secondly, estimating the value of $|v_x|$ at a maximum of the auxiliary function $z = \zeta^2 w_x^2$, with ζ a cutoff function.

The method here presented seems to be better adapted to recognize the different nonlinear terms involved in the equation and so, it allows to obtain *sharp* pointwise gradient estimates. Moreover, our method provides quantitative estimates. In many cases these estimates have an important physical meaning and allow to deduce many other qualitative properties of the solutions.

2000 *Mathematics Subject Classification*: 35K65, 35K10, 35R35, 76D27.

Key words: Pointwise gradient estimates, one-dimensional parabolic equations, non linear diffusion-convection equation, regularizing effects, unilateral estimates, interfaces.

We shall limit the exposition to a general class of equations in divergence form

$$u_t = \sigma(u, u_x)_x, \quad (1)$$

but the method also applies to more general equations (see the exposition made in [7]). In order to explain some additional details, we consider the special case of the nonlinear diffusion-convection equation

$$u_t = \varphi(u)_{xx} + \psi(u)_x. \quad (2)$$

Equation (2) appears in many different contexts. For instance, unsaturated soil-moisture flows are modelled by (2) under the condition $\varphi'(0) = 0$. In that case the equation becomes degenerate and due to that a very rich structure arise concerning many different aspects: finite speed of propagation, stability results, etc. In the context of gas flow in a porous medium, the expression

$$V(t, x) = - \frac{\varphi(u(x, t))_x + \psi(u(x, t))}{u(x, t)}$$

represents the Eulerian description of the velocity at the position x and the time t of the “homogenized gas” (see, [22], [18]). It seems natural to expect boundedness of V which, in fact, leads to “optimal gradient bounds” for the solution u of (2) in some special cases as, for instance, $\varphi(u) = u^m$, $\psi(u) = u^n$, $1 < m$, $1 \leq n$. In that case, it is possible to find out some special solutions for which the modulus of continuity found via this gradient estimate can not be improved (see, e.g., [30], [25] and [33]).

The nature of “optimal gradient bounds” is much less evident in other contexts in which equation (2) arises (as, for instance, in Plasma Physics where $\varphi(u) = u^m$ and $0 < m < 1$ [15]) or even in the context of equation (2) when convection dominates over diffusion: $\varphi(u) = u^m$, $\psi(u) = u^n$ and $0 < n \leq 1 < m$: [20]).

One of the main features of the method, as presented here, is that it endows us with a “simple” analytical argument that allows to find sharp gradient bounds independently on the physical arguments involved in the concrete formulation.

The main idea of the method is to find an auxiliary nonlinear parabolic operator \mathcal{L} in such a way that a pointwise gradient estimate of the type

$$\sigma(u, u_x) \leq k(t, x, u) \quad (3)$$

holds as a consequence of the comparison principle applied to a function of the form $p = \sigma(u, u_x)/k(\cdot, \cdot, u)$, in the kernel of operator \mathcal{L} , and a constant function, i.e.

$$\mathcal{L}(p) = 0 \leq \mathcal{L}(1). \quad (4)$$

In this way, the possible choices of the function $k(t, x, u)$ are determined by the condition $\mathcal{L}(1) \geq 0$. Another characteristic of the method is that it can be used *locally in time and*

space and, in particular, it illustrates some regularizing effects (for instance some gradient bounds take place for any $t > 0$, independently of whether it already holds at $t = 0$ or not). In particular, we get some pointwise gradient estimates when $\varphi(u)$ is merely a maximal monotone graph (see Corollary 6).

The content of the paper is the following: Section 2 is devoted to a general presentation of the method for equation (1) and its special case (2) in the context of classical solutions. The application to the case of weaker notions of solution is carried out in Section 3. Pointwise gradient estimates are first obtained for the generalized solution of the Cauchy problem associated to (2) in Subsection 3.1. An unidirectional pointwise gradient estimate, that cease to be true if the direction of the spatial variable x is changed, is obtained in Subsection 3.2. The case of unbounded initial data and the proof of regularizing effects in the class of mild solutions are developed in Subsection 3.3. The derivation of pointwise gradient estimates for the case of a finite spatial interval is considered in Subsection 3.4. Finally, some comments on the connection between pointwise gradient estimates and the study of the free boundaries defined as the boundaries of the support of the solution are collected in Subsection 3.5.

Perhaps, some remarks on the long process of elaboration of this paper can be useful to some readers. The discussions on this subject by the authors were started after the stay of the first author at the University of Kentucky (during the 1980/1981 academic year). There he gave a general course on *Evolution Equations and Accretive Operators* (see [7]) and wrote a manuscript (without title and unpublished which we shall refer in the following as Bénilan [8]). In those days the authors were preparing their joint paper Bénilan-Díaz [10] submitted on May 18, 1981. The common point of view was a consequence of the results by the second author (in collaboration with R. Kersner) on the problem $u_t = \varphi(u)_{,xx} + \psi(u)_x$ for a case including $\varphi(u) = u^m$, $\psi(u) = u^n$ with $0 < n < 1 \leq m$. The first version of the Díaz-Kersner paper [20] appeared as internal report of the MRC at Wisconsin-Madison in 1982 although the definitive version waited until in 1987. In the period from 1980 to 1985 the authors of the present paper visited each other at least once a year thanks to the economical support received from the French Embassy at Madrid (see the Acknowledgments of [10]).

The first application of the present technique, which was not included in [8] was an alternative proof to Theorem 1 of the paper Díaz-Kersner [20] (correspondence maintained in March, 20, 1983). A second written document on the elaboration of the present paper was a letter of Bénilan to Díaz containing the statement in of Theorem 1 (i.e. for any φ including the case of a maximal monotone graph). Simultaneously to this period, both authors developed their normal activity teaching postgraduate courses in their universities (for instance, a part of the results of this paper was exposed by the second author during the 1985/1986 academic year). In this way, the connection of the (at this time unpublished) results with the ones by other authors was becoming a little bit barroque. For instance, a first version of the results was sent to B. H. Gilding (among many other colleagues) who mentions the notes in his paper (Gilding [27] submitted on September 17, 1986) as P. Bénilan -J. I. Díaz, *Unpublished notes* (see reference [3] of [27]). A first typed draft of the paper was discussed

by the authors at Montreal, in June of 1990 at the occasion of the participation by both authors at the Free Boundary Meeting. Then, they decided to add to the manuscript a part on the $L^1 - L^\infty$ smoothing effect incorporating some variations (occupying several pages written at Montreal and now in possession of the second author) to another unpublished manuscript by Bénilan (this time intitled by him as “On $u_t = \varphi(u)_{xx} + \psi(u)_x$: unfinished, 1978: [5]). Some related ideas of those modifications were published later in the paper by Bénilan and Bouillet [9]. The process of writing the definitive version of this paper was accelerated in the last five years before the death of Bénilan as the correspondence hold between both authors prove. In fact, Bénilan mentions this paper in the list of his publications (*paper No.104*) which he sent to the Departamento de Matematica Aplicada at the occasion of his nomination as Doctor Honoris Causa by the Universidad Complutense de Madrid which, although officially and definitively approved, it was programed for a too late date: two months after his death. The last meeting of both authors was in the “Journées Analyse non linéaire”, held at occasion of the 60th birthday of Philippe Bénilan, in October 27–29, 2000 at Les Moussières, Jura (France). We were talking on the last details to incorporate in to the paper and decided to publish it in some (not yet decided) journal.

2. The general method

In this section we introduce the main idea of the method. As usual, pointwise gradient estimates will be obtained as *a priori* estimates for any possible solution u . So, at the first step, they will be obtained in the more favorable setting of smooth solutions. Since the resulting estimates will be independent of the regularity used in the proof, they can be extended to the framework of weaker solutions. This second step will be illustrated in the next section.

2.1. A general quasilinear equation

Let Q be an open regular set of $\mathbb{R} \times \mathbb{R}$ and let u be a classical solution of the equation

$$u_t = \sigma(u, u_x)_x \text{ in } Q. \quad (5)$$

We assume $u \in C^{2,3}(Q)$ such that

$$m_- \leq u(t, x) \leq m_+ \text{ on } Q, \quad (6)$$

for some $m_- \leq m_+$. We also assume that $\sigma \in C^2([m_-, m_+] \times \mathbb{R})$.

Our main goal is to obtain pointwise estimates as (3) for suitable auxiliary functions k . As mentioned before, we start by assuming the regularity conditions: $k \in C^2(Q \times [m_-, m_+])$

and $k \neq 0$. The principal ingredient of the method is to find a (possibly nonlinear) parabolic operator, \mathcal{L} , vanishing when applied to the function

$$p(t, x) := \frac{\sigma(u(t, x), u_x(t, x))}{k(t, x, u)} \quad (t, x) \in Q \tag{7}$$

(some other choices of function p are also of interest: see [7]).

Using (5) we find

$$u_t = (kp)_x = (k_x + k_u + k_u u_x)p + kp_x \tag{8}$$

and differentiating (8) with respect to x ,

$$\begin{aligned} u_{tx} &= (k_{xx} + 2k_{xu}u_x + k_{uu}u_x^2 + k_u u_{xx})p + 2(k_x + k_u u_x)p_x \\ &\quad + kp_{xx} = \sigma_u u_x + \sigma_{u_x} u_{xx}. \end{aligned} \tag{9}$$

Differentiating (7) with respect to t we have

$$p_t k + p(k_t + k_u u_t) = \sigma_u u_t + \sigma_{u_x} u_{tx}.$$

Multiplying (8) by σ_u and (9) by σ_{u_x} and adding the resulting expressions we find that

$$\tilde{\mathcal{L}}(p, u_x) = 0$$

where

$$\begin{aligned} \tilde{\mathcal{L}}(p, u_x) &= p_t k + pk_t + (pk_u - \sigma_u)\{(k_x + k_u u_x)p + kp_x\} \\ &\quad - \sigma_{u_x}[(k_{xx} + 2k_{xu}u_x + k_{uu}u_x^2)p + 2(k_x + k_u u_x)p_x + kp_{xx}] \\ &\quad - k_u p[p_x k + p(k_x + k_u u_x) - \sigma_u u_x]. \end{aligned}$$

In order to obtain an expression independent of u_x we assume

$$\sigma_{u_x} > 0. \tag{10}$$

Thus we can define $S(u, \cdot) := \sigma(u, \cdot)^{-1}$ and so $u_x = S(u, kp)$. Finally, in order to find an operator \mathcal{L} such that $\mathcal{L}(p) = 0$ we define \mathcal{L} by

$$\mathcal{L}(p) := \tilde{\mathcal{L}}(p, S(u, kp))$$

(the dependence on u of the right hand side is understood as a (t, x) -dependent coefficient). The exact form of $\mathcal{L}(p)$ is irrelevant for our purposes. Our main aim is to compare p with 1 and so the relevant expression is $\mathcal{L}(1)$, i.e.

$$\mathcal{L}(1) = k_t + (k_u - \sigma_u(\tilde{S}(k))k_x - \sigma_{u_x}(\tilde{S}(k))\{k_{xx} + 2k_{xu}\tilde{S}(k) + k_{uu}\tilde{S}(k)^2\}) \tag{11}$$

where we used the notation $\sigma_u(\tilde{S}(k)) = \sigma_u(u, \tilde{S}(k))$ and $\tilde{S}(k) := S(u, k)$.

The following result is a trivial consequence of the maximum principle (see, e.g., the presentation made in [6]):

LEMMA 1. Assume (10),

$$k(t, x, u) > 0 \text{ on } Q \times (m_-, m_+), \quad (12)$$

$$\mathcal{L}(1) \geq 0 \text{ in } Q \quad (13)$$

$$\sigma(u, u_x)(t, x) \leq k(t, x, u(t, x)), \quad (t, x) \in \partial_p Q, \quad (14)$$

where $\partial_p Q$ denotes the parabolic boundary of Q . Then

$$\sigma(u, u_x)(t, x) \leq k(t, x, u(t, x)) \text{ for any } (t, x) \in \overline{Q}. \quad (15)$$

A concrete boundary value problem for which it is easy to check assumption (14) corresponds to the case of Neumann type boundary conditions.

LEMMA 2. Let $I = (a, b)$ with $-\infty \leq a < b \leq +\infty$ and let $Q = (0, T) \times I$ where $0 < T \leq +\infty$. Let u be a classical solution of (5) on Q . Assume (10), (12), (13),

$$\sigma(u, u_x)(0, x) \leq k(u)(0, x) \text{ for all } x \in I, \quad (16)$$

$$\begin{cases} (u_t - k_x - k_u u_x)(t, a) \geq 0 \text{ if } a > -\infty, \text{ for all } t \in (0, T), \\ (u_t - k_x - k_u u_x)(t, b) \leq 0 \text{ if } b < +\infty, \text{ for all } t \in (0, T), \end{cases} \quad (17)$$

and

$$\left\| \frac{k_x + k_u u_x}{k} \right\|_{L^\infty((0, T) \times \partial I)} < \infty. \quad (18)$$

Then estimate (15) holds.

Proof. From (8) we have that

$$-p_x + p \left[- \left\{ \frac{k_x + k_u u_x}{k} \right\} \right] = \frac{-u_t}{k}$$

So, if ν denotes the exterior “unit outer normal” to ∂I (i.e. $\nu(a) = -1$, $\nu(b) = 1$), we deduce that

$$-p_x \nu + p \left[-\nu \left\{ \frac{k_x + k_u u_x}{k} \right\} \right] = -\frac{u_t \nu}{k} \text{ on } (0, T) \times \partial I.$$

Since we assume (18) we can apply the comparison principle. Taking $\bar{p}(t, x) = 1$, assumption (16) implies that $p \leq \bar{p}$ on $\{0\} \times (a, b)$ and (17) implies that

$$-\bar{p}_x \nu + \bar{p} \left[-\nu \left\{ \frac{k_x + k_u u_x}{k} \right\} \right] \leq -\frac{u_t \nu}{k} \text{ on } (0, T) \times \partial I.$$

In consequence $p \leq \bar{p}$ on $[0, T] \times \bar{I}$. □

In the special case of

$$k(t, x, u) = \theta(u), \quad (19)$$

we deduce, from (11), that

$$\mathcal{L}(1) = -\sigma_{u_x}(\tilde{S}(\theta))\tilde{S}(\theta)^2\theta'' \quad (20)$$

and then the following result holds:

COROLLARY 1. *Let u be a smooth solution of (5) satisfying (6). Assume (10) and let $\theta \in C^2([m_-, m_+])$ such that*

$$\theta > 0 \text{ and } \theta'' \leq 0 \text{ on } [m_-, m_+]. \quad (21)$$

Then

$$\frac{\sigma(u, u_x)}{\theta(u)} \leq \sup_{\partial_p Q} \frac{\sigma(u, u_x)}{\theta(u)}. \quad (22)$$

REMARK 1. Conclusion (22) expresses a “conservation of the regularity”: if the estimate is true on the parabolic boundary of Q it also holds in the interior of Q . We emphasize the large generality on θ assumed in (21) and that, by passing to the limit, it is possible to extend the conclusion for θ concave such that $\theta \geq 0$. As a general remark, we point out that the Bernstein technique, as described before, still works in several space dimensions for radially symmetric solutions (see, for instance, [28], for a related result). We also mention that the method can be applied to a general class of quasilinear equations (see the unpublished notes by the first author [8] and the application made in [45]).

In the rest of this section we illustrate the applicability of the method by considering equation (2).

2.2. A nonlinear diffusion-convection equation

As it was already mentioned, a physically relevant special case of equation (1) is

$$u_t = \varphi(u)_{xx} + \psi(u)_x \text{ in } Q, \quad (23)$$

where φ is a nondecreasing function with $\varphi(0) = 0$. For the sake of simplicity, we assume in this subsection that $\varphi \in C^2([m_-, m_+])$, $\varphi' > 0$ and $\psi \in C^1([m_-, m_+])$. In the applications we shall present here, we assume the (t, x) -dependence of function k of the form

$$k(t, x, u) = \rho(t, x)\theta(u). \quad (24)$$

Then, Lemma 1 leads to the following result

LEMMA 2. Let $Q \subset \mathbb{R}^+ \times \mathbb{R}$. Let u be a smooth solution of (23) satisfying (6). Let $\rho > 0$ and $\theta > 0$ be smooth functions such that

$$0 \leq \frac{\rho_t}{\rho} + \left[\theta' - \varphi'' \left(\frac{\rho\theta - \psi}{\varphi'} \right) - \psi'' \right] \frac{\rho_x}{\rho} - \varphi' \left[\frac{\rho_{xx}}{\rho} + 2 \frac{\rho_x}{\rho} \frac{\theta'}{\theta} \left(\frac{\rho\theta - \psi}{\varphi'} \right) + \frac{\theta''}{\theta} \left(\frac{\rho\theta - \psi}{\varphi'} \right)^2 \right] \text{ in } Q \tag{25}$$

$$\varphi(u)_x + \psi(u) \leq \rho\theta(u) \text{ on } \partial_\rho Q. \tag{26}$$

Then

$$\varphi(u)_x + \psi(u) \leq \rho\theta(u) \text{ on } Q. \tag{27}$$

In the next section we shall give several applications of Lemma 2 to the case of weak solutions of (23). Nevertheless, some previous consequences for the cases in which condition (26) is trivially satisfied (the Cauchy and the Neumann problems) can be easily derived for the special choice $\rho(t, x) = 1/\sqrt{t}$.

COROLLARY 2. A. Let $Q = (0, T) \times \mathbb{R}$ with $0 < T \leq \infty$ and $\theta \in C^2([m_-, m_+])$ satisfying $\theta > 0$ and

$$\frac{\varphi'}{2} + \theta\theta'' \left(1 - \psi \frac{\sqrt{t}}{\theta} \right)^2 \leq 0 \text{ for any } t \in (0, T] \text{ on } [m_-, m_+]. \tag{28}$$

Then, if u is a smooth solution of (23) on Q satisfying (6), we have the estimate

$$\varphi(u)_x + \psi(u) \leq \frac{\theta(u)}{\sqrt{t}} \text{ on } (0, T] \times \mathbb{R}. \tag{29}$$

B. Let $I = (a, b)$ with $-\infty < a < b < +\infty$ and $Q = (0, T) \times I$. Assume that θ satisfies the conditions of part A and let u be a smooth solution of equation (23) on Q verifying (6) and the homogeneous Neumann boundary conditions

$$0 = (\varphi(u)_x + \psi(u))(t, a) = (\varphi(u)_x + \psi(u))(t, b) \text{ for } t \in (0, T). \tag{30}$$

Then inequality (29) holds on $(0, T] \times I$. The same conclusion holds if we replace the boundary condition (30) by

$$0 = \varphi(u)_x(t, a) = \varphi(u)_x(t, b) \tag{31}$$

and assume that $\psi \leq 0$ on (m_-, m_+) .

REMARK 2. In contrast to Corollary 1 the above result shows the first “regularizing effect”: estimate (29) holds for any $t > 0$, independently of either it takes place at $t = 0$ or not. Other regularizing effects will be given in Subsection 3.3.

Another interesting choice of function $\rho(t, x)$ leading to local effects, independently of the boundary conditions, is $\rho(t, x) = \xi(x)/\sqrt{t}$ with $\xi(x)$ vanishing on the boundary of the spatial domain.

COROLLARY 3. Let $I = (a, b)$ with $-\infty < a < b \leq +\infty$ and $Q = (0, T) \times I$. Assume that $\theta \in C^2([m_-, m_+])$, $\theta > 0$, $\xi \in C^2(I)$, $\xi > 0$ on I and $\xi = +\infty$ on ∂I . Assume that

$$0 \leq -\frac{1}{2t} + \left[\theta' - \varphi'' \left(\frac{\theta \xi / \sqrt{t} - \psi}{\varphi'} \right) - \psi'' \right] \frac{\xi_x}{\xi} - \varphi' \left[\frac{\xi_{xx}}{\xi} + 2 \frac{\xi_x}{\xi} \frac{\theta'}{\theta} \left(\frac{\theta \xi / \sqrt{t} - \psi}{\varphi'} \right) + \frac{\theta''}{\theta} \left(\frac{\theta \xi / \sqrt{t} - \psi}{\varphi'} \right)^2 \right] \tag{32}$$

Let u be a smooth solution of equation (23) on Q satisfying (6). Then we have the inequality

$$\varphi(u)_{,x} + \psi(u) \leq \frac{\theta(u)\xi(x)}{\sqrt{t}} \quad \text{on } (0, T] \times I. \tag{33}$$

Condition (32) leads to interesting results once we know more about the functions φ and ψ , for instance, if $\varphi(u) = u^m$, $\psi(u) = u^n$ for some positive m and n (see Subsection 3.4).

3. Applications to generalized and mild solutions

Corollaries 2 and 3 will be used in this section for two different objectives: (i) to extend estimates (29) and (33) to weaker notions of solution of the Cauchy and Dirichlet problems associated to equation (23) (and so under much more generality on the data φ , ψ and u_0), and, (ii) to present explicitly different choices of the function θ according φ and ψ .

3.1. Generalized solutions

We start by considering the class of nonnegative bounded generalized solutions of the Cauchy problem

$$0 \leq u_0(x) \leq M_0 \quad \text{for any } x \in \mathbb{R}. \tag{34}$$

$$\begin{cases} u_t = \varphi(u)_{,xx} + \psi(u)_{,x} & \text{in } (0, T) \times \mathbb{R}, \\ u(0, x) = u_0(x) & x \in \mathbb{R}, \end{cases} \tag{35}$$

with

$$0 \leq u_0(x) \leq M_0 \text{ for } x \in \mathbb{R}, \tag{36}$$

for some M_0 . We assume that

$$\varphi, \psi \in C^0([0, M_0]), \varphi(0) = 0, \varphi \text{ strictly increasing} \tag{37}$$

as well as the following technical conditions (see Subsection 3.3)

$$\begin{cases} \exists \delta_0 > 0 \text{ such that } \varphi, \psi \in C^2((0, M_0 + \delta_0]) \text{ and} \\ \varphi'' \text{ and } \psi'' \text{ are locally Hölder continuous on } (0, M_0 + \delta_0]. \end{cases} \tag{38}$$

We recall that given $0 < T \leq +\infty$, a function $u(t, x)$ is said to be a *nonnegative bounded generalized solution* of equation (23) if $u \in C([0, T] \times \mathbb{R})$,

$$0 \leq u(t, x) \leq M_0 \text{ for } x \in \mathbb{R} \text{ and } t \in [0, T], \tag{39}$$

for some $m > 0$ and satisfies the integral identity

$$\begin{aligned} & \int_{t_1}^{t_2} \int_{x_1}^{x_2} \{u\phi_t + \varphi(u)\phi_{xx} - \psi(u)\phi_x\} dx dt \\ &= \int_{x_1}^{x_2} \{u(t_2, x)\phi(t_2, x) - u(t_1, x)\phi(t_1, x)\} dx \\ &+ \int_{t_1}^{t_2} \{u(t, x_2)\phi(t, x_2) - u(t, x_1)\phi(t, x_1)\} dt \end{aligned}$$

for all non-empty bounded rectangles $(t_1, t_2) \times (x_1, x_2)$ and nonnegative functions $\phi \in C^{1,2}([t_1, t_2] \times [x_1, x_2])$ such that $\phi(t, x_1) = \phi(t, x_2) = 0$ for all $t \in [t_1, t_2]$.

We recall (see, e.g., [20]) that any generalized solution is a weak solution in the sense that the above integral identity can be replaced by the condition $u_t = \varphi(u)_{xx} + \psi(u)_x$ in $D'(Q)$.

Different results on the existence and uniqueness of a generalized solution u of the Cauchy Problem (35) are today available in the literature (see e.g. [13] [24] and [17]). For our present purposes we shall make use of the results of Gilding [27] requiring, merely, the continuity of the initial datum. We have:

THEOREM 1. *Assume that φ and ψ satisfy (37) and (38). Let $u_0 \in C^0(\mathbb{R})$ satisfying (36). Let $u \in C([0, T] \times \mathbb{R})$ be the generalized solution to the Cauchy problem (35). Assume that there exists a function $\lambda \in C^0([0, M_0])$, concave, increasing and such that*

$$M < \infty \tag{40}$$

where

$$M := \int_0^{M_0} \frac{d\varphi(s)}{\lambda(s)\lambda'(s)}.$$

Finally, assume that there exists a constant $C \geq 0$ such that

$$\psi(s) \leq C\lambda(s) \text{ for any } s \in (0, M_0). \tag{41}$$

Then

$$\varphi(u)_x + \psi(u) \leq \left[\sqrt{4C^2T + 2M} + 2C\sqrt{T} \right] \frac{\lambda(u)}{\sqrt{t}} \text{ in } D'((0, T) \times \mathbb{R}). \tag{42}$$

Before giving the proof we shall state two immediate consequences.

COROLLARY 4. *Let φ and ψ satisfying (37) and (38). Assume ψ be such that*

$$\psi(s) \leq C\sqrt{s} \text{ for any } s \in (0, M_0), \tag{43}$$

for some constant $C \geq 0$. Then, if $u_0 \in C^0(\mathbb{R})$ satisfies (36) and u is the generalized solution to the Cauchy Problem (35) we have the estimate

$$\varphi(u)_x + \psi(u) \leq 2\sqrt{\frac{u\varphi(M_0)}{t}} \text{ in } D'((0, T) \times \mathbb{R}). \tag{44}$$

COROLLARY 5. *Let φ and ψ satisfying (37) and (38). Let $u_0 \in C^0(\mathbb{R})$ satisfying (36) and u be the generalized solution of the Cauchy Problem (35). Then $\varphi(u(t, \cdot))_x \in L^\infty(\mathbb{R})$ and*

$$\|\varphi(u(t, \cdot))_x\|_{L^\infty(\mathbb{R})} \leq 2 \left[\frac{\varphi(M_0)M_0}{2} \right]^{1/2} + N \tag{45}$$

for any $t > 0$, where

$$N = \sup_{[0, M_0]} \psi - \inf_{[0, M_0]} \psi. \tag{46}$$

Proof of Corollaries 4 and 5. Corollary 4 follows from the fact that function $\lambda(s) = \sqrt{s}$ satisfies (40) for any continuous nondecreasing function φ . To prove Corollary 5 we point out that u is also a solution of the equation

$$u_t = \varphi(u)_{xx} + \psi_d(u)_x, \tag{47}$$

where $\psi_d(s) := \psi(s) + d$ with $d \in \mathbb{R}$ arbitrary. By choosing $d = -d_1$ with $d_1 = \sup\{\psi(s) : s \in [0, M_0]\}$ condition (43) is verified for ψ_d (since by the maximum principle u we know that u satisfies $0 \leq u \leq M_0$). Then (44) implies

$$\varphi(u)_x \leq \left(\frac{2M_0\varphi(M_0)}{t} \right)^{1/2} + d_1 - d_2$$

where $d_2 = \inf\{\psi(s) : s \in [0, M_0]\}$. To obtain the estimate in the reverse sense we define $\tilde{u}(t, x) = u(t, -x)$. Then \tilde{u} satisfies

$$\tilde{u}_t = \varphi(\tilde{u})_{,xx} + \tilde{\psi}_d(\tilde{u})_{,x} \tag{48}$$

with $\tilde{\psi}_d(s) = -\psi(s) - d$. Then, by choosing $d = -d_2$, we conclude that

$$-\varphi(u)_{,x} = \varphi(\tilde{u})_{,x} \leq 2 \left(\frac{M_0 \varphi(M_0)}{t} \right)^{1/2} + d_1 - d_2,$$

and so estimate (45) holds.

Proof of Theorem 1. To get the result we shall perform an approximation process according classical ideas. Our main purpose is to approximate u by a sequence of smooth functions. Let δ_0 be given in (38). Fix $\delta \in (0, \delta_0)$ arbitrary and let $k_0 \in \mathbb{N}$ be such that

$$2^{-k_0+1} \leq \varphi(M_0 + \delta) - \varphi(M_0). \tag{49}$$

For $k \geq k_0$ we define the cylinder $Q_k = (0, T) \times (-k, k)$ and consider the boundary value problem

$$u_t = \varphi(u)_{,xx} + \psi(u)_{,x} \text{ in } Q_k, \tag{50}$$

$$u(t, \pm k) = M_0 \text{ for all } t \in (0, T), \tag{51}$$

$$u(0, x) = u_{0,k}(x) \text{ on } (-k, k), \tag{52}$$

where the sequence $\{u_{0,k}\}$ satisfies the following properties: (i) $u_{0,k}(x) \leq M_0 + \delta$ for all $x \in \mathbb{R}$, (ii) there exists $\varepsilon_k > 0$ such that $\varepsilon_k \leq u_{0,k}(x)$ for all $x \in \mathbb{R}$, (iii) there exists $\alpha_k \in (0, 1]$ such that $u_{0,k} \in C^{2+\alpha_k}(\mathbb{R})$, (iv) $u_{0,k+l}(x) \leq u_{0,k}(x)$ for all $x \in \mathbb{R}$, (v) $u_{0,k} \downarrow u_0$ as $k \uparrow \infty$ uniformly on compact subsets of \mathbb{R} , (vi) $u_{0,k}(x) = M_0 + \delta$ for all $|x| > k - 1/2$.

This approximation process was made explicit in [27]. Since properties (ii), (iii) and (vi) hold we can apply the theory of nondegenerate parabolic problems (see e.g. [38]) and so (50), (51), (52) has a unique classical solution $u_k(t, x)$. Moreover from (i) and (ii) we deduce that $\varepsilon k \leq u_k(t, x) \leq M_0 + \delta$ for all $(t, x) \in Q_k$. By the maximum principle, property (iv) implies, that $u_{k+l}(t, x) \leq u_k(t, x)$ for all $(t, x) \in \overline{Q}_k$ (assumed $k \geq k_0$). Hence we can define

$$u^*(x, t) = \lim_{k \nearrow \infty} u_k(x, t) \text{ for all } (t, x) \in [0, T] \times \mathbb{R}.$$

It is easy to see that u^* is a generalized solution. It is clear that it remains to show that $u^* \in C^0([0, T] \times \mathbb{R})$ since by the uniqueness of the solution, $u = u^*$. The continuity of u^*

will be obtained similarly to the proof of Theorem 1 of [27]: as in the proof of Corollary 4, we first conclude the gradient estimate for $t \in (0, T]$

$$\|\varphi(u_k(t, \cdot))_x\|_{L^\infty((-k, k))} \leq 2 \left(\frac{\varphi(M_0 + \delta)(M_0 + \delta)}{t} \right)^{1/2} + N$$

with N given by (46). This implies the existence of a modulus of continuity of u_k in x and (by the results of Kruzkov [37] and [24]) also in t . This modulus of continuity is independent of k and so, by passing to the limit in k , it also applies to $u(t, x)$ for $t \in (0, T]$ and $x \in \mathbb{R}$. The continuity at $t = 0$, i.e. the inequalities

$$u_0(x_0) \leq \lim_{(t,x) \rightarrow (0,x_0)} \inf u(t, x) \leq \lim_{(t,x) \rightarrow (0,x_0)} \sup u(t, x) \leq u_0(x_0)$$

for any $x_0 \in \mathbb{R}$, were proved in [24] by using a local barrier function argument which literally applies to our case.

The rest of the proof consists in obtaining the gradient estimate (42), on $(0, T) \times (-k, k)$, for the smooth solutions u_k but replacing M_0 by M_δ in (42), where

$$M_\delta := \int_0^{M_0+\delta} \frac{d\varphi}{\lambda(s)\lambda'(s)}.$$

Then, passing to the limit in k , we deduce the same estimate for u (but now in the sense of $D'((0, T) \times \mathbb{R})$). Finally, as δ is arbitrary we get (42).

In order to estimate $\varphi(u_k)_x + \psi(u_k)$ we apply Lemma 2 with $k(s) = \theta(s)/\sqrt{t}$, where $\theta(s)$ is a convex, increasing and positive function to be determined. Notice that condition (16) trivially holds. On the other hand, from the boundary condition (51) we deduce that $(u_k(t, \pm k))_t = 0$ for $t \in (0, T)$. Moreover as $u_k(t, x) \leq M_0 + \delta$ we deduce that $u_x(t, -k) \leq 0$ and $u_x(t, k) \geq 0$ for all $t \in (0, T)$. Then, conditions (17) holds. Finally, since u_k is a positive classical solution, condition (18) is also satisfied.

Thus, it only remains to check condition (13) or, equivalently, (28). We introduce the function

$$\theta(s) := e\lambda(s - cj(c)) \tag{53}$$

where e and c are positive constants to be chosen later and j is the convex function defined by

$$j(s) := \int_0^s \int_0^r \frac{d\varphi(z)}{\lambda(z)\lambda'(z)} dr. \tag{54}$$

Our first condition on c is $c < 1/M$. So from the convexity of j we deduce that $\theta''(s) < 0$ and $\theta'(s) > 0$ if $s \in (0, M_0 + \delta]$. Moreover

$$\theta(s) \geq e\lambda((1 - cM)s) \geq e(1 - cM)\lambda(s) > 0 \quad \text{if } s \in (0, M_0 + \delta]. \tag{55}$$

In order to check condition (28), we observe that

$$\theta''(s) = e\lambda''(s - cj(s))(1 - cj''(s))^2 - ec\lambda'(s - cj(s))\frac{\varphi'(s)}{\lambda(s)\lambda'(s)}.$$

Then, (28) is equivalent to

$$ec\frac{\lambda(s - cj(s))}{\lambda(s)}\frac{\lambda'(s - cj(s))}{\lambda'(s)}\left(1 - \frac{\psi(s)\sqrt{t}}{e\lambda(s - cj(s))}\right)^2 \geq \frac{1}{2}. \tag{56}$$

But the concavity of λ implies that $\lambda'(s - cj(s)) \geq \lambda'(s)$. Then, from assumption (41)

$$\left(1 - \frac{\psi(s)\sqrt{t}}{e\lambda(s - cj(s))}\right) \geq \left(1 - \frac{C\sqrt{t}}{e(1 - cM_\delta)}\right).$$

So, condition (56) is verified if

$$e^2c(1 - cM_\delta)\left(1 - \frac{2C\sqrt{T}}{e(1 - cM_\delta)}\right) \geq \frac{1}{2}. \tag{57}$$

Taking $c = 1/(2M_\delta)$, (57) reduces to the inequality

$$e^2 - 4eC\sqrt{T} - 2M_\delta \geq 0,$$

and, finally, the choice

$$e = 2C\sqrt{T} + \sqrt{4C^2T + 2M_\delta}$$

leads to the desired estimate (42) completing the proof of Theorem 1.

3.2. Unidirectional estimates

It is important to point out that sometimes, it is possible to get pointwise gradient estimates of *unidirectional type* for the solutions of the diffusion-convection Cauchy problem (35), i.e. inequalities which become false if we reverse the direction of the spatial variable x by making the transformation $\tilde{x} = -x$. This is illustrated in the next result which exhibits an alternative proof to a result derived in [19].

THEOREM 2. *Let $u_0 \in C^0(\mathbb{R})$ satisfies (36), with M_0 small enough, and let u be the generalized solution to the Cauchy problem (35) corresponding to*

$$\varphi(u) = u^m, \psi(u) = -u^n \text{ and } 0 < n < 1 \leq m.$$

Then

$$(u^m)_x \geq -K\frac{u^m}{t} \text{ in } D((0, T) \times \mathbb{R}), \tag{58}$$

for some $K > 0$ depending only on M_0, m , and n .

Proof. As in the proof of the previous theorem we can assume u to be positive and smooth enough. Let $\lambda(r)$ be a regular function, to be chosen later, such that $\lambda(r)$ is a smooth function transforming $[0, w_0]$ onto $[0, m \ln M_0]$, for some $w_0 > 0$ and such that

$$K = \sup_{w \in [0, w_0]} \lambda'(w) > \inf_{w \in [0, w_0]} \lambda'(w) = C > 0. \tag{59}$$

Then, the equation

$$m \ln u(t, x) = \lambda(w(t, x))$$

defines a regular function $w(t, x)$ such that $0 < w(t, x) \leq w_0$ for all $(t, x) \in [0, T] \times \mathbb{R}$. In order to prove the estimate we introduce the function

$$z(t, x) = w_x(t, x)t.$$

Then, it is clear that the conclusion of the theorem will follow once we prove that $z(t, x) \geq -1$ on $[0, T] \times \mathbb{R}$, since

$$\frac{(u^m)_x}{u^m} = \lambda(w)_x = \lambda'(w)w_x \geq -\frac{K}{t}.$$

It is not difficult to see that z satisfies $\mathcal{L}(z) = 0$ where

$$\begin{aligned} \mathcal{L}(z) = & z_t - mu^{m-1}z_{xx} - \frac{u^{m-1}}{t} \left[(3m-1)\lambda'(w) + m \frac{\lambda''(w)}{\lambda'(w)} \right] z z_x \\ & - nu^{n-1}z_x - \frac{u^{m-1}}{t^2} \left[(m-1)\lambda'(w)^2 + m \left(\frac{\lambda''(w)}{\lambda'(w)} \right)_w + (2m-1)\lambda''(w) \right] z^3 \\ & + \frac{n(n-1)}{mt} u^{n-1} \lambda'(w) z^2 - \frac{z}{t}. \end{aligned}$$

Hence,

$$\begin{aligned} \mathcal{L}(-1) = & \frac{u^{m-1}}{t^2} \left[(m-1)\lambda'(w)^2 + m \left(\frac{\lambda''(w)}{\lambda'(w)} \right)_w + (2m-1)\lambda''(w) \right] \\ & - \frac{n(n-1)}{mt} u^{n-1} \lambda'(w) z^2 + \frac{1}{t}. \end{aligned}$$

So, we have to choose $\lambda(r)$ such that $\mathcal{L}(-1) \leq 0$ (notice that the comparison at $t = 0$ is obviously satisfied and that we can argue starting with u_k and then passing to the limit as in the previous theorem). The above condition holds once that

$$(m-1)\lambda'(w)^2 + m \left(\frac{\lambda''(w)}{\lambda'(w)} \right)_w + (2m-1)\lambda''(w) \leq 0 \tag{60}$$

and

$$-\frac{n(1-n)C}{mM_0^{1-n}} + 1 \leq 0, \tag{61}$$

with C given by (59). The condition (61) is satisfied by taking

$$C \geq \frac{mM_0^{1-n}}{n(1-n)}.$$

To simplify the study of condition (60) we shall take $\lambda(r)$ such that $\lambda''(r) \equiv -\gamma < 0$ for any $r \in [0, w_0]$. Then

$$\left(\frac{\lambda''(w)}{\lambda'(w)} \right)_w = \frac{-\gamma^2}{\lambda'(w)^2} < \frac{-\gamma^2}{K^2},$$

and (60) is satisfied once

$$(m-1)K^2 - m\frac{\gamma^2}{K^2} - (2m-1)\gamma \leq 0. \tag{62}$$

Let us take $\lambda(r) = ar^2 + br$, with $a = (-\gamma)/2$ and $b = K$. By taking $\gamma = K$, (62) is transformed into

$$(m-1)K^2 - (2m-1)K - m \leq 0,$$

and so, it is fulfilled once that

$$0 < K \leq 2 < \frac{(2m-1) + ((2m-1)^2 + 4m)^{1/2}}{2(m-1)}.$$

Finally, it is a routine matter to check that, among the many possible choices, $\lambda(r) = r(2-r)$ satisfies all the requirements once we assume

$$M_0^{1-n} < \min \left\{ \frac{1}{2m}, \frac{1}{\frac{m}{2} + \frac{m}{2n(1-n)}} \right\}.$$

□

As we shall comment later, in Subsection 3.4, estimate (58) ceases to be true if we reverse the direction of x .

3.3. Mild solutions: smoothing effects

In the case of $u_0 \in L^1(\mathbb{R})$ it is possible to show some regularizing effect for the, so called, *mild solution* of the Cauchy problem (35). To introduce this notion we consider the implicit time-discretization

$$(ITD) \left\{ u \frac{w_i - w_{i-1}}{t_i - t_{i-1}} = \varphi(w_i)_{xx} + \psi(w_i)_x + f_i \text{ in } D'(\mathbb{R}), \right.$$

for a given $\epsilon > 0$ and a time discretization $t_0 = 0 < t_1 < \dots < t_n \leq T, t_i - t_{i-1} < \epsilon, T - t_n < \epsilon$, and for given data $w_0, f_i \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$. We say that u is a mild solution of (35) if $u \in C([0, +\infty) : L^1(\mathbb{R}))$, $u(0, \cdot) = u_0(\cdot)$, and, for any $\epsilon > 0$ there exists $(t_0, t_1, \dots, t_n, w_0, w_1, \dots, w_n)$ satisfying (ITD) with

$$\|w_0 - u_0\|_1 \leq \epsilon, \sum_i \int_{t_{i-1}}^{t_i} \|f_i\|_1 dt \leq \epsilon$$

and such that $\|u(t) - w_i\|_1 \leq \epsilon$ for any $t \in (t_{i-1}, t_i], i = 1, \dots, n$. The existence of a mild solution is due to B enilan and Tour e [13]. Moreover, it was proven there that, under suitable additional regularity on the data φ, ψ and u_0 , the mild solution is also a weak solution.

THEOREM 3. *Let φ and ψ satisfy (37) and let $u_0 \in L^1(\mathbb{R}), u_0 \geq 0$. Let $u \in C([0, T] : L^1(\mathbb{R}))$ be the mild solution to the Cauchy problem (35). Assume that*

$$R(I) = \mathbb{R}, \tag{63}$$

where

$$I(u) = \text{sign}(u) \int_0^u (\varphi(u) - \varphi(r)) dr.$$

Then

$$\|u(t, \cdot)\|_{L^\infty(\mathbb{R})} \leq I^{-1} \left(\frac{\|u_0\|_{L^1(\mathbb{R})}^{3/2}}{4t} \right). \tag{64}$$

Moreover; $\varphi(u(t, \cdot))_x \in L^\infty(\mathbb{R})$ and

$$\|\varphi(u(t, \cdot))_x\|_{L^\infty(\mathbb{R})} \leq 2 \left[\frac{\varphi(m(t))m(t)}{2} \right]^{1/2} + N \tag{65}$$

for any $t > 0$, with

$$m(t) = I^{-1} \left(\frac{\|u_0\|_{L^1(\mathbb{R})}^{3/2}}{4t} \right)$$

and N given by (46).

Proof. Using the continuity of the solution with respect to the data and the functions φ and ψ (see [12]) we may assume that $u_0, \varphi, \psi \in C^\infty(\mathbb{R})$ with $\varphi' > 0$, so that u is a classical solution with $u_x, u_t \in C([0, T] : L^1(\mathbb{R}))$. From the definition of I we deduce that $I(u)_x = |u| \varphi(u)_x$. Moreover

$$\frac{d}{dt} \int_{\mathbb{R}} \int_0^u I(r) dr = \int_{\mathbb{R}} I(u) u_t = - \int_{\mathbb{R}} I(u)_x \varphi(u)_x.$$

But, using Hölder inequality and the L^1 -contraction of the associated semigroup ([13], [12]) we get that

$$\begin{aligned} \|I(u(t, \cdot))\|_{L^\infty(\mathbb{R})}^2 &\leq \frac{1}{4} \|I(u)_x\|_{L^1}^2 \leq \frac{1}{4} \left[\int_{\mathbb{R}} |u|^{1/2} |u|^{1/2} \varphi(u)_x \right]^2 \\ &\leq \frac{1}{4} \|u_0\|_{L^1(\mathbb{R})} \left[\int_{\mathbb{R}} |u| \varphi(u)_x^2 \right] = -\frac{1}{4} \|u_0\|_{L^1(\mathbb{R})} \int_{\mathbb{R}} I(u)_x \varphi(u)_x \\ &= -\frac{1}{4} \|u_0\|_{L^1(\mathbb{R})} \frac{d}{dt} \int_{\mathbb{R}} \int_0^u I(r) dr. \end{aligned}$$

On the other hand, if we define the function

$$l(t) = \int_{\mathbb{R}} \left(\int_0^{u(t,x)} I(r) dr \right) dx,$$

then

$$l'(t) \leq \|I(u(t, \cdot))\|_{L^\infty(\mathbb{R})} \|u(t)\|_{L^1(\mathbb{R})} \leq \|I(u(t, \cdot))\|_{L^\infty(\mathbb{R})} \|u_0\|_{L^1(\mathbb{R})}.$$

So, $l(t)$ satisfies the ordinary differential inequality

$$l'(t) + \frac{4l(t)^2}{\|u_0\|_{L^1(\mathbb{R})}} \leq 0.$$

Dividing by $l(t)^2$, integrating and using that $l(t_0) > 0$ for any $t_0 \geq 0$ we get that

$$l(t) \leq \frac{\|u_0\|_{L^1(\mathbb{R})}^2}{4t},$$

and finally

$$\|I(u(t))\|_{L^\infty(\mathbb{R})} \leq \frac{\|u_0\|_{L^1(\mathbb{R})}^{3/2}}{4t},$$

which leads to estimate (64). The rest is an obvious application of the arguments of Corollary 4. □

REMARK 3. Some related results can be found in [9] and [39]. It is possible to apply the general method to the study of one-side estimates on u_t . Among the many papers devoted to this task, we only mention here the presentation made in [7] and the articles [4] and [45] (many other references can be found in the surveys [35] and [46]). This allows to prove stronger differentiability on the solutions of (2) leading to the existence of the, so called, *strong solutions* satisfying that u_t , $\varphi(u)_{x,x}$ and $\psi(w_i)_x$ are in $L^1((0, T) \times \mathbb{R})$ (see also the different approach made in [11]).

3.4. The case of a bounded interval

Concerning the case in which the solution is not defined on \mathbb{R} but on a bounded interval, we can get interior gradient estimates without using any concrete information on the boundary conditions. We illustrate it by considering the special case of $\psi \equiv 0$.

PROPOSITION 1. *Let $I = (a, b)$ with $-\infty < a < b < +\infty$ and $Q = (0, T) \times I$. Let $u \in L^\infty(Q)$ be a positive solution of*

$$u_t = (u^m)_{xx} \text{ in } Q.$$

Let $\xi \in C^2(I)$, $\xi > 0$ on I and $\xi = +\infty$ on ∂I such that

$$|\xi'(x)| \leq \xi(x)^2, \quad |\xi''(x)| \leq \xi(x)^3 \quad \text{for any } x \in I.$$

Then, if $M_0 := \sup_Q u$,

a) if $0 < m < 1$,

$$|(u^m)_x| \leq \frac{\xi(x)u^m}{\sqrt{t}} \left(\frac{\sqrt{T}}{1-m} + \sqrt{\frac{M_0}{2(1-m)}} \right),$$

b) if $m = 1$, for any $\varepsilon > 0$,

$$|(u^m)_x| \leq \frac{\xi(x)u}{\sqrt{t}} \left(\varepsilon + \log \frac{M_0}{u} \right) \left(2\sqrt{T} \left(1 + \sup \left(1, \frac{1}{\varepsilon} \right) \right) \right) + \sqrt{\frac{2}{\varepsilon}},$$

c) if $m > 1$,

$$|(u^m)_x| \leq \frac{\xi(x)\lambda_0(\alpha u - u^m)}{\sqrt{t}}$$

with $\alpha > \alpha_0 := \sup_Q u^{m-1}$ and λ_0 such that

$$\lambda_0 \leq \frac{2\sqrt{T}}{(m-1)(\alpha - \alpha_0)} (m(\alpha - \alpha_0) + 2(m-1)\alpha_0) + \sqrt{\frac{2}{(m-1)(\alpha - \alpha_0)}}.$$

Proof. In the case a) we take $\theta(u) = \lambda u^m$. Then condition (32) is satisfied if.

$$-\lambda^2(1-m) + \frac{u^{1-m}}{2} + \lambda\sqrt{t}\frac{\xi'}{\xi^2}(2m-1) + mT\frac{\xi''}{\xi^3} \leq 0,$$

which is implied by

$$-\lambda^2(1-m) + \frac{M_0^{1-m}}{2} + |\lambda|\sqrt{T}\frac{\xi'}{\xi^2}|2m-1| + mT \leq 0.$$

This last condition holds once that

$$|\lambda| \geq \lambda_0 = \frac{1}{1-m} \left(\left| m - \frac{1}{2} \right| \sqrt{T} + \left(T \left| m - \frac{1}{2} \right|^2 + (1-m) \left(2mT + \frac{M_0^{1-m}}{2} \right) \right)^{1/2} \right).$$

A simple computation shows that

$$\lambda_0 \leq \frac{\sqrt{T}}{1-m} + \sqrt{\frac{M_0^{1-m}}{2(1-m)}}.$$

Case b). We take $\theta(u) = \lambda u \log \frac{M}{u}$ with $M > M_0$. Then, (32) leads to

$$-\lambda^2 \log \frac{M}{u} + \frac{1}{2} + \frac{\xi'}{\xi^2} \sqrt{t} \lambda \left(\log \frac{M}{u} - 1 \right) + \frac{\xi''}{\xi^3} t \leq 0.$$

Assuming $|\lambda| \geq \sqrt{T}$, then

$$\frac{\xi'}{\xi^2} \sqrt{t} \lambda \leq \lambda^2$$

and the above condition becomes

$$-\lambda^2 \log \frac{M}{u} + \frac{1}{2} + \sqrt{T} |\lambda| \left| \log \frac{M}{u} - 1 \right| + 2T \leq 0.$$

This is satisfied once that

$$|\lambda| \geq \lambda_0 = \frac{1}{\log \frac{M}{u}} \left(\left| \log \frac{M}{u} - 1 \right| \sqrt{T} + \left(T \left(\log \frac{M}{u} - 1 \right)^2 + 4 \log \frac{M}{u} \left(2T + \frac{1}{2} \right) \right)^{1/2} \right).$$

Again, a simple computation shows that

$$\lambda_0 \leq 2\sqrt{T} \left(1 + \frac{1}{\log \frac{M}{u}} \right) + \left(\frac{2}{\log \frac{M}{u}} \right)^{1/2}.$$

Case c). We take $\theta(u) = \lambda(\alpha u - u^m)$ with $\alpha > \alpha_0 := \sup_Q u^{m-1}$. Then, (32) leads to

$$-\lambda^2(m-1)(\alpha - u^{m-1}) + \frac{1}{2} + \lambda \sqrt{t} \frac{\xi'}{\xi^2} (m\alpha - (2m-1)u^{m-1}) + m t \frac{\xi''}{\xi^3} u^{m-1} \leq 0.$$

Assuming $|\lambda| \geq \frac{(2m-1)}{m-1}\sqrt{T}$, then

$$(2m - 1) \frac{\xi'}{\xi^2} \sqrt{t} \lambda \leq (m - 1) \lambda^2$$

and the above condition becomes

$$-\lambda^2(m - 1)(\alpha - \alpha_0) + \frac{1}{2} + |\lambda| \sqrt{T} |m\alpha - (2m - 1)\alpha_0| + 2mT\alpha_0 \leq 0.$$

This condition is satisfied if

$$|\lambda| \geq \lambda_0 = \frac{1}{(m - 1)(\alpha - \alpha_0)} \left(\sqrt{T} |m\alpha - (2m - 1)\alpha_0| + \left(T |m\alpha - (2m - 1)\alpha_0|^2 + 4(m - 1)(\alpha - \alpha_0) \left(\frac{1}{2} + 2mT\alpha_0 \right) \right)^{1/2} \right),$$

and we have

$$\lambda_0 \leq \frac{2\sqrt{T}}{(m - 1)(\alpha - \alpha_0)} (m(\alpha - \alpha_0) + 2(m - 1)\alpha_0) + \left(\frac{2}{(m - 1)(\alpha - \alpha_0)} \right)^{1/2}.$$

□

In the case of homogeneous Dirichlet boundary conditions and for a general maximal monotone graph φ and $\psi \equiv 0$, it is not difficult to show that the conclusions of Corollary 4 and Proposition still hold for mild solutions. This leads to a gradient estimate which implies a regularizing effect stronger than the fact that $\varphi(u(t, \cdot))_x \in L^2(a, b)$, for any $t > 0$, shown by H. Brézis in [16] for initial data $u_0 \in H^{-1}(a, b)$. For the sake of the exposition, let us assume that

$$D(\varphi) = R(\varphi) = \mathbb{R}. \tag{66}$$

Then we have:

COROLLARY 6. *Let φ be a maximal monotone graph of \mathbb{R}^2 with $0 \in \varphi(0)$ and satisfying (66). Let $u_0 \in L^1(a, b)$, $u_0 \geq 0$. Then the mild solution u of problem*

$$\begin{cases} u_t = w_{xx}, w \in \varphi(u) & \text{in } (0, T) \times (a, b), \\ w(t, a) = w(t, b) = 0 & t \in (0, T), \\ u(0, x) = u_0(x) & x \in (a, b), \end{cases}$$

satisfies

$$\|w(t, \cdot)_x\|_{L^\infty(a,b)} \leq 2 \left[\frac{\varphi^+(m(t))m(t)}{2} \right]^{1/2} \tag{67}$$

for any $t > 0$, with $\varphi^+(r) = \max\{s \in \mathbb{R} : s \in \varphi(r)\}$ and

$$m(t) = I^{-1} \left(\frac{\|u_0\|_{L^1(a,b)}^{3/2}}{4t} \right).$$

3.5. Applications to the study of the free boundaries

We shall end the paper with several comments on the connections between pointwise gradient estimates and the study of the free boundaries defined as the boundaries of the support of the solution of the equation (2).

As it was already mentioned in the introduction, in the context of the gas flow in porous media the expression

$$V(t, x) = - \frac{\varphi(u(x, t))_x + \psi(u(x, t))}{u(x, t)}$$

represents the Eulerian description of the velocity of the “homogenized gas” at the position x and the time t once we identify the parabolic equation with the continuity equation for the mass conservation. In particular, if we denote by $x = \zeta(t)$ any of those interfaces, by standard fluid mechanics arguments we deduce, at least formally, that if $\zeta(t)$ is differentiable, then the particles of the gas at the boundary of the support at time $t = 0$ must remain at the boundary of the support of the gas for any $t > 0$ and so they must move with velocity $V(t, \zeta(t))$. Hence if, for instance, $\zeta(t) = \sup\{x \in \mathbb{R} : u(t, x) > 0\}$ then

$$\zeta'(t) = - \frac{\varphi(u(t, \zeta(t)))_x + \psi(u(t, \zeta(t)))}{u(t, \zeta(t))}. \tag{68}$$

The question of justifying the above formula, in a rigorous way, is not an easy task. To start with, the occurrence of the free boundaries must be correctly proved without invoking any physical argument (recall that the equation holds for an homogenized medium). Among many papers dealing with the study of the interfaces we would start by mentioning the short paper by A. S. Kalashnikov [34] in which he shows that the solutions of the Cauchy problem (35) with $\psi \equiv 0$ and $u_0 \geq 0$ with compact support remain with compact support for any $t > 0$ (a property that, in what follows, we shall call as *finite speed of propagation*), provided that the pointwise gradient estimate

$$|\varphi(u)_x| \leq Mu \tag{69}$$

holds. We recall that, by using completely different methods, it was shown that the finite speed of propagation holds if and only if

$$\int_{0^+} \frac{d\varphi(s)}{s} < \infty \tag{70}$$

(see [41] for the sufficient part and [31] and [42] for the necessity part). So, we conclude that estimate (69) only remains true by assuming the condition (70). Although a direct derivation of estimate (69) was already shown in [3] and [32], we mention that this can be deduced also from our Theorem 1 by taking $\lambda(u) \equiv u$. Notice that condition (40) coincides with (70) and that the estimate can be also proved for the class of mild solutions when φ is a maximal monotone graph satisfying (70) since this condition make sense even if φ is not C^1 .

In the case of $\psi \neq 0$ the analysis is more sophisticated since the equation loses its invariance after changing x by $-x$. The convection term privileges one direction in contrast with the other one. The *finite speed of propagation* (i. e. the existence of a minimal and maximal interface) was shown in [30] and [33] when $\varphi(u) = u^m$ and $\psi(u) = u^n$ with $m > 1$ and $n \geq 1$. Nevertheless, it was shown (by means of suitable pointwise gradient estimate) in [19] that if $\varphi(u) = u^m$ and $\psi(u) = -u^n$ with $0 < n < 1 \leq m$ then one of the interfaces ($\zeta_+(t) = \sup\{x \in \mathbb{R} : u(t, x) > 0\}$) does not appear ($\zeta_+(t) = +\infty$ for any $t > 0$) although the other one ($\zeta_-(t) = \inf\{x \in \mathbb{R} : u(t, x) > 0\}$) exists ($-\infty < \zeta_-(t) < +\infty$ for any $t > 0$). Notice that this can be also derived from Theorem 2 since, by a comparison argument, without loss of generality, we can assume M_0 small enough and from estimate (58) we deduce that

$$u(t, x)^m \geq u(t, x_0)^m e^{-\frac{K}{t}(x-x_0)}$$

for any $t > 0$ and $x \geq x_0$. Taking x_0 such that $u(t, x_0) > 0$ we get the result. We point out that the existence of the interface $\zeta_-(t)$ shows the impossibility to have the reverse directional symmetric pointwise gradient estimate

$$(u^m)_x \leq K \frac{u^m}{t} \text{ on } (0, T) \times \mathbb{R}.$$

The occurrence of the interfaces was studied by B. H. Gilding [26], in the general case of φ and ψ as in Theorem 1 He proved that the interface $\zeta_+(t)$ exists if and only if there is a real number σ such that

$$\sigma s + \psi(s) > 0 \text{ for all } s \in (0, M_0]$$

and

$$\int_{0^+} \frac{\varphi'(s)}{\sigma s + \psi(s)} < \infty.$$

Notice that this proves the non existence of $\zeta_+(t)$ if $\varphi(u) = u^m$ and $\psi(u) = -u^n$ with $0 < n < 1 \leq m$ and that, by changing x by $-x$, it proves the existence of $\zeta_-(t)$. It was proved also in [26] that if $V(t, x)$ is bounded for $t = 0$ then it remains bounded for any $t > 0$ assumed (70) and

$$\int_{0^+} \left[\frac{-\psi(s)}{s} \right] ds < \infty.$$

Nevertheless, when $\varphi(u) = u^m$ and $\psi(u) = -u^n$ with $0 < n < 1 \leq m$, it was shown in [20] that the best pointwise gradient estimate has the form

$$|(u^{m-n})_x| \leq M \quad (71)$$

so that estimate (69) fails. Estimate (71) was later extended in [26] to more general φ and ψ but satisfying additional conditions which are satisfied when $\varphi(u) = u^m$, $\psi(u) = -u^n$ and $0 < n < 1 \leq m$ (see also the work [29] for the study of the dependence of the Bernstein estimates on epsilon if one puts such a parameter in front of the diffusion term). We mention that, as kindly indicated in [24] and [26], the boundedness of V and the generalization of the estimate (71) were obtained by applying the general idea of the method presented in this paper (a preliminar version of it circulate, as unpublished notes, among the specialists in the field since the early eighties).

The justification of the differential equation was carried out, when $\varphi(u) = u^m$ and $\psi(u) = -u^n$ in [43] and [21] for the case $0 < n < 1 \leq m$ (see also the formal asymptotic analysis made in [36]). It was shown ([21]) that in absence of *waiting time phenomena* ([2], [1]) $\zeta_- \in C([0, T]) \cap Lip(0, T)$,

$$\lim_{x \rightarrow \zeta_-(t)-0} \frac{\varphi(u(t, x))_x}{\psi(u(t, \zeta(t)))} = -1$$

and that formula (68) holds in the sense that

$$\begin{aligned} \zeta'(t) &= - \lim_{x \rightarrow \zeta_-(t)-0} \frac{\varphi(u(t, x))_x + \psi(u(t, x))}{u(t, x)} \\ &= - \lim_{x \rightarrow \zeta_-(t)-0} \left[\frac{\psi(u(t, x))}{u(t, x)} \right] \left[\frac{\varphi(u(t, x))_x}{\psi(u(t, x))} + 1 \right] \end{aligned}$$

for *a.e.* $t \in [0, T]$. It is easy to see that applying (formally) the l'Hopital rule to the right-hand side of this interface equation we get a second-order equation. The interface equations of this type can also be derived by means of a special technique based on local comparison of the space profile of a solution with a set of the space profiles of a complete family of explicit solutions to the same equation, say, the travelling wave solutions: see [23], Section 5.

$C^{1+\alpha}$ and further regularity of the interface was obtained in [44] (see also [40]) under additional assumptions on the initial datum.

REFERENCES

- [1] ALVAREZ, L. and DÍAZ, J. I., *Sufficient and necessary initial mass conditions for the existence of a waiting time in nonlinear diffusion-convection processes*, Journal of Mathematical Analysis and Applications, 155 (1991), 378–392.
- [2] ALVAREZ, L., DÍAZ, J. I. and KERSNER, R., *On the initial growing of the interfaces in filtration processes, in Nonlinear Diffusion Equations and Their Equilibrium States I*, W. M. Ni, L. A. Peletier and J. Serrin eds.), Springer-Verlag, New York 1988, 1–20.

- [3] ARONSON, D. G., *Regularity properties of flows through porous media*, SIAM J. Appl. Math. 17 (1969), 461–467.
- [4] ARONSON, D. G. and BÉNILAN, PH., Régularité des solutions de l'équation des milieux poreux dans \mathbb{R}^N , C. R. Acad. Sc. Paris, 288 (1979), 103–105.
- [5] BÉNILAN, PH., On $u_t = \varphi(u)_{xx} + \psi(u)_x$, Unpublished 1978.
- [6] BÉNILAN, PH., *Opérateurs diffé rentiales linéaires*, in *Analyse mathématique et calcul numé rique pour les sciences et les techniques*, R. Dautray and J.-L. Lions eds., Masson, Paris 1984, 947–1085.
- [7] BÉNILAN, PH., *Evolution Equations and Accretive Operators*, Lecture Notes (taken by S. Lenhart), Univ. of Kentucky 1981.
- [8] BÉNILAN, PH., Personal Notes, Unpublished 1981.
- [9] BÉNILAN, PH. and BOUILLET, J., *On a slow fast diffusion equation*, Nonlinear Analysis, 26 (1996), 813–822.
- [10] BÉNILAN, PH. and DÍAZ, J. I., *Comparison of solutions of nonlinear evolutions equations with different nonlinear terms*, Israel Journal of Mathematics, 42 (1982), 241–257.
- [11] BÉNILAN, PH. and GARIEPY, R., *Strong solutions in L^1 of degenerate parabolic equations*, Journal of Differential Equations, 119 (1995), 473–502.
- [12] BÉNILAN, PH. and TOURÉ, H., *Sur l'équation générale $u_t = \varphi(\cdot, u)_{xx} - \psi(u)_x + v$* , C.R. Acad.Sc. Paris, 299 (1984), 91–922.
- [13] BÉNILAN, PH. and TOURÉ, H., *Sur l'é quation générale $u_t = a(\cdot, u, \varphi(\cdot, u)_x)_x + v$ dans L^1 : II- Le problème d'évolution*, Ann. Inst. Henri Poincaré: Analyse non linéaire, 12, (1995) 727–761.
- [14] BERNSTEIN, S. N., *Collected works, vol III: Differential equations, Calculus of Variations and Geometry (1903 – 1947)*, Izdat. Akad. Nauk. SSSR, Moscow, 1960, (in russian).
- [15] BERRYMAN, J. G. and HOLLAND, C. J., *Stability of the separable solution for fast diffusion*, Arch. Rational Mech. Anal. 74 (1980), 379–388.
- [16] BRÉZIS, H., *Monotonicity Methods in Hilbert Spaces and Some Applications to Nonlinear Partial Differential Equations*, in *Contributions to Nonlinear Functional Analysis*, E. Zarantonello, ed., Academic Press, New York 1971, 101–156.
- [17] CARRILLO, J., *Entropy solutions for nonlinear degenerate problems*, Arch. Ration. Mech. Anal. 147 (1999), 269–361.
- [18] DÍAZ, J. I., *Two Problems in Homogenization of Porous Media*, Extracta Mathematica, 14 (1999), 141–155.
- [19] DÍAZ, J. I. and KERSNER, R., *Non existence d'une des frontières libres dans une équation dégénérée en theorie de la filtration*, C.R. Acad. Sc. Paris, 296 (1983), 505–508.
- [20] DÍAZ, J. I. and KERSNER, R., *On a nonlinear degenerate parabolic equation in filtration or evaporation through a porous medium*, Journal of Differential Equations, 69 (1987), 368–403.
- [21] DÍAZ, J. I. and SHMAREV, S. I., *On the behaviour of the interface in nonlinear processes with convection dominating diffusion via Lagrangian coordinates*, Advances in Mathematical Sciences and Applications, 1 (1992), 19–45. Corrections 2 (1993), 503–506.
- [22] GAGNEUX, G. and MADAUNE-TORT, M., *Analyse mathématique de modèles non linéaires de l'ingenierie pétrolière. (Mathematical analysis of nonlinear models of petrol engineering)*. Mathematiques & Applications, 22, Springer-Verlag, Paris, 1995.
- [23] GALAKTIONOV, V. A., SHMAREV, S. I. and VAZQUEZ, J. L., *Second-order interface equations for nonlinear diffusion with very strong absorption*, Commun. Contemp. Math., 1 (1999), 51–64.
- [24] GILDING, B. H., *Hölder continuity of solutions of parabolic equations*, J. London Math. Soc. 13 (1976), 103–106.
- [25] GILDING, B. H., *Properties of solutions of an equation in the theory of infiltration*. Arch. Ration. Mech. Anal. 65 (1977), 203–225.
- [26] GILDING, B. H., *The Ocurrance of Interfaces in Nonlinear Diffusion-Advection Processes*. Arch. Ration. Mech. Anal. 100 (1988), 243–263.
- [27] GILDING, B. H., *Improved Theory for a Nonlinear Degenerate Parabolic Equation*. Annali Scu. Norm. Sup. Pisa, Serie IV. 14 (1989), 165–224.
- [28] GILDING, B. H. and GONCERZEWICZ, J., *Localization of solutions of exterior domain problems for the porous media equation with radial symmetry*. SIAM, J. Math. Anal. 31 (2000), 862–893.
- [29] GILDING, B. H., NATALINI, R. and TESI, A., *How parabolic free boundaries approximate hyperbolic fronts*. Trans. Amer. Math. Soc, 352 (2000), 1797–1824.

- [30] GILDING, B. H. and PELETIER, L. A., *The Cauchy problem for an equation in the theory of infiltration*, Arch. Ration. Mech. Anal. *61* (1976), 127–140.
- [31] KALASHNIKOV, A. S., *Equations of the type of a nonstationary filtration with infinite rate of propagation of perturbations*, Vestnik Moskov. Univ. Ser. I Mat. Meh. *27* (1972), 45–49.
- [32] KALASHNIKOV, A. S., *On the differential properties of generalized solutions of equations of the nonsteady-state filtration type*, Vestnik Moskov Univ. Ser. I Mat. Fiz. *29* (1974), 62–68.
- [33] KALASHNIKOV, A. S., *The nature of the propagation of perturbations in processes that can be described by quasilinear degenerate parabolic equations*, Trudy Sem. Petrovsk., *1* (1975), 135–144, (in russian).
- [34] KALASHNIKOV, A. S., *On the concept of finite rate of propagation of perturbations*, Uspekhi Mat. Nauk, *34* (1979), 199–200.
- [35] KALASHNIKOV, A. S., *Some problems of the qualitative theory of second-order nonlinear degenerate parabolic equations*, Uspekhi Mat. Nauk, *42* (1987), 135–176.
- [36] KAMIN, S. and ROSENAU, PH., *Thermal waves in an absorbing and convective medium*, Physica *8D*, (1983) 717–737.
- [37] KRUIZHNIKOV, S. N., *Nonlinear parabolic equations with two independent variables*, Trudi Mosk. Matem. ob. (Proceedings of Moscow Math. Society), *6* (1967), 329–346.
- [38] LADYZHENSKAJA, O. A., SOLONNIKOV, V. A., and URAL'TSEVA, N. N., *Linear and quasilinear equations of parabolic type*, American Mathematical Society, Providence, R.I., 1967.
- [39] LAURENCOT, PH. and SIMONDON, F., *Source-type solutions to porous medium equations with convection*, Comm. in Appl. Anal., *1*, 1997.
- [40] MEIRMANOV, A. M., PUKHINACHOV, V. V. and SHMAREV, S.I., *Evolution Equations and Lagrangian Coordinates*, Walter de Gruyter, Berlin, 1997.
- [41] OLEINIK, O. A., KALASHNIKOV, A.S. and CHZHOU Y.-L., *The Cauchy problem and boundary problems for equations of the type of non-stationary filtration*, Izv. Akad. Nauk SSR, Ser. Mat. *22* (1958), 667–704 (in russian).
- [42] PELETIER, L. A., *A necessary and sufficient condition for the existence of an interface in flows through porous media*, Arch. Ration. Mech. Anal., *56* (1974), 183–190.
- [43] SHMAREV S. I., *Instantaneous appearance of singularities of a solution of a degenerate parabolic equation*, Sib. Math. J., *31* (1990), 671–682.
- [44] SHMAREV, S. I., *Global $C^{1+\alpha}$ regularity of interface for $u_t = (u^m)_{xx} + (u^\lambda)_x$* , Advances in Mathematical Sciences and Applications, *5* (1995), 1–29.
- [45] SIMONDON, F., *Strong solutions for $u_t = (\varphi(u))_{xx} - f(t)(\psi(u))_x$* , Comm. in PDE, *13* (1988), 1337–1354.
- [46] VAZQUEZ, J. L., *An introduction to the mathematical theory of the porous medium equation*, in *Shape Optimization and Free Boundaries*, M. Delfour, ed., Kluwer Acad. Publ., Dordrecht, Netherlands, 1992, 347–389.

Jesús Ildefonso Díaz
 Facultad de Matemáticas
 Universidad Complutense de Madrid
 20040 Madrid
 Spain