



Contents lists available at ScienceDirect

Nonlinear Analysis

journal homepage: www.elsevier.com/locate/na

Free boundaries touching the boundary of the domain for some reaction–diffusion problems

Jesús Ildefonso Díaz^{a,*}, Tommaso Mingazzini^b

^a Instituto de Matemática Interdisciplinar, Universidad Complutense de Madrid, Plaza de las Ciencias 3, 28040 Madrid, Spain

^b Universidad Complutense de Madrid, Dpto. de Matemática Aplicada, Plaza de las Ciencias 3, 28040 Madrid, Spain

ARTICLE INFO

Communicated by Enzo Mitidieri

Keywords:

Semilinear elliptic and parabolic problems

Free boundary

Compact support

Boundary behaviour

ABSTRACT

We give conditions on the behaviour of the trace datum near the boundary of its support in order to know whether the free boundary given by the boundary of the support of the solution of suitable elliptic or parabolic semilinear problem is connected or not with the boundary of the support of the boundary datum.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

We study the way the free boundary of solutions to some partial differential equations behaves depending on the trace of the solutions. The free boundary problems we consider are of two different types:

(i) *Elliptic reaction–diffusion type problems*, as

$$\begin{cases} -Lu + \lambda u^q = 0 & \text{in } \Omega, \\ u = h & \text{on } \partial\Omega, \end{cases} \quad (1)$$

under the fundamental assumption

$$q \in (0, 1), \quad (2)$$

which guarantees the formation of the free boundary (at least for $\lambda > 0$ large enough, if Ω is bounded, or for any $\lambda > 0$, if Ω is unbounded). Such problem arises, for instance, in Chemical Engineering when a catalytic chemical reactor occupying a domain Ω has a reactant feed channel (entrance boundary) which is represented by the part $\Gamma_+ \subset \partial\Omega$, where the reactant concentration is $h(x) > 0$ and the rest of walls of the chemical reactor are isolated in such a way that, if we denote by $\Gamma_0 := \partial\Omega \setminus \Gamma_+$, then $h(x) = 0$ on Γ_0 . Here we assume that there is no exit boundary (see Fig. 1). The exponent q is called the order of the reaction.

(ii) *The obstacle problem*

$$\begin{cases} -Lu \geq f(x), & u \geq 0 \quad \text{and} \quad (-Lu - f(x))u = 0 & \text{in } \Omega, \\ u(x) = h(x) & & \text{on } \partial\Omega. \end{cases} \quad (3)$$

Here the free boundary is given by the boundary of the *coincidence set* (the set of points where $u = 0$); according for instance to [17] a sufficient condition for the existence of the free boundary is that $f(x) \leq -\mu$ for some $\mu > 0$ on a large enough

* Corresponding author.

E-mail addresses: ildefonso.diaz@mat.ucm.es (J.I. Díaz), tommaso.mingazzini@gmail.com (T. Mingazzini).




Fig. 1. Chemical reactor scheme.

open subset of Ω (see, for instance, [28] for a full treatment of the obstacle problem). Among the many frameworks in which the obstacle problem arises we could mention, for instance, the unilateral problem of the stationary shape of a membrane which is forced downwards by a constant force f , is fixed on the boundary to a height $h(x)$ and constrained to lie over the hyperplane $u = 0$. Actually, here we shall consider the special case in which (3) can be formulated in terms of

$$\begin{cases} -Lu + \lambda\beta(u) \ni \varepsilon & \text{in } \Omega, \\ u = h & \text{on } \partial\Omega, \end{cases} \tag{4}$$

for some constant $\varepsilon \in [0, \lambda)$, where $\beta(u)$ is the maximal monotone graph of \mathbb{R}^2 given by

$$\beta(u) = \begin{cases} 0 & \text{for } u < 0, \\ [0, 1] & \text{for } u = 0, \\ 1 & \text{for } u > 0. \end{cases} \tag{5}$$

If u “solves” problem (4) (the rigorous definition of solution will be given later) then u is also a solution of the obstacle problem (3) with $f = -\lambda + \varepsilon$: indeed, we will see that $\varepsilon \geq 0$ and $h \geq 0$ imply that $u \geq 0$. Then, if $u > 0$, $-Lu + \lambda = \varepsilon$ which is the same as $-Lu - f = 0$. Finally, since there is uniqueness of solution for both formulations we get that the solutions must be the same.

Another interesting application of problem (4) arises also in the context of Chemical Engineering (as problem (1) with $q = 0$: see, e.g., [7]).

For some general purposes, such as the existence, uniqueness and regularity of the solutions, the domain Ω will be assumed to be an open regular set of \mathbb{R}^N . Nevertheless, when studying the qualitative properties of the solutions we focus on the bi-dimensional case, and we adopt as domain Ω both a bounded rectangle and the upper half plain in \mathbb{R}^2 , i.e., $\Omega = \mathbb{R} \times [0, \infty)$. In the unbounded setting we use the following notation: $x := (x_1, x_2)$ with $x_1 \in \mathbb{R}$ and $x_2 \in [0, \infty)$. The unbounded boundary of the domain is then $\partial\Omega = \mathbb{R} \times \{0\}$ and so the boundary function h will depend only on the variable x_1 .

In general, L denotes a second order elliptic operator of the form

$$Lu = \sum_{i,j=1}^N \frac{\partial}{\partial x_i} \left(a_{ij}(x) \frac{\partial}{\partial x_j} u \right) = \text{div}(\mathbf{A}(x)\nabla u), \tag{6}$$

with $a_{ij} \in C^{1,\alpha}(\overline{\Omega})$ for some $\alpha \in (0, 1)$, such that the corresponding matrix $\mathbf{A}(x)$ is symmetric and positive definite. Actually, in the parts concerning the behaviour of the support and free boundary of the solutions we shall restrict to the case of constant coefficients. This restriction serves merely to simplify the calculations and does not affect the local behaviour. For what concerns the boundary datum h , we assume that

$$h \in L^\infty(\partial\Omega) \text{ and } h \geq 0 \text{ on } \partial\Omega,$$

even though the existence and uniqueness results on a bounded domain hold for $h \in L^1(\partial\Omega)$ (and even for signed boundary measures).

A general exposition containing many references on both problems can be found in the monograph [17]. One can see that both problems are special cases of the wider formulation

$$\begin{cases} -Lu + \lambda\beta(u) \ni f & \text{in } \Omega, \\ u = h & \text{on } \partial\Omega, \end{cases} \tag{7}$$

where $\beta(u)$ is a maximal monotone graph of \mathbb{R}^2 such that $0 \in \beta(0)$: β is given by

$$\beta(u) = |u|^{q-1} u \tag{8}$$

in case of problem (1) and by (5) in case of problem (3). We define, as usual, the domain of β as $D(\beta) = \{r \in \mathbb{R} : \beta(r) \neq \emptyset\}$ where \emptyset stands for the empty set.

We also consider the associated parabolic problem

$$\begin{cases} u_t - Lu + \lambda\beta(u) \ni f(x, t) & \text{in } Q_\infty, \\ u = h(t, x) & \text{on } \Sigma_\infty, \\ u(x, 0) = u_0(x) & \text{on } \Omega, \end{cases} \tag{9}$$

where $Q_\infty = \Omega \times (0, \infty)$, $\Sigma_\infty = \partial\Omega \times (0, \infty)$ and for some $f \in L^\infty(Q_\infty) \cap L^1_{loc}((0, \infty); L^1_{loc}(\Omega))$, $h \in L^\infty(\Sigma_\infty) \cap L^1_{loc}((0, +\infty); L^1(\partial\Omega))$, with $f, h \geq 0$ respectively on Q_∞ and on Σ_∞ , and $u_0 \in L^\infty(\Omega)$ with $u_0 \geq 0$ on Ω .

As mentioned before, the above problems, both elliptic and parabolic, give rise to a free boundary defined as the boundary of the support of the solution. If we denote the positivity set of a non-negative function u by $\mathcal{S}(u) := \{x \in \Omega : u(x) > 0\}$, then the free boundary is defined as $\mathcal{F}(u) = \partial\mathcal{S} \cap \Omega$ (we also introduce the null set of u as $\mathcal{N}(u) := \{x \in \Omega : u(x) = 0\}$ and the support of u as $\overline{\mathcal{S}(u)}$). Similar notations can be introduced also for the parabolic problem, applying the definitions to $u(t, \cdot)$.

Our main goal in this work is to study the behaviour of the free boundary near the support of the boundary datum h (respectively $h(t, \cdot)$). For this purpose, we shall assume that

$$\mathcal{S}(h) \subsetneq \partial\Omega,$$

respectively

$$\mathcal{S}(h(t, \cdot)) \subsetneq \partial\Omega, \quad \text{for a.e. } t > 0.$$

The main question we investigate is whether the free boundary $\mathcal{F}(u)$ is connected or not with the boundary of the support of the boundary datum h (and similar question for the parabolic formulation). In some sense, this research can be considered as a natural continuation of the study of the so called *non-diffusion of the support property* (see [17,2]) in the case where $h \equiv 0$; under a suitable behaviour of f near the boundary of its support $\mathcal{S}(u) = \mathcal{S}(f)$. In the case of parabolic free boundary problems this question is related with the behaviour of the free boundary for small times (the so called *waiting time property*) and received a great attention in the last 40 years (see, e.g., the monographs [31,3] and their many references). Another study, not too far from our interest is the paper by Martel and Souplet [26] regarding the behaviour of solutions of linear parabolic problems with incompatible initial data.

To be more precise, our main goal is to find some sufficient criterion on the behaviour of h near the boundary of its support ensuring that the free boundary $\mathcal{F}(u)$ is in contact with $\partial\mathcal{S}(h)$. In this way the support of the datum is not diffused on the boundary of the domain and we would have

$$\partial\mathcal{S}(u) \cap \partial\Omega = \mathcal{S}(h). \tag{10}$$

It is what we can call the *non-diffusion on the boundary of the support property*. In addition, we want to give some sufficient conditions ensuring the opposite qualitative behaviour, i.e., to find conditions on h implying that there is a strict expansion of the support $\mathcal{S}(h)$ on the boundary $\partial\Omega$. In other words, we want to know cases in which $\mathcal{F}(u)$ has no contact with $\partial\mathcal{S}(h)$ and so

$$\mathcal{S}(h) \subsetneq \partial\mathcal{S}(u) \cap \partial\Omega.$$

We call this phenomenon the *expansion on the boundary of the support property*. The only paper in the previous literature about such boundary qualitative behaviour we are aware of is [30] in which they proved the *expansion on the boundary of the support property* for problem (1) in the special case of $Lu = \Delta u$, h given by the Heaviside function and Ω the half plane $\mathbb{R} \times \mathbb{R}^+$. As we shall see later, this property holds even for suitable continuous boundary data h .

We point out that there is an extensive literature dealing with the regularity of the free boundary when it touches the fixed boundary for the special case of the obstacle problem (both in the elliptic and parabolic case): see, e.g. [4–6,29]. In some of those papers the authors also consider more general diffusion operators (arriving to consider even the case of fully nonlinear second order operators). Nevertheless, the reaction term here involved, $\beta(u)$, is never similar to the one considered in those papers except for the case of zero-order reaction. In any case, our main interest is not the regularity of the free boundary but its qualitative behaviour depending on the data of the problem (specially the behaviour of the boundary datum near its support). To stretch the value of our results, we underline that the techniques of proof in this paper are quite different to the ones used in the above mentioned papers on the obstacle problem.

Before stating our main results we need to make precise the notion of solution. The delicate point in our study is that we want to allow the boundary datum to be discontinuous and so the notion of the trace of the solution must be taken in a very general framework (something which, in our opinion, is not discussed enough in [30]).

We recall that the notion of boundary trace of a function u in Ω depends on the regularity properties of such function u . For instance, when $u \in C(\overline{\Omega})$ the boundary trace $u|_{\partial\Omega}$ is clearly well defined and belongs to $C(\partial\Omega)$. If u is in some

Sobolev space $W^{1,p}(\Omega)$, for some $p > 1$, then the boundary trace can be defined and is a function in the space $L^p(\partial\Omega)$ (more precisely in the Sobolev space $W^{1-\frac{1}{p},p}(\partial\Omega)$: see, e.g., [22,1]). Nevertheless, the identification of the elements of the trace space $W^{1-\frac{1}{p},p}(\partial\Omega)$ is not always easy and leads to some pathological results against intuition. For instance in the book by Mikhailov [27] one can see that already when $\Omega = B$, the unit ball of \mathbb{R}^2 , there are continuous functions $h \in C(\partial B)$ which are not the trace of any function in $H^1(\Omega)$ (i.e., $C(\partial B) \not\subseteq H^{\frac{1}{2}}(\partial B)$).

A different approach was proposed by Haïm Brezis, in an unpublished paper (1972) profusely mentioned in the literature (see [32,25,19]), which holds for semilinear second order boundary value problems with boundary data in $L^1(\partial\Omega)$ (later extended to measures on $\partial\Omega$). The main idea is to multiply by a “regular” test function ($\varphi \in W^{2,\infty}(\Omega) \cap W_0^{1,\infty}(\Omega)$) and to integrate twice by parts. We introduce the adjoint operator

$$L^*u = \sum_{i,j=1}^N \frac{\partial}{\partial x_j} \left(a_{ij}(x) \frac{\partial}{\partial x_i} u \right) = \operatorname{div}(\mathbf{A}^* \nabla u)$$

(\mathbf{A}^* the transposed matrix of \mathbf{A}) and for $x \in \partial\Omega$ we define

$$\partial_A u := (\mathbf{A}^* \nabla u) \cdot \mathbf{n},$$

where $\mathbf{n}(x)$ is the outward normal vector to $\partial\Omega$ in x . A solution is then a function u which satisfies

$$-\int_{\Omega} u L^* \varphi \, dx + \lambda \int_{\Omega} b \varphi \, dx = \int_{\Omega} f \varphi \, dx - \int_{\partial\Omega} h \partial_A \varphi \, d\sigma, \tag{11}$$

for all $\varphi \in W^{2,\infty}(\Omega) \cap W_0^{1,\infty}(\Omega)$ and for some $b \in L^1_{\text{loc}}(\Omega)$ such that $b(x) \in \beta(u(x))$ for a.e. $x \in \Omega$ (in case of problem (1), it is $b = u^q$). In order to give a meaning to all the above integrals, it is useful to recall that, since $\varphi \in W_0^{1,\infty}(\Omega)$, it holds $c_1 \rho(x) \leq \varphi(x) \leq c_2 \rho(x) \, \forall x \in \Omega$, where

$$\rho(x) := \operatorname{dist}(x, \partial\Omega)$$

and c_1, c_2 are positive constants. Thus we must require at least that

$$f \in L^1(\Omega; \rho),$$

where

$$L^1(\Omega; \rho) := \left\{ f \in L^1_{\text{loc}}(\Omega) : \int_{\Omega} |f(x)| \rho(x) \, dx < +\infty \right\}.$$

1.1. Elliptic case

Definition 1.1. Given $f \in L^1(\Omega; \rho)$ and $h \in L^1(\partial\Omega)$, we say that u is a very weak solution of problem (7) if $u \in L^1(\Omega)$ and there exists $b \in L^1(\Omega; \rho)$ such that $b(x) \in \beta(u(x))$ for a.e. $x \in \Omega$, and for any test function $\varphi \in W^{2,\infty}(\Omega) \cap W_0^{1,\infty}(\Omega)$ identity (11) holds.

It is not too difficult to adapt to our framework some existence and uniqueness results in the literature (see [12,32,25]).

Theorem 1.2. Let Ω be a bounded regular open set of \mathbb{R}^N , let β be a maximal monotone graph of \mathbb{R}^2 such that $0 \in \beta(0)$ and let $f \in L^1(\Omega; \rho)$ and $h \in L^1(\partial\Omega)$. Then there exists a unique very weak solution u of problem (7). Moreover, there exists a constant C , only dependent of Ω , such that if \widehat{u} is the very weak solution corresponding to the data $\widehat{f} \in L^1(\Omega; \rho)$ and $\widehat{h} \in L^1(\partial\Omega)$, with $\widehat{b}(x) \in \beta(\widehat{u}(x))$ for a.e. $x \in \Omega$ as in Definition 1.1, then we have

$$\| [u - \widehat{u}]_+ \|_{L^1(\Omega)} + \lambda \| [b - \widehat{b}]_+ \|_{L^1(\Omega; \rho)} \leq C \left(\| [f - \widehat{f}]_+ \|_{L^1(\Omega; \rho)} + \| [h - \widehat{h}]_+ \|_{L^1(\partial\Omega)} \right) \tag{12}$$

and

$$\| u - \widehat{u} \|_{L^1(\Omega)} + \lambda \| b - \widehat{b} \|_{L^1(\Omega; \rho)} \leq C \left(\| f - \widehat{f} \|_{L^1(\Omega; \rho)} + \| h - \widehat{h} \|_{L^1(\partial\Omega)} \right). \tag{13}$$

In particular, $f \leq \widehat{f}$ and $h \leq \widehat{h}$ imply that $u \leq \widehat{u}$ on Ω .

To study the behaviour of the solution close to the boundary of $\mathcal{S}(h)$ we consider two different frames, an unbounded case and a bounded one.

In the first case we set $\Omega = \mathbb{R} \times [0, \infty)$, a_{ij} is constant for $i, j \in \{1, 2\}$, $f \equiv 0$, and $\beta(u)$ is given by (8) or (5). For what concerns the boundary datum, we are interested in the case of h satisfying, $h \in L^\infty(\partial\Omega)$, $h(x_1) = 0$ on $(-\infty, 0)$ and $h(x_1) > 0$ on $(0, +\infty)$.

The reason why we consider boundary data in $L^\infty(\partial\Omega)$ instead of in $L^1(\partial\Omega)$ (remember that now $\partial\Omega$ is unbounded, so $L^\infty(\partial\Omega) \not\subseteq L^1(\partial\Omega)$) is that we know the explicit solution in the unperturbed linear case ($\lambda = 0, L = \Delta, f \equiv 0$) with




Fig. 2. Expansion on the boundary of the support.

boundary data given by the Heaviside function (52). Such solution is given by

$$u(x_1, x_2) = 1 - \frac{1}{\pi} \arctan\left(\frac{x_1}{x_2}\right) \tag{14}$$

(the result can be found in [30, formula (2.6)]). Having at disposal an explicit solution like (14) is really useful in the study of the behaviour of general solutions close to the point $x = (0, 0)$. In addition, since our main interest, as already said, is specifically the behaviour near the boundary of the support $\partial \mathcal{S}(h)$ and not in the whole Ω , we shall assume also that h is non decreasing and that $h(x_1) = c_+ > 0$ for $x_1 \geq \delta > 0$. We can resume this set of hypothesis in

$$\begin{cases} \Omega = \mathbb{R} \times [0, \infty), & a_{ij} \text{ constant for } i, j \in \{1, 2\}, & f \equiv 0, \\ \beta(u) \text{ is given by (8) or (5),} \\ h \in L^\infty(\partial\Omega), & h(x_1) = 0 \text{ on } (-\infty, 0), \\ h(x_1) > 0 \text{ on } (0, \infty) \text{ and } h(x_1) = c_+ > 0 \text{ on } (\delta, +\infty). \end{cases} \tag{H_{hp}}$$

To give a definition of solution for the new setting H_{hp} , we first introduce the family of rectangles

$$R_n = \{(x_1, x_2) : |x_1| < n, 0 < x_2 < \bar{x}\},$$

where \bar{x} is a constant which will be made explicit later on. We define now the bounded domain Ω_n , which is a sufficiently smooth regularization of the rectangular R_n . We call Γ_n^1 the horizontal boundary of Ω_n , i.e., $\Gamma_n^1 := \{(x_1, x_2) \in \partial\Omega_n : x_2 = 0 \text{ or } x_2 = \bar{x}\}$, and $\Gamma_n^2 = \partial\Omega_n \setminus \Gamma_n^1$. Then we consider the problem

$$\begin{cases} -Lu + \beta(u) \ni 0 & \text{in } \Omega_n, \\ u(x_1, 0) = h(x_1), & u(x_1, \bar{x}) = 0, & |x_1| \leq n, \\ u(x_1, x_2) = 0 & (x_1, x_2) \in \Gamma_n^2, & x_1 < 0, \\ u(x_1, x_2) = z(x_2) & (x_1, x_2) \in \Gamma_n^2, & x_1 > 0, \end{cases} \tag{15}$$

where z is given by (51). We define the class of “limit very weak solutions”.

Definition 1.3. In the framework of H_{hp} , we say that u is a limit very weak solution of problem (7) if $u = \lim_{n \rightarrow \infty} u_n$, where u_n is the solution of the truncated problem (15).

Theorem 1.4. Assume (H_{hp}) . There exists a unique limit very weak solution of problem (7) on the hyperplane $\Omega = \mathbb{R} \times [0, \infty)$. Moreover that solution satisfies the comparison principle with respect to the boundary data h : if $h \leq \hat{h}$ then the corresponding limit very weak solutions satisfy $u \leq \hat{u}$ on Ω .

Our main result concerns the qualitative behaviour of the solution of (7) under the assumption (H_{hp}) .

Theorem 1.5. Assume (H_{hp}) . Then there exist four positive constants $\underline{C} < \bar{C}$, $\underline{\varepsilon} < \bar{\varepsilon}$ and two boundary points $\underline{x}_{1,\varepsilon}, \bar{x}_{1,\varepsilon} > 0$, such that:

- (i) If $h(x_1) \geq \bar{C}x_1^{\frac{2}{1-q}}$ for a.e. $x_1 \in (0, \bar{x}_{1,\varepsilon})$ and $h(x) \geq \bar{\varepsilon}$ for a.e. $x_1 \in (\bar{x}_{1,\varepsilon}, +\infty)$, then the expansion on the boundary of the support property holds.
 - (ii) If $h(x_1) \leq \underline{C}x_1^{\frac{2}{1-q}}$ for a.e. $x_1 \in (0, \underline{x}_{1,\varepsilon})$ and $h(x_1) \leq \underline{\varepsilon}$ for a.e. $x_1 \in (\underline{x}_{1,\varepsilon}, +\infty)$, then the non-diffusion on the boundary of the support property holds.
- In both cases, $q \in (0, 1)$ when β is given by (8) and $q = 0$ when β is chosen as (5).

The indicative qualitative behaviour of solutions illustrated in Theorem 1.5 is resumed in Figs. 2(i) and 3(ii).




Fig. 3. Non-diffusion on the boundary of the support.

Corollary 1.6. *In the same framework of Theorem 1.5 we have the additional consequences:*

- (1) Under condition (i), $u = \frac{\partial}{\partial n} u = 0$ on $(-\infty, 0) \times \{0\}$.
- (2) Under condition (ii), $\frac{\partial}{\partial n} u > 0$ on $(-\zeta, 0) \times \{0\}$ for some $\zeta > 0$. If in particular h is a multiple of the Heaviside function, then $\frac{\partial}{\partial n} u \notin C(\partial\Omega)$ and $u \notin C(\overline{\Omega})$.

In both situations, $n = (0, 1)$ is the normal vector to $\partial\Omega$.

For the bounded case, we set $\Omega \subset \mathbb{R}^2$ to be the rectangle of vertexes

$$R = \{(-B_l, 0), (B_r, 0), (B_r, B_u), (-B_l, B_u)\} \quad \text{with } B_l, B_r, B_u > 0.$$

Let us call

$$\begin{aligned} \partial_1 R &= [(-B_l, 0), (B_r, 0)], & \partial_2 R &= [(B_r, 0), (B_r, B_u)], \\ \partial_3 R &= [(B_r, B_u), (-B_l, B_u)], & \partial_4 R &= [(-B_l, B_u), (-B_l, 0)], \end{aligned}$$

the four sides of R . The internal datum f is set to zero. We have the following result.

Theorem 1.7. *Assume that*

$$h = \begin{cases} \varepsilon H(x_1) & \text{on } \partial_1 R, \\ \max(0, \psi(x_2)) & \text{on } \partial_2 R, \\ 0 & \text{on } \partial_3 R \cup \partial_4 R, \end{cases}$$

with $H(x_1) = 0$ for $x_1 < 0$, $H(x_1) = 1$ for $x_1 \geq 0$ and

$$\psi(x_2) = \varepsilon \left(\frac{1}{2} + \frac{1}{\pi} \arctan \left((\det A)^{-1} \left(a_{22} \frac{B_r}{x_2} - a_{12} \right) \right) \right) + \frac{\lambda \varepsilon^q}{2} x_2^2 + C x_2 \tag{16}$$

where C is a negative constant. If

$$C \leq \min \left(-\frac{\varepsilon}{B_u} - \frac{\lambda \varepsilon^q}{2} B_u, -\frac{\varepsilon (\det A)^{1/2}}{\pi a_{22} B_l} \right),$$

the solution of (7) with boundary datum h and internal datum $f \equiv 0$ satisfies the expansion on the boundary of the support property. In both cases, $q \in (0, 1)$ when β is given by (8) and $q = 0$ when β is chosen as (5).

Proof. First of all we want to point out that since the boundary conditions are all non-negative and bounded by ε , then $0 \leq u \leq \varepsilon$.

We consider the subsolution given by $\underline{u} = u_1 + u_2$, where

$$u_1 = \varepsilon \left(\frac{1}{2} + \frac{1}{\pi} \arctan \left((\det A)^{-1/2} \left(a_{22} \frac{x_1}{x_2} - a_{12} \right) \right) \right), \quad u_2 = \frac{\lambda \varepsilon^q}{2} x_2^2 + C x_2. \tag{17}$$

One can easily check that $Lu_1 = 0$ and $Lu_2 = \lambda \varepsilon^q$ and as a consequence

$$-Lu = -\lambda u^q \geq -\lambda \varepsilon^q = -L\underline{u}.$$

Now let us check the boundary conditions for \underline{u} . On $\partial_1 R$ we have that $\underline{u}(x_1) = \varepsilon H(x_1)$. On $\partial_2 R$ it is exactly $\psi(x_2)$. On $\partial_3 R$, thanks to the hypothesis on C , it holds

$$\underline{u}(x_1, B_u) \leq \varepsilon + \frac{\lambda \varepsilon^q}{2} B_u^2 + C B_u \leq 0.$$

On $\partial_4 R$, $\underline{u}(-B_I, 0) = 0$. We compute the x_2 -derivative

$$\frac{\partial \underline{u}}{\partial x_2}(x_1, x_2) = \lambda \varepsilon^q x_2 + C - \frac{\varepsilon}{\pi} \frac{a_{22}(\det A)^{1/2} x_1}{\det A x_2^2 + (a_{22}x_1 - a_{12}x_2)^2}.$$

With a direct computation and thanks to the condition on C , $\partial \underline{u} / \partial x_2(-B_I, 0) < 0$ and $\partial \underline{u} / \partial x_2(-B_I, x_1)$ is an increasing function. Hence, since $\underline{u}(-B_I, B_U) \leq 0$, $\underline{u}|_{\partial_4 R} \leq 0$. We have proved that $\underline{u} \leq u$.

Now let us check that the expansion on the boundary of the support holds for \underline{u} and as a consequence for u . We know that $\underline{u}(x_1, 0) = 0$ for $x_1 < 0$. The derivative

$$\frac{\partial \underline{u}}{\partial x_2}(x_1, 0) = C - \frac{\varepsilon \det A}{\pi a_{22} x_1} < 0$$

for $x_1 > \frac{\varepsilon \det A}{\pi a_{22} C} = -\eta$. It follows that $u > 0$ in a neighbourhood in \mathbb{R}^2 of $(-\eta, 0) \times \{0\}$. \square

Corollary 1.8. *The same result holds under the same conditions of the previous theorem but with $u|_{\partial_2 R} \geq \psi$.*

1.2. Parabolic case

As far as the parabolic problem is concerned, our main interest consists in analysing the stabilization of the solution to the solution of stationary problem in order to well understand the *expansion on the boundary of the support* property. When Ω is a general open bounded set the notion of very weak solution is quite similar to the elliptic case (see Fig. 4).

Definition 1.9. Take $T > 0$, $f \in L^1(0, T; L^1(\Omega; \rho))$, $h \in L^1(0, T; L^1(\partial\Omega))$ and $u_0 \in L^1(\Omega; \rho)$ with $u_0(x) \in \overline{D(\beta)}$. We say that u is a very weak solution of problem (9) if $u \in L^1(0, T; L^1(\Omega))$ and there exists $b \in L^1(0, T; L^1(\Omega; \rho))$ such that $b(t, x) \in \beta(u(t, x))$ for a.e. $(t, x) \in (0, T) \times \Omega$, and for every test function $\varphi \in W^{1,\infty}([0, T]; L^\infty(\Omega)) \cap L^\infty(0, T; W^{2,\infty}(\Omega) \cap W_0^{1,\infty}(\Omega))$ with $\varphi(T, \cdot) = 0$ the following identity holds

$$\begin{aligned} & - \int_0^T \int_\Omega u \frac{\partial \varphi}{\partial t} dxdt + \int_0^T \int_\Omega u L^* \varphi dx + \lambda \int_0^T \int_\Omega b \varphi dxdt \\ & = \int_\Omega u_0(x) \varphi(0, x) dx + \int_0^T \int_\Omega f \varphi dxdt - \int_0^T \int_{\partial\Omega} h \partial_A \varphi d\sigma dt. \end{aligned} \tag{18}$$

Once again, it is not too difficult to adapt to our framework some existence and uniqueness results in the literature (see [23]).

Theorem 1.10. (i) *For data f , h and u_0 as in Definition 1.9 there exists a unique very weak solution of (9). Moreover there holds a smoothing effect (same as [23]).*

(ii) *If in addition $h \in W^{1,1}(0, T; L^1(\partial\Omega))$, then the very weak solution satisfies $u \in C([0, T]; L^1(\Omega; \rho))$.*

Our result on the asymptotic behaviour, for $t \rightarrow +\infty$, seems to be new in the context of very weak solutions (check [18,21] for similar results on more regular solutions).

Remark 1.11. Whenever we are dealing at the same time with the parabolic and the elliptic problem, as we are going to do, we use the symbols $u_\infty, h_\infty, f_\infty$ to denote the solution and the data of the elliptic boundary value problem.

For the next result we will add the following hypothesis:

$$\exists q \in [0, 1) \text{ such that } |b| \leq C |r|^q \quad \text{for any } b \in \beta(r) \text{ and for any } r \in \mathbb{R}. \tag{19}$$

In the above condition the case $q = 0$ means that $R(\beta)$ (the range of β , i.e., $r \in \mathbb{R}$ such that there exists $x \in \mathbb{R}$ for which $r \in \beta(x)$) is bounded.

Theorem 1.12. *Consider the case of β satisfying condition (19). Assume $h \in W^{1,1}(0, T; L^1(\partial\Omega))$ for any $T > 0$ and that there exists a sequence $t_n \rightarrow +\infty$, as $n \rightarrow +\infty$, such that*

$$\int_{t_n-1}^{t_n+1} \int_\Omega |f(s, x) - f_\infty(x)| \rho(x) dxds \rightarrow 0 \quad \text{as } n \rightarrow +\infty \tag{20}$$

and

$$\int_{t_n-1}^{t_n+1} \int_{\partial\Omega} |h(s, x) - h_\infty(x)| d\sigma ds \rightarrow 0 \quad \text{as } n \rightarrow +\infty. \tag{21}$$




Fig. 4. Convergence of $\mathcal{S}(u(t, \cdot))$ to $\mathcal{S}(u_\infty)$.

Assume in addition that

$$u \in L^\infty(0, \infty; L^1(\Omega; \rho)). \tag{22}$$

Then $u(t_n, \cdot) \rightarrow u_\infty$ in $L^1(\Omega; \rho)$ with u_∞ the very weak solution of (7) with data f_∞ and h_∞ .

For the qualitative behaviour of the solutions, we consider now the half plane case under the assumptions

$$\begin{cases} \Omega = \mathbb{R} \times [0, \infty), a_{ij} \text{ are constants, } f \equiv 0, u_0 \geq 0, \\ \beta(u) \text{ is given by (8) or (5),} \\ h \in W_{loc}^{1,1}(0, +\infty; L^1(\partial\Omega)) \cap L^\infty((0, +\infty) \times \partial\Omega), \\ h(t, \cdot) = 0 \text{ on } (-\infty, 0) \text{ and } h(t, \cdot) > 0 \text{ on } (0, +\infty), \\ h(t, x_1) = c_+ \quad \forall x_1 \in [\delta, \infty), t > 0. \end{cases} \tag{\widehat{H}_{hp}}$$

In this setting we adapt the definition of limit very weak solution from the elliptic case using truncated-in-space solutions. Existence and uniqueness can be obtained in similar way.

Corollary 1.13. Assume (\widehat{H}_{hp}) . Then there exist four positive constants $\underline{C} < \bar{C}$, $\underline{\varepsilon} \leq \bar{\varepsilon}$ and two boundary points $x_{1,\varepsilon}, \bar{x}_{1,\varepsilon} > 0$ such that

- (i) if $u_0 = 0$, $h(t, x_1) \leq \underline{C}x_1^{\frac{2}{1-q}}$ for a.e. $x_1 \in (0, x_{1,\varepsilon})$ and $h(t, x_1) \leq \underline{\varepsilon}$ for a.e. $x_1 \in (x_{1,\varepsilon}, +\infty)$, for any $t \geq 0$, then the non-diffusion on the boundary of the support property holds for any $t \geq 0$, i.e., $\partial \mathcal{S}(u(t, \cdot)) \cap \partial\Omega = \mathcal{S}(h(t, \cdot))$ for any $t \geq 0$ (infinite waiting time property).
- (ii) Assume that $u_0(x_1, x_2) \geq u_1(x_1, x_2) + u_2(x_1, x_2)$, with u_1 and u_2 solutions of the problems (57) and (58), and that $h(t, x_1) \geq \bar{C}x_1^{\frac{2}{1-q}}$ for a.e. $x_1 \in (0, \bar{x}_{1,\varepsilon})$ and $h(t, x_1) \geq \bar{\varepsilon}$ for a.e. $x_1 \in (\bar{x}_{1,\varepsilon}, +\infty)$, for any $t \in (0, T)$.

Then the expansion on the boundary of the support property holds for any $t \in (0, T]$, i.e., $\mathcal{S}(h(t, \cdot)) \subsetneq \partial \mathcal{S}(u(t, \cdot)) \cap \partial\Omega := [-\delta_0, \infty] \times \{0\}$, for some $\delta_0 > 0$.

Remember that $q \in (0, 1)$ when $\beta(u)$ is given by (8) and $q = 0$ when β is given by (5).

Corollary 1.14. The conclusions of Corollary 1.6 remain valid for $u(t, \cdot)$ under the corresponding assumptions.

Remark 1.15. For the special case of $q = 0$ it is illustrative to compare the conclusions of Theorem 1.5 (respectively Corollary 1.13) with the complementary information given in Theorem B of [29] (respectively Theorem I of [4]).

The organization of the rest of the paper is the following. Section 2 is devoted to the proof of the general existence and uniqueness results, Theorems 1.2 and 1.10. The stabilization of very weak solutions, when $t \rightarrow +\infty$, is considered in Section 3 and, in particular Theorem 1.12 is proved there. Finally the special case of the half plane is considered in Section 4. After proving Theorem 1.4 we present the proof of Theorem 1.5 in Sections 4.2 and 4.3. The special case of discontinuous boundary data plays an important role in such proof and so it is previously discussed there.

2. On the existence and uniqueness of very weak solutions

In this part we omit to write the parameter λ as it can be thought as already included in β .

2.1. Proof of Theorem 1.2

We need to introduce first a result on the corresponding linear problem

$$\begin{cases} -Lu = f & \text{in } \Omega, \\ u = h & \text{on } \partial\Omega. \end{cases} \tag{23}$$

Definition 2.1. Assume $f \in L^1(\Omega; \rho)$ and $h \in L^1(\partial\Omega)$. A function $u \in L^1(\Omega)$ is a weak solution of (23) if it satisfies

$$\int_{\Omega} uL^*\phi \, dx = \int_{\partial\Omega} h \partial_A\phi \, d\sigma - \int_{\Omega} f\phi \, dx$$

for every function $\phi \in C_0^2(\bar{\Omega})$.

The following lemma is a known result and we cite it for further needs.

Lemma 2.2. Assume that u solves (23) with $h = 0$ and $f \in L^2(\Omega)$. Then $u \in H_0^1(\Omega)$ and

$$\int_{\Omega} -Lu \cdot u \, dx \geq C\|u\|_{L^2(\Omega)}^2.$$

Next proposition is a consequence of a generalization of the estimates of Brezis [12] applied to the operator L . The proof when L is the Laplacian can be found in [25]. The case of an even more general second order linear operator of the form

$$Lu = -\text{div}(\mathbf{A}\nabla u) + \mathbf{b} \cdot \nabla u - \text{div}(\mathbf{c}u) + du,$$

under appropriate structural and regularity assumptions on the coefficients $\mathbf{A} \in M^{n \times n}(\mathbb{R})$, $\mathbf{b}, \mathbf{c} \in \mathbb{R}^n$, $d \in \mathbb{R}$ (essentially the maximum principle should hold), is contained in [32].

Proposition 2.3. Let $f \in L^1(\Omega; \rho)$ and $h \in L^1(\partial\Omega)$. Then there exists a unique solution $u \in L^1(\Omega)$ of problem (23) in the sense of Definition 2.1. Moreover there exists $C = C(\Omega, L) > 0$ such that

$$\|u\|_{L^1(\Omega)} \leq C(\|f\|_{L^1(\Omega; \rho)} + \|h\|_{L^1(\partial\Omega)}) \tag{24}$$

and u satisfies

$$-\int_{\Omega} u_+L^*\phi \, dx \leq \int_{\Omega} f(\text{sgn}_+u)\phi \, dx - \int_{\partial\Omega} \partial_A\phi h_+ \, d\sigma \tag{25}$$

and

$$-\int_{\Omega} |u|L^*\phi \, dx \leq \int_{\Omega} f(\text{sgn}u)\phi \, dx - \int_{\partial\Omega} \partial_A\phi |h| \, d\sigma, \tag{26}$$

for every non negative $\phi \in C_0^2(\bar{\Omega})$. We have used the notation

$$\text{sgn}r = \begin{cases} 1 & \text{if } r > 0, \\ 0 & \text{if } r = 0, \\ -1 & \text{if } r < 0, \end{cases} \quad \text{sgn}_+r = \begin{cases} 1 & \text{if } r \geq 0, \\ 0 & \text{if } r < 0. \end{cases}$$

Proof of Theorem 1.2. Uniqueness, monotonicity and estimate (12) follow from Proposition 2.3. Indeed, assume that u_1 and u_2 are solutions of (7) with data f_1, h_1 and f_2, h_2 respectively. It means that there exist $b_1(x) \in \beta(u_1(x))$ and $b_2(x) \in \beta(u_2(x))$ such that (11) holds. This implies that $w = u_1 - u_2$ is a solution of

$$\begin{cases} -Lw = f^* = f_1 - f_2 - b_1 + b_2 & \text{in } \Omega, \\ w = h^* = h_1 - h_2 & \text{on } \partial\Omega. \end{cases} \tag{27}$$

Then, estimate (12) follows from (25) when applied to w with test function ϕ_0 solution of

$$\begin{cases} -L^*\phi_0 = 1 & \text{in } \Omega, \\ \phi_0 = 0 & \text{on } \partial\Omega. \end{cases} \tag{28}$$

Also monotonicity follows from estimate (25) when applied to w with test function ϕ_0 . Uniqueness can be derived with the same procedure from (26).

Existence. We consider the Yosida approximation β_μ of β , maximal monotone graph of \mathbb{R}^2 , which we know to be a Lipschitz increasing function (see [13]). We look for a solution of the problem

$$\begin{cases} -Lu + \beta_\mu(u) = f & \text{in } \Omega, \\ u = h & \text{on } \partial\Omega. \end{cases} \tag{29}$$

The solution of such a problem is a straightforward generalization of Proposition 2.1.2 in [25].

Let us call u_μ the solution of (29), and let consider $f \in L^\infty(\Omega)$ and $h \in L^\infty(\partial\Omega)$. Then, by the monotonicity of solutions, one have the upper bound $u_\mu \leq M = \max(\sup_\Omega f, \sup_{\partial\Omega} h)$ for all $\mu > 0$. Also $\{\beta_\mu(u_\mu)\}$ is uniformly bounded in $L^\infty(\Omega)$ (see [14]).

We now show that $\{u_\mu\}$ and $\{\beta_\mu\}$ are Cauchy sequences in $L^2(\Omega)$. Given $\lambda, \mu > 0$, we subtract the equations for u_λ and u_μ , multiply the result for $u_\lambda - u_\mu$ and integrate to obtain, with the use of Lemma 2.2,

$$0 = \int_\Omega -L(u_\lambda - u_\mu)(u_\lambda - u_\mu) + \int_\Omega (\beta_\lambda(u_\lambda) - \beta_\mu(u_\mu))(u_\lambda - u_\mu) \\ \geq C \|u_\lambda - u_\mu\|_{L^2(\Omega)}^2 + (\beta_\lambda(u_\lambda) - \beta_\mu(u_\mu), (u_\lambda - u_\mu))_{L^2(\Omega)},$$

which, following [14], gives

$$C \|u_\lambda - u_\mu\|_{L^2(\Omega)}^2 + (\beta_\lambda(u_\lambda) - \beta_\mu(u_\mu), \lambda\beta_\lambda(u_\lambda) - \mu\beta_\mu(u_\mu))_{L^2(\Omega)} \leq 0.$$

Sending $\lambda, \mu \rightarrow 0$ and remembering that $\{\beta_\mu(u_\mu)\}$ is uniformly bounded in $L^\infty(\Omega)$, we get that $\|u_\lambda - u_\mu\|_{L^2(\Omega)} \rightarrow 0$. We set $u := \lim_{\mu \rightarrow 0} u_\mu$. By Lemma 2.4 of [16] also $\beta_\mu(u_\mu)$ is a Cauchy sequence in $L^2(\Omega)$ and its limit $b \in L^\infty(\Omega)$ satisfies that $b(x) \in \beta(u(x))$ since β is maximal. Passing to the limit in the definition of solution we have

$$0 = - \int_\Omega u_\mu L^* \varphi \, dx + \lambda \int_\Omega \beta_\mu(u_\mu) \varphi \, dx - \int_\Omega f \varphi \, dx + \int_{\partial\Omega} h \partial_A \varphi \, d\sigma \\ \rightarrow - \int_\Omega u L^* \varphi \, dx + \lambda \int_\Omega b \varphi \, dx - \int_\Omega f \varphi \, dx + \int_{\partial\Omega} h \partial_A \varphi \, d\sigma \tag{30}$$

for any $\varphi \in C_0^2(\bar{\Omega})$. Hence u is a solution of (7).

If $(f, h) \in L^1(\Omega; \rho) \times L^1(\partial\Omega)$ we consider $\{(f_n, h_n)\} \subset L^\infty(\Omega) \times L^\infty(\partial\Omega)$ which converges to (f, h) in $L^1(\Omega; \rho) \times L^1(\partial\Omega)$. Call u_n the solution of (7) with data f_n and h_n . Thanks to (13), u_n and b_n are Cauchy sequences in $L^1(\Omega)$ and hence converges to functions u, b respectively. Since β is maximal $b(x) \in \beta(u(x))$ and passing to the limit in the definition of solution we find that u solves (7). \square

2.2. Proof of Theorem 1.10

We start this part by giving a result on the corresponding linear problem

$$\begin{cases} u_t - Lu = f(t, x) & \text{in } Q_T, \\ u = h(t, x) & \text{on } \Sigma_T, \\ u(0, x) = u_0(x) & \text{on } \Omega. \end{cases} \tag{31}$$

Proposition 2.4. Assume $f \in L^1(Q_T; \rho) = L^1(0, T; L^1(\Omega; \rho))$, $h \in L^1(\Sigma_T)$ and $u_0 \in L^1(\Omega; \rho)$. Problem (31) possesses a unique very weak solution $u \in L^1(Q_T)$, in the sense that

$$\int_{Q_T} -(\zeta_t + L\zeta)u - f\zeta \, dxdt = - \int_{\Sigma_T} h \partial_A \zeta \, d\sigma dt + \int_\Omega \zeta(x, 0)u_0 \, dx$$

for every $\zeta \in C^{2,1}(\bar{Q}_T)$. Additionally, there hold

$$\int_{Q_T} -(\zeta_t + L\zeta)u_+ - f\zeta \operatorname{sgn}_+(u) \, dx dt \leq - \int_{\Sigma_T} h_+ \partial_A \zeta \, d\sigma dt + \int_\Omega \zeta(x, 0)(u_0)_+ \, dx \tag{32}$$

and

$$\int_{Q_T} -(\zeta_t + L\zeta)|u| - f\zeta \operatorname{sgn}(u) \, dx dt \leq - \int_{\Sigma_T} |h| \partial_A \zeta \, d\sigma dt + \int_\Omega \zeta(x, 0)|u_0| \, dx, \tag{33}$$

for every non-negative $\zeta \in C_0^{2,1}(Q_T)$.

Proof. The proof is exactly the same of [24] with Δ replaced by L . \square

Lemma 2.5. Let $f \in L^1(Q_T; \rho)$, $h \in L^1(\Sigma_T)$ and $u_0 \in L^1(\Omega; \rho)$. Then problem (9) has at most one solution. If u_1, u_2 are solutions with data f_1, h_1, u_{01} and f_2, h_2, u_{02} respectively, then

$$\|u_1 - u_2\|_{L^1(Q_T)} + \|b_1 - b_2\|_{L^1(Q_T; \rho)} \leq C (\|f_1 - f_2\|_{L^1(Q_T; \rho)} + \|h_1 - h_2\|_{L^1(\Sigma_T)} + \|u_{01} - u_{02}\|_{L^1(\Omega; \rho)}), \tag{34}$$

with $C > 0$ and $b_1 \in \beta(u_1)$, $b_2 \in \beta(u_2)$. Furthermore, if $f_1 \leq f_2$, $h_1 \leq h_2$ and $u_{01} \leq u_{02}$ then $u_1 \leq u_2$.

Proof. As for the elliptic case, we set $w = u_1 - u_2$, which is solution of

$$\begin{cases} w_t - Lw = f_1 - f_2 - b_1 + b_2 & \text{in } Q_T, \\ w = h_1 - h_2 & \text{on } \Sigma_T, \\ w(0, x) = u_{01} - u_{02} & \text{in } \Omega. \end{cases}$$

Take ψ_T solution of

$$\begin{cases} -\psi_t - L\psi = 1 & \text{in } Q_T, \\ \psi = 0 & \text{on } \Sigma_T, \\ \psi(T, x) = 0 & \text{in } \Omega. \end{cases}$$

Estimate (34) and monotonicity follow from (32) with ψ_T as test function and uniqueness follows from (33) with the same method. \square

Proof of Theorem 1.10. The proof of existence for the case of β a continuous function is again an easy adaptation of previous results in the literature concerning the special case of $L = \Delta$ (see, e.g., [23, Lemma 2.7] and [24, Lemmas 1.3 and 1.7]: notice that the assumption of β Lipschitz assumed at the beginning of the paper is not needed in both lemmas). The adaptation to the case of L given by (6) and β multivalued is completely similar to the one presented in the stationary case.

The continuity in t of the very weak solution will be used in our study of the asymptotic behaviour of solutions and can be obtained by reformulating the parabolic semilinear problem as an abstract Cauchy problem on the Banach space $X = L^1(\Omega; \rho)$,

$$(AP) \begin{cases} \frac{du}{dt}(t) + A(t)u(t) \ni f(t) & \text{in } X, \\ u(0) = u_0, \end{cases}$$

where $A(t) : D(A(t)) \rightarrow \mathcal{P}(X)$ is the operator defined by $(w, z) \in A(t) \subset X \times X$ iff $w \in L^1(\Omega)$ is the very weak solution of

$$\begin{cases} -Lw + \beta(w) \ni z & \text{in } \Omega, \\ w(x) = h(t, x) & \text{on } \partial\Omega. \end{cases}$$

Here, in the definition of the operator $A(t)$, $t \in (0, T)$ is a parameter (remember that $h \in W^{1,1}(0, T; L^1(\partial\Omega))$, hence $h(t, \cdot)$ makes sense). For a.e. $t \in (0, T)$, this operator is $T - \omega$ -accretive on X (see, e.g., [8] or [10]) for some $\omega \geq 0$ large enough. Indeed, we must show that, for some $\mu > 0$, $(I + \mu(A(t) + \omega I))^{-1}$ is a contraction on X and this is equivalent to show that if w^i , with $i = 1, 2$, are the very weak solutions of

$$\begin{cases} -\mu Lw^i + \mu\beta(w^i) + \mu\omega w^i \ni z^i(x) & \text{in } \Omega, \\ w^i(x) = h(t, x) & \text{on } \partial\Omega, \end{cases} \tag{35}$$

for some $z^i \in L^1(\Omega; \rho)$, then

$$\| [w^1 - w^2]_+ \|_{L^1(\Omega; \rho)} \leq \| [z^1 - z^2]_+ \|_{L^1(\Omega; \rho)}.$$

But this is a trivial consequence of the estimates proven in Theorem 1.2 once that ω is taken large enough (in particular $\omega > C$, the constant appearing in estimate (12) which was only dependent on Ω). In addition, for a.e. $t \in (0, T)$ this operator is m -accretive (see [8]) in the sense that $R(I + \mu A(t)) = X$. Indeed, we must prove that problem (35) (i.e., (7)) for a given right hand side in $z^i \in L^1(\Omega; \rho)$ has a unique solution, which, again, is a consequence of Theorem 1.2. Finally, since $h \in W^{1,1}(0, T; L^1(\partial\Omega))$ we get that the t -dependence of the solution has the same regularity: $w \in W^{1,1}(0, T; L^1(\Omega))$ and so the Crandall–Evans theorem [15,20] can be applied ensuring the existence and uniqueness of a mild solution $u \in C([0, T]; L^1(\Omega; \rho))$ of the abstract problem (AP). Finally, since we have uniqueness of the very weak solution of the parabolic problem, it is easy to see (as, for instance, in [11]) that both solutions must coincide and thus we get the desired time regularity result. \square

3. On the stabilization when $t \rightarrow +\infty$

Remember that the solutions of the parabolic problem (9) and the elliptic problem (7) will be indicated with u and u_∞ respectively.

Proof of Theorem 1.12. We follow some of the ideas contained in [18] (see also references therein). We define

$$U_n(s, x) = u(t_n + s, x), \quad F_n(s, x) = f(t_n + s, x), \quad H_n(s, x) = h(t_n + s, x),$$

where $t_n \rightarrow \infty$ when $n \rightarrow \infty$. By Theorem 1.10 we know that there exists $b \in L^1(0, T; L^1(\Omega; \rho))$ such that $b(t, x) \in \beta(u(t, x))$ a.e. in Q_T . Thus we also define $B_n(s, x) = b(t_n + s, x)$. Then it is clear that

$$\begin{cases} \frac{\partial U_n}{\partial s} - LU_n + \lambda B_n(s, x) = F_n(s, x) & \text{in } (-1, 1) \times \Omega, \\ U_n = H_n(s, x) & \text{on } (-1, 1) \times \partial\Omega \end{cases} \quad (36)$$

for all $n > 1$. Then from the estimate of Theorem 1.10 (which coincides with (1.26) of [24] easily adapted to the case in which L is given by (6)),

$$\|U_n\|_{L^1((-1,1) \times \Omega)} + \lambda \|B_n\|_{L^1((-1,1); L^1(\Omega; \rho))} \leq C (\|F_n\|_{L^1((-1,1); L^1(\Omega; \rho))} + \|H_n\|_{L^1((-1,1); L^1(\partial\Omega))} + \|U_n(-1, \cdot)\|_{L^1(\Omega; \rho)}).$$

Assumptions (20) and (21) imply that $F_n \rightarrow F_\infty$, and $H_n \rightarrow H_\infty$ strongly in $L^1((-1, 1); L^1(\Omega; \rho))$ and $L^1((-1, 1); L^1(\partial\Omega))$ respectively, where F_∞ and H_∞ are defined as $F_\infty(s, x) = f_\infty(x)$ and $H_\infty(s, x) = H_\infty(x)$. Moreover, from assumption (22), $\|U_n(-1, \cdot)\|_{L^1(\Omega; \rho)}$ is bounded independently of n . Consequently, due to assumption (19), $\{\rho B_n\}$ is a bounded sequence in $L^p((-1, 1); L^p(\Omega))$ with $p = 1/q$ if $q \in (0, 1)$ and for every $p > 1$ if $q = 0$. Thus,

$$\rho B_n \rightharpoonup \rho B_\infty$$

weakly in $L^p((-1, 1); L^p(\Omega))$ (after passing to a subsequence), for some B_∞ in $L^p((-1, 1); L^p(\Omega))$. Then, by [24] (Lemma 1.6(ii) easily adapted to the case in which $L \neq \Delta$), we get that $U_n \rightarrow U_\infty$ (strongly) in $C([-1 + \varepsilon, 1]; L^1(\Omega; \rho))$ for some $U_\infty \in C([-1 + \varepsilon, 1]; L^1(\Omega; \rho))$ for any $\varepsilon \in (0, 2)$. Indeed, since problem (36) is linear we can use the decomposition

$$U_n = \mathbb{P}(F_n - \lambda B_n, 0, 0) + \mathbb{P}(0, H_n, 0) + \mathbb{P}(0, 0, U_n(-1)),$$

where \mathbb{P} is the solution mapping (see [24, Lemma 1.6]). The operator $\mathbb{P}(F_n - \lambda B_n, 0, 0)$ is compact from $L^p((-1 + \varepsilon, 1) \times \Omega) \times \{0\} \times \{0\} \rightarrow C([-1 + \varepsilon, 1] \times \Omega)$, since in general $w = \mathbb{P}(q, 0, 0)$ is given by

$$w(s, x) = \int_{-1}^1 \int_{\Omega} G_L(x, y, s, \tau) q(y, \tau) dy d\tau$$

(remember that the Green function $G_L(x, \cdot, s, \cdot) \in C_0([-1 + \varepsilon, 1] \times \bar{\Omega})$). The compactness of the other terms $\mathbb{P}(0, H_n, 0)$ and $\mathbb{P}(0, 0, U_n(-1))$ was shown in Lemma 1.6(ii) of the mentioned reference. In our case, we know the continuity in time of the functions (Theorem 1.10). In particular, since $U_n \rightarrow U_\infty$ in $C([-1 + \varepsilon, 1]; L^1(\Omega))$, we find that $\{U_n(-1 + \varepsilon, \cdot)\}$ is a Cauchy sequence in $L^1(\Omega; \rho)$.

Then, by applying an estimate similar to (13) but for the parabolic problem (see estimate (1.26) of [24] easily adapted to the case in which L is given by (6)), we get

$$\begin{aligned} &\|U_n - U_m\|_{L^1((-1+\varepsilon,1) \times \Omega)} + \lambda \|B_n - B_m\|_{L^1((-1+\varepsilon,1); L^1(\Omega; \rho))} \\ &\leq C (\|F_n - F_m\|_{L^1((-1+\varepsilon,1); L^1(\Omega; \rho))} + \|H_n - H_m\|_{L^1((-1+\varepsilon,1); L^1(\partial\Omega))} + \|U_n(-1 + \varepsilon) - U_m(-1 + \varepsilon)\|_{L^1(\Omega; \rho)}), \end{aligned}$$

which proves that $\{\rho B_n\}$ is a Cauchy sequence in $L^1((-1 + \varepsilon, 1); L^1(\Omega))$ and so $\rho B_n \rightarrow \rho B_\infty$ strongly in $L^1((-1 + \varepsilon, 1); L^1(\Omega))$. Then, since β is maximal monotone, we conclude (see [9]) that $B_\infty(s, x) \in \beta(U_\infty(s, x))$ for a.e. $(s, x) \in (-1 + \varepsilon, 1) \times \Omega$.

It only remains to prove that $U_\infty(s, x) = u_\infty(x)$, with u_∞ the (unique) very weak solution of (7) with data f_∞, h_∞ . Since

$$\text{ess sup}_{t_n+s \in (0, +\infty)} \|u(t_n + s, \cdot)\|_{L^1(\Omega; \rho)} \leq C,$$

then there exists a (stationary) Radon measure $\mu_\infty \in M(\Omega; \rho)$ such that $u(t_n + s, \cdot) \rightharpoonup \mu_\infty$ weakly in $M(\Omega; \rho)$. Moreover, $u(t_n + s, \cdot) \rightarrow U_\infty(s, x)$ (strongly) in $C([-1 + \varepsilon, 1]; L^1(\Omega; \rho)) \subset C([-1 + \varepsilon, 1]; M(\Omega; \rho))$. Then, by the uniqueness of the limit, we deduce that $U_\infty(s, \cdot) = \mu_\infty(\cdot)$ for any $s \in [-1 + \varepsilon, 1]$, so that the singular part of the measure $\mu_\infty(\cdot)$ vanishes (i.e., $\mu_\infty \in L^1(\Omega; \rho)$). Let us denote now $u_\infty(x) \equiv \mu_\infty(x)$. Then $U_\infty(s, x) = u_\infty(x)$ for any $s \in [-1 + \varepsilon, 1]$. By the same reasons (thanks to the assumption on β) we get that $B_\infty(s, x) = b_\infty(x)$ for a.e. $s \in [-1 + \varepsilon, 1]$, for some $b_\infty \in L^1(\Omega; \rho)$ such that $b_\infty(x) \in \beta(u_\infty(x))$ for a.e. $x \in \Omega$. Finally, we take as test function in the definition of very weak solution of the parabolic problem $\varphi(s, x) = \psi(s)\zeta(x)$, with $\zeta \in W^{2,\infty}(\Omega) \cap W_0^{1,\infty}(\Omega)$ and $\psi \in C^1([-1, 1])$ such that $\psi|_{[-1, -1+\varepsilon]} = \psi(1) = 0$ and such that $\int_{-1+\varepsilon}^{+1} \psi(s) ds = 1$. Obviously such special $\varphi(s, x)$ is a correct test function since $\varphi \in W^{1,\infty}(-1, 1; L^\infty(\Omega)) \cap L^\infty(-1, 1; W^{2,\infty}(\Omega) \cap W_0^{1,\infty}(\Omega))$ and $\varphi(1, \cdot) = 0$. Then, from the definition of very weak solution, we get that

$$\begin{aligned} &-\int_{-1+\varepsilon}^1 \int_{\Omega} U_n(s, x) \psi'(s) \zeta(x) dx ds + \int_{-1+\varepsilon}^1 \int_{\Omega} \psi(s) U_n(s, x) L^* \zeta(x) dx ds + \lambda \int_{-1+\varepsilon}^1 \int_{\Omega} B_n(s, x) \psi(s) \zeta(x) dx ds \\ &= \int_{\Omega} U_n(-1, x) \psi(-1) \zeta(x) dx + \int_{-1+\varepsilon}^1 \int_{\Omega} F_n(s, x) \psi(s) \zeta(x) dx ds - \int_{-1+\varepsilon}^1 \int_{\partial\Omega} \psi(s) H_n(s, \sigma) \frac{\partial \zeta(\sigma)}{\partial n} d\sigma ds. \end{aligned}$$

Passing to the limit, as $n \rightarrow +\infty$, and using that $\psi(-1 + \varepsilon) = 0$, we arrive to

$$\begin{aligned} & - \left(\int_{-1+\varepsilon}^1 \psi'(s) \, ds \right) \left(\int_{\Omega} u_{\infty}(x) \zeta(x) \, dx \right) + \int_{-1+\varepsilon}^1 \psi(s) \, ds \int_{\Omega} u_{\infty}(x) L^* \zeta(x) \, dx + \lambda \int_{-1+\varepsilon}^1 \int_{\Omega} B_{\infty}(s, x) \psi(s) \zeta(x) \, dx \\ & = \int_{-1+\varepsilon}^1 \psi(s) \, ds \int_{\Omega} f_{\infty}(x) \zeta(x) \, dx - \int_{-1+\varepsilon}^1 \psi(s) \, ds \int_{\partial\Omega} h_{\infty}(\sigma) \frac{\partial \zeta(\sigma)}{\partial n} \, d\sigma. \end{aligned}$$

But

$$\int_{-1+\varepsilon}^1 \psi'(s) \, ds = 0,$$

and since $\int_{-1+\varepsilon}^{+1} \psi(s) \, ds = 1$ we get that

$$\int_{\Omega} u_{\infty}(x) L^* \zeta(x) \, dx + \lambda \int_{\Omega} b_{\infty}(x) \zeta(x) \, dx = \int_{\Omega} f_{\infty}(x) \psi(s) \zeta(x) \, dx - \int_{\partial\Omega} h_{\infty}(\sigma) \frac{\partial \zeta(\sigma)}{\partial n} \, d\sigma,$$

which shows that u_{∞} coincides with the (unique) very weak solution of the stationary problem. □

Remark 3.1. Notice that the boundedness of the trajectories assumption is considerably weaker than the usual for weak solutions (see, e.g., [18]) which is of the type $u \in L^{\infty}(0, +\infty; H^1(\Omega))$. Notice also that this condition is necessary once we assume that the conclusion of Theorem 1.12 holds.

A sufficient condition leading to the boundedness of the trajectories (assumption (22)) can be obtained by the method of super and subsolutions as in Proposition 3 of [18].

Proposition 3.2. Assume that the stationary problem (7) admits a bounded weak solution u_{∞} . Let f, f_{∞} and h, h_{∞} satisfy (20) and (21) respectively. Suppose the existence of $f_u, f_d \in L^1((0, T) \times \Omega)$ for any $T > 0$ and $h_u, h_d \in L^1((0, T) \times \partial\Omega)$ with $f_u, h_u (f_d, h_d)$ non-increasing (decreasing) in t such that

$$\begin{aligned} -\bar{f}(x) &\leq f_d(t, x) \leq f(t, x) \leq f_u(t, x) \leq \bar{f}(x), \\ -\bar{h}(x) &\leq h_d(t, x) \leq h(t, x) \leq h_u(t, x) \leq \bar{h}(x), \end{aligned}$$

for $0 \leq \bar{f} = \operatorname{div} \mathbf{c}$ with $\mathbf{c} \in L^p(\Omega)^N$, $0 \leq \bar{h} \in L^1(\partial\Omega)$ and

$$\begin{aligned} \lim_{t \rightarrow \infty} f_u(t, x) &= \lim_{t \rightarrow \infty} f_d(t, x) = f_{\infty}(x) \quad \text{in } L^1(\Omega; \rho), \\ \lim_{t \rightarrow \infty} h_u(t, x) &= \lim_{t \rightarrow \infty} h_d(t, x) = h_{\infty}(x) \quad \text{in } L^1(\partial\Omega). \end{aligned} \tag{37}$$

Let u, u_u, u_d be bounded weak solutions of (9) associated to the data (f, h, u_0) , (f_u, h_u, \bar{u}_0) and $(f_d, h_d, \underline{u}_0)$ with $\bar{u}_0, \underline{u}_0$ solutions of (7) with data \bar{f}, \bar{h} and $-\bar{f}, -\bar{h}$ respectively. If $u_u, u_d \in L^{\infty}((t_0, \infty); L^1(\Omega))$ for some $t_0 > 0$ then the conclusion of Theorem 1.12 holds.

4. On the half plane problems

Before starting with proving existence and uniqueness we shall show some results concerning the boundedness of the support of solutions. For this purpose we assume that

$$\beta(u) = u^q, \quad \lambda = 1, \quad \sup h = 1. \tag{38}$$

This hypothesis on λ and h is not a restriction as, taking it off, all calculations can be performed in the same way. Moreover the proof of Theorems 1.4 and 1.5 for the multivalued case (which formally corresponds to make $q = 0$ in all the above expressions) follows, word by word, the same proof of the case in which $q \in (0, 1)$ and replacing the identity symbol $=$ by the one of containing \supset . The details about local super and subsolutions can be seen also in the book [17, Theorem 2.16, Chapter 2].

In this first part we study some comparison functions which are essential, first of all, to give sense to the formulation of problem (15), especially to its boundary conditions, fundamental in the definition of “limit very weak solution”. Secondly, they are important also in the study of the behaviour of solutions near the origin.

Assuming (38), we repeat the same procedure of [30], looking for local supersolutions, which are solutions of

$$\begin{cases} -Lu + u^q = 0 & \text{in } B_R(x_0), \\ u = 1 & \text{on } \partial B_R(x_0), \end{cases} \tag{39}$$

where $B_R(x_0)$ is the ball with radius R and centred in x_0 . The problem is that, differently from [30], we do not know exact radial solution for (39). So we introduce a family of radial supersolution for (39).

Assume that $y(x) = \eta(|x - x_0|)$ is a radially symmetric function defined in $B_R(x_0)$. Then, if we denote with $r = |x - x_0|$, we have

$$Ly(x) = \eta'' \sum_{ij} a_{ij} \frac{x_i x_j}{r^2} + \frac{\eta'}{r} \left(\sum_i a_{ii} - \sum_{ij} a_{ij} \frac{x_i x_j}{r^2} \right). \tag{40}$$

Considering that

$$\sum_i a_{ii} - \nu \leq \sum_i a_{ii} - \sum_{ij} a_{ij} \frac{x_i x_j}{r^2} \leq \sum_i a_{ii} - \mu, \tag{41}$$

we can define the quantity

$$B_A = \sup_{x \in \Omega} \left(\sum_i a_{ii} - \sum_{ij} a_{ij} \frac{x_i x_j}{r^2} \right).$$

In particular, if we assume $\eta', \eta'' \geq 0$,

$$-Ly \geq -\nu \eta'' - \frac{B_A}{r} \eta'. \tag{42}$$

We introduce the operator

$$L_\nu(\eta) = \nu \eta'' + \frac{B_A}{r} \eta', \tag{43}$$

which operates on functions of a real scalar variable and we study the properties of the solutions to the problem

$$\begin{cases} -L_\nu \eta + \eta^q = 0 & r \in (0, R), \\ \eta(0) = 0, \quad \eta(R) = 1, \end{cases} \tag{44}$$

whenever $R \in \mathbb{R}^+$. We set the constants

$$C_0 = \left(\frac{(1-q)^2}{2\nu(q+1) + 2B_A(1-q)} \right)^{1/(1-q)}, \quad R_0 = \frac{1}{C_0^{(1-q)/2}} \tag{45}$$

and introduce the function

$$\eta_0(r) = C_0 r^{\frac{2}{1-q}}, \quad r \in [0, R_0].$$

It is a direct computation to see that η_0 is the solution of (44). For $R > R_0$ we do not know the analytic form of the solution η_R of (44) but we know that the function

$$\bar{\eta}_R(r) = \begin{cases} 0 & r \in [0, R - R_0], \\ \eta_0(r - (R - R_0)) & r \in [R - R_0, R], \end{cases} \tag{46}$$

is a supersolution. The next lemma gives the proof of this fact.

Lemma 4.1. *The function $\bar{\eta}_R$, for $R > R_0$, defined by (46) is a supersolution of (44).*

Proof. For $r \in (0, R - R_0)$, $-L_\nu \bar{\eta}_R + \bar{\eta}_R^q = 0$. For $r \in (R - R_0, R)$, calling $s = r - (R - R_0)$,

$$\begin{aligned} -L_\nu \bar{\eta}_R + \bar{\eta}_R^q &= -\nu \eta_0''(s) - \frac{B_A}{s + R - R_0} \eta_0'(s) + \eta_0^q(s) \\ &\geq -\nu \eta_0''(s) - \frac{B_A}{s} \eta_0'(s) + \eta_0^q(s) \\ &= -L_\nu \eta_0 + \eta_0^q = 0, \end{aligned}$$

where the inequality is due to the fact that $\eta_0' \geq 0$ and $R - R_0 \geq 0$. This inequality combined with the values of $\bar{\eta}_R$ at the boundary, i.e., $\bar{\eta}_R(0) = 0$ and $\bar{\eta}_R(R) = 1$, makes of $\bar{\eta}_R$ a supersolution for problem (44). □

The following lemma is the conclusion of this line of reasoning.

Lemma 4.2. *Suppose that u_R is solution of (39). Then the function $y_R(x) = \bar{\eta}_R(|x - x_0|)$ satisfies $y_R \geq u_R$ in $\bar{B}_R(x_0)$.*

Proof. The proof uses the comparison principle. Just notice, recalling (42) and Lemma 4.1, that

$$-Ly_R + y_R^q \geq -L_v \bar{\eta}_R + \bar{\eta}_R^q \geq 0 = -Lu_R + u_R^q.$$

Checking the boundary conditions we obtain the statement. \square

Using the results on local supersolutions just given, we can find a function $\rho : (R_0, \infty) \rightarrow \mathbb{R}^+$ such that if $R > R_0$ and if η_R is the solution of (44) in $(0, R)$ then

$$\eta_R = 0 \text{ in } [0, \rho(R)] \text{ and } \eta_R > 0 \text{ in } (\rho(R), R].$$

We define the function

$$d(R) = R - \rho(R), \quad R > R_0, \tag{47}$$

and the following properties hold:

Lemma 4.3. *It holds*

- (i) $R_0 < R_1 < R_2 \Rightarrow d(R_1) \geq d(R_2)$,
- (ii) $\lim_{R \rightarrow \infty} d(R) = c_{vq}$,

with

$$c_{vq} = \frac{(2\nu(1+q))^{1/2}}{1-q}. \tag{48}$$

Proof. Point (i) is exactly the same as in [30]. The second statement follows the same line too but we sketch it to show the details. We already know that on $[0, \rho(R)]$ the solution is zero since $\bar{\eta}_R$ is zero. We focus on the problem

$$\begin{cases} -L_v \eta + \eta^q = 0 & r \in (\rho(R), R), \\ \eta(\rho(R)) = 0, & \eta(R) = 1. \end{cases}$$

We set $w(r) = \eta(R - r)$ and we transform the problem into

$$\begin{cases} -\nu w'' + \frac{B_s}{R-r} w' + w^q = 0 & \text{in } (0, d(R_0)), \\ w \geq 0, \\ w(0) = 1, & w(d(R_0)) = 0. \end{cases} \tag{49}$$

We note that the second extreme of the domain in (49) should be $d(R) = R - \rho(R)$, which, according to point (i), is smaller than $d(R_0)$. This does not affect the result since $w = 0$ in $(d(R), d(R_0))$. We multiply (49) by w' and integrate over $(r, d(R_0))$

$$\begin{aligned} 0 &= -\frac{\nu}{2} \int_r^{d(R_0)} \frac{d}{ds} (w')^2 ds + \int_r^{d(R_0)} \frac{B_s}{R-r} (w')^2 ds + \frac{1}{q+1} \int_r^{d(R_0)} \frac{d}{ds} w^{q+1} ds \\ &\geq \frac{\nu}{2} (w')^2(r) - \frac{1}{q+1} w^{q+1}(r). \end{aligned}$$

The inequality in the second line is due to the non-negativity of the second term of the right-hand side in the first line. Since $0 \leq w \leq 1$ we conclude that

$$|w'(r)| \leq \left(\frac{2}{\nu(q+1)} \right)^{1/2}, \quad 0 \leq r \leq d(R_0).$$

We have found that the family of solutions $\{w(r, R) : R > R_0\}$ is equicontinuous in $[0, d(R_0)]$ and from compactness argument we can extract a subsequence $\{w(\cdot, R_n)\}$ with $R_n \rightarrow \infty$ and a function $\bar{w} \in C([0, \infty))$ such that $w(\cdot, R_n) \rightarrow \bar{w}$ uniformly on compact sets. Actually what really matters is the convergence on $[0, d(R_0)]$ as $w(r, R) = 0$ for $r > d(R_0)$ and $R > R_0$. In the limit, the equation for \bar{w} becomes

$$\begin{cases} -\nu w'' + w^q = 0, \\ w \geq 0, \\ w(0) = 1, & w(\infty) = 0, \end{cases} \tag{50}$$

whose unique solution is

$$z(t) = \left[\left(1 - \frac{t}{c_{vq}} \right)_+ \right]^{2/(1-q)}. \tag{51}$$

The convergence of $w(\cdot, R)$ to the solution of (50) implies that $d(R) \rightarrow c_{vq}$ as $R \rightarrow \infty$. \square

Lemma 4.4. Let u_R be the solution of (39). Then $u_R(x) \leq y_R(x) = \eta_R(|x - x_0|)$.

Remark 4.5. If in (39) we set the value at the boundary to be ε instead of one, all results just shown change just in the value of the constants. To be more specific, the constant C_0 appearing in (45) remains the same while R_0 should be changed into $R_\varepsilon = \varepsilon^{(1-q)/2} C_0$ and c_{vq} in $c_{vq\varepsilon} = \varepsilon^{(1-q)/2} c_{vq}$.

We remind that the Heaviside function H_v is given by

$$H_v(x_1) = \begin{cases} 0 & x_1 \in (-\infty, 0), \\ 1 & x_1 \in (0, \infty). \end{cases} \tag{52}$$

Lemma 4.6. Let u be a solution of (7) in the setting of H_{hp} with h the Heaviside function. Then

$$\mathfrak{J}(u) \subset \{(x, y) \in \Omega : x \geq 0, y < c_{vq}\} \cup \{(x, y) \in \Omega : x \leq 0, r < c_{vq}\}.$$

Proof. The proof is the same as the one proposed in [30] and uses the technique of local supersolutions. We start by giving a bound in the x_2 direction. For $R > R_0$ we consider the function y_R as in Lemma 4.4 and we set

$$\bar{u}(x_1, x_2; \xi) = y_R(x_1 - \xi, x_2 - R)$$

defined in $B_R(\xi, R)$. Since $\bar{u} = 1$ in $\partial B_R(\xi, R)$, by the comparison principle we obtain that $\bar{u} \geq u$ in $B_R(\xi, R)$. As we have chosen $\xi \in \mathbb{R}$ arbitrarily we deduce that

$$u(x_1, x_2) = 0 \quad \text{for all } x_1 \in \mathbb{R}, d(R) \leq x_2 \leq 2R - d(R).$$

Letting $R \rightarrow \infty$ and thanks to Lemma 4.3 we have that

$$u(x_1, x_2) = 0 \quad \text{for all } x_1 \in \mathbb{R}, x_2 \geq c_{vq}.$$

The boundedness for $x_1 < 0$ works similarly. Again we set

$$\bar{u}(x_1, x_2; \theta) = y_R(x_1 - R \cos \theta, x_2 - R \sin \theta) \quad \text{for } \pi/2 \leq \theta \leq \pi,$$

this time defined on $\Sigma_\theta = B_R(R \cos \theta, R \sin \theta) \cap \Omega$. The boundary of Σ_θ consists of the subset of $\partial B_R(R \cos \theta, R \sin \theta)$ which is in Ω and where $\bar{u} = 1$ and $u \leq 1$ and a part of $\partial \Omega$ with $x_1 < 0$ where $\bar{u} \geq 0$ and $u = 0$. Once again, because we can move $\theta \in [\pi/2, \pi]$, by the comparison principle we obtain that

$$u(x_1, x_2) = 0 \quad \text{in } \{(x_1, x_2) : x_1 \leq 0, d(R) < r < 2R - d(R)\},$$

with $r = (x_1^2 + x_2^2)^{1/2}$. Letting $R \rightarrow \infty$,

$$u(x_1, x_2) = 0 \quad \text{in } \{(x_1, x_2) : x_1 \leq 0, r \geq c_{vq}\}. \quad \square$$

Remark 4.7. As in Remark 4.5, if in Lemma 4.6 we substitute H_v with εH_v , the result is the same but with c_{vq} replaced by $c_{vq\varepsilon}$.

4.1. Proof of Theorem 1.5

For the proof of Theorem 1.5 we start by showing (ii) for $h = \varepsilon H_v$. We will use this result later to prove (ii) for a general boundary datum. In a second moment we will show (i) and give a numerical representation of the behaviour of $\bar{C}(q)$ and $\underline{C}(q)$ when $\varepsilon = 1$.

4.2. Heaviside function

Problem (7) under assumption H_{hp} and with $h = \varepsilon H_v$ is

$$\begin{cases} -Lu + u^q = 0 & \text{in } \Omega, \\ u(x_1, 0) = \varepsilon H_v(x_1) & x_1 \in \mathbb{R}. \end{cases} \tag{53}$$

We remind that the symmetric matrix A

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

satisfies

$$\mu |\xi|^2 \leq \xi^T A \xi \leq \nu |\xi|^2, \quad \text{for all } \xi \in \mathbb{R}^2, \tag{54}$$

for some $\mu, \nu > 0$.

Remark 4.8. The existence of $\mu > 0$ such that condition (54) holds is equivalent to the fact that

$$a_{12}^2 < a_{22}a_{11}. \tag{55}$$

Indeed, if we set $\xi = (\xi_1, \xi_2)$, we have

$$\begin{aligned} \xi^T A \xi &= a_{11}\xi_1^2 + 2a_{12}\xi_1\xi_2 + a_{22}\xi_2^2 \\ &= \mu(\xi_1^2 + \xi_2^2) + ((a_{11} - \mu)\xi_1^2 + 2a_{12}\xi_1\xi_2 + (a_{22} - \mu)\xi_2^2). \end{aligned}$$

In order for the second term to be greater or equal to zero for all $\xi_1, \xi_2 \in \mathbb{R}$, it must hold

$$a_{12}^2 < (a_{11} - \mu)(a_{22} - \mu) < a_{11}a_{22}.$$

This fact is used later on to obtain a subsolution.

In order to study the positivity set for the solution of (53) we look for a proper subsolution. What we are really interested in is the behaviour of the solution in a neighbourhood of the origin, i.e., the point $(0, 0)$. We will show that, although the boundary datum εH_v is zero for $x_1 < 0$, the solution is positive for $x_1 > -\zeta_{\mu\varepsilon}$ and x_2 sufficiently small, for some $\zeta_{\mu\varepsilon} > 0$.

The procedure to obtain a proper lower bound \underline{u} for u is the same as the one in [30]. We look for a \underline{u} solution of

$$\begin{cases} -L\underline{u} = -\varepsilon^q & \text{in } \mathbb{R} \times (0, c_{vq}), \\ \underline{u}(x_1, 0) = \varepsilon H(x_1) & x_1 \in \mathbb{R}, \\ \underline{u}(x_1, c_{vq\varepsilon}) \leq 0 & x_1 \in \mathbb{R}. \end{cases} \tag{56}$$

If such \underline{u} exists, and remembering the bounds $0 \leq u \leq \varepsilon$ where u is solution of (53), we get

$$-L(\underline{u} - u) = -\varepsilon^q + u^q \leq 0.$$

This property, by the comparison principle, assures that $u \geq \underline{u}$.

To find out an explicit formula for \underline{u} , we split $\underline{u} = u_1 + u_2$ with

$$\begin{cases} -Lu_1 = -\varepsilon^q & x_1 \in \mathbb{R}, 0 < x_2 < c_{vq\varepsilon}, \\ u_1(x_1, 0) = 0, & u_1(x_1, c_{vq\varepsilon}) = -\varepsilon & x_1 \in \mathbb{R}, \end{cases} \tag{57}$$

and

$$\begin{cases} -Lu_2 = 0 & x_1 \in \mathbb{R}, 0 < x_2 < c_{vq\varepsilon}, \\ u_2(x_1, 0) = \varepsilon H(x_1), & u_2(x_1, c_{vq\varepsilon}) \leq \varepsilon & x_1 \in \mathbb{R}. \end{cases} \tag{58}$$

We try, as in [30], to find a u_2 which depends only on the ratio $m = x_1/x_2$. Calling f such a function, we have

$$\sum_{i,j=1}^2 a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} f(x_1/x_2) = \frac{1}{x_2^2} ((a_{22}m^2 - 2a_{12}m + a_{11})f'' + 2(a_{22}m - a_{12})f').$$

Since $x_2 > 0$, we look for a solution of the differential equation

$$(a_{22}m^2 - 2a_{12}m + a_{11})f'' + 2(a_{22}m - a_{12})f' = 0,$$

which we see to be equivalent to $gf'' + g'f' = (gf')' = 0$, where $g(m) = a_{22}m^2 - 2a_{12}m + a_{11}$. We deduce that $f' = Cg^{-1}$, where C is a constant. Hence f is given by the indefinite integral

$$f = C \int \frac{1}{a_{22}m^2 - 2a_{12}m + a_{11}}.$$

From (55) we know that $a_{12}^2 < a_{11}a_{22}$, which implies that

$$f(m) = \bar{C} + C \frac{1}{(a_{11}a_{12} - a_{12}^2)^{1/2}} \arctan \left(\frac{a_{22}m - a_{12}}{a_{11}a_{12} - a_{12}^2} \right).$$

We set

$$u_2(x_1, x_2) = f_\varepsilon(x_1/x_2),$$

where f_ε is f with the constant $\bar{C} = \bar{C}_\varepsilon$ and $C = C_\varepsilon$ chosen for the specific problem. Let us check the boundary conditions: we fix $x_1 \neq 0$ and send $x_2 \rightarrow 0$. If $x_1 < 0$, then $x_1/x_2 \rightarrow -\infty$ and $f_\varepsilon(x_1/x_2) \rightarrow \bar{C}_\varepsilon - C_\varepsilon \frac{\pi}{2} (a_{11}a_{12} - a_{12}^2)^{-1/2}$. Setting $C_\varepsilon = \varepsilon\pi^{-1} (a_{11}a_{12} - a_{12}^2)^{1/2}$ and $\bar{C}_\varepsilon = \varepsilon/2$ we have that $f_\varepsilon(x_1/x_2) \rightarrow 0$. For $x_1 > 0$, we see that $f_\varepsilon(x_1/x_2) \rightarrow \varepsilon$. Since, with these values for the constants, $0 \leq f_\varepsilon \leq \varepsilon$, also the other boundary condition is satisfied.

For u_1 we choose

$$u_1(x_1, x_2) = - \left(\frac{\varepsilon}{c_{vq\varepsilon}} + \frac{\varepsilon^q c_{vq\varepsilon}}{2} \right) x_2 + \frac{\varepsilon^q}{2} x_2^2. \tag{59}$$

It is immediate to verify that u_1 given by (59) is solution of (57).

Now we check that \underline{u} is positive in a neighbourhood of the origin. We know that, for $x_2 = 0$ and $x_1 < 0$, \underline{u} is zero. We want to understand if it is positive for some $x_2 > 0$. We compute

$$\frac{\partial \underline{u}}{\partial x_2} = -\left(\frac{\varepsilon}{c_{vq\varepsilon}} + \frac{c_{vq\varepsilon}}{2}\right) + x_2 - \frac{C_\varepsilon x_1}{a_{22}x_1^2 - a_{12}x_1x_2 + a_{11}x_2^2},$$

and evaluate it in $x_2 = 0$,

$$\frac{\partial \underline{u}}{\partial x_2}(x_1, 0) = -\left(\frac{\varepsilon}{c_{vq\varepsilon}} + \frac{c_{vq\varepsilon}}{2}\right) - \frac{C_\varepsilon}{a_{22}x_1}.$$

One can see that it is strictly positive for $x_1 < 0$ sufficiently close to zero, precisely in a neighbourhood of $(-\zeta_{v\varepsilon}, 0) \times \{0\}$, where

$$\zeta_{v\varepsilon} = \frac{2C_\varepsilon c_{vq\varepsilon}}{a_{22}(2\varepsilon + c_{vq\varepsilon}^2)}.$$

4.3. General data

In this part we show that the behaviour displayed by the solution of (53) can be found also in the solutions of (1) where h is continuous, depending on the decay rate of h near zero. If h is sufficiently big when $x_1 \sim 0$, then the free boundary is not connected with $\mathcal{S}(h)$.

We select the family of functions $\{h_\delta\} \subset C(\mathbb{R})$ of the form

$$h_\delta(x_1) = \begin{cases} 0 & x_1 \leq 0, \\ Cx_1^{\frac{2}{1-q}} & 0 < x_1 \leq x_\delta, \\ 1 & x > x_\delta. \end{cases} \tag{60}$$

Proof of Theorem 1.5. For what concerns (i), consider the solution \underline{u} of (53) with $\varepsilon = 1$ and boundary data given by $H(x_1 - x_\delta)$. It is immediate to check that $H(x_1 - x_\delta) \leq h_\delta(x_1)$ for all $x_1 \in \mathbb{R}$. This implies that $\underline{u} \leq u_\delta$ where u_δ is the solution of (53) with boundary data h_δ . So if

$$x_\delta = \left(\frac{1}{C}\right)^{\frac{1-q}{2}}$$

is such that $x_\delta < \zeta_v$ then there exists V , neighbourhood of $(0, 0)$ in \mathbb{R}^2 , such that $\underline{u} > 0$ in $V \cap \{x_1 < 0, x_2 > 0\}$. But this is true whenever

$$C > \left(\frac{1}{\zeta_v}\right)^{\frac{2}{1-q}}.$$

It is enough to set $\bar{C} = \zeta_v^{-2/(1-q)}$, $\bar{\varepsilon} = 1$ and $\bar{x}_{1,\varepsilon} = (1/\bar{C})^{(1-q)/2}$. The statement follows since $\underline{u} \leq u_\delta$.

One can also try to repeat the same proof with $\varepsilon H(x_1 - x_\delta)$ and compare the results to find the best lower bound for which the expansion on the boundary of the support property holds.

For the proof of (ii) it is enough to consider the special case of $h_\varepsilon(x_1) = \underline{C}x_1^{\frac{2}{1-q}}$ for a.e. $x_1 \in (0, x_{1,\varepsilon})$ and $h_\varepsilon(x_1) = \underline{\varepsilon}$ for a.e. $x_1 \in (x_{1,\varepsilon}, +\infty)$. Indeed, if $h \leq h_\varepsilon$ then we know that the correspondent limit very weak solutions u and u_ε satisfy that $0 \leq u \leq u_\varepsilon$. So, if the non-diffusion of the boundary of the support property holds for u_ε then it also holds for u .

Consider the function $\bar{u} = w(x_1 + vx_2)$ where

$$w(s) = Cs_+^{\frac{2}{1-q}},$$

with $C > 0$. We compute

$$\begin{aligned} -L\bar{u} + \bar{u}^q &= -(a_{11} + 2a_{12}v + a_{22}v^2)w'' + w^q \\ &= x_+^{2/(1-q)} \left(-(a_{11} + 2a_{12}v + a_{22}v^2) \frac{2(1+q)}{(1-q)^2} C + C^q \right). \end{aligned} \tag{61}$$

If we take

$$C \leq \frac{(1-q)^2}{2(1+q)(a_{11} + 2a_{12}v + a_{22}v^2)}, \tag{62}$$

we have that $-L\bar{u} + \bar{u}^q \geq 0$, hence \bar{u} is a supersolution. We notice that

$$\bar{u}(\theta, r) = 0 \quad \text{for } \theta \in \Theta = (-\pi/2, \arctan(-v)),$$




Fig. 5. Graphs of $\underline{C}(q)$ and $\bar{C}(q)$.

where $\theta = \arctan(x_1/x_2)$. Again, taking \underline{C} smaller than (62) and choosing $x_{1,\varepsilon}$ and ε such that $h_\varepsilon \leq \bar{u}$, we have that $u \leq u_\varepsilon \leq \bar{u}$; so we obtain that u is zero in the sector θ . \square

Let us restrict our attention on the family of boundary data h_δ . Each function is defined completely by the value of C , since x_δ can be deduced by C and the continuity property. Of course for consistency we expect $\underline{C} \leq \bar{C}$. But can we affirm that equality holds? Actually, with our method we cannot say. In fact, let us consider for example the case of $L = \Delta$, which means $A = Id$. We computed the values

$$\bar{C} = \left(\frac{\pi(q^2 - q + 2)}{(1-q)\sqrt{2(q+1)}} \right)^{\frac{2}{1-q}} \quad \text{and} \quad \underline{C} = \frac{(1-q)^2}{2(1+q)}.$$

We see from the graphs in Fig. 5 that $\underline{C} < \bar{C}$ for $q \in (0, 0.2)$, and the difference is quite big. For $q \in (0.2, 1)$ the difference becomes even bigger as \underline{C} is decreasing while \bar{C} is increasing. The question of what happens when $\underline{C} < C < \bar{C}$ is still open and a different approach or a finer analysis is needed.

Proof of Corollary 1.6. The additional properties are a by-product of the proof of Theorem 1.5. Indeed, for (1), since the supersolution w already satisfies this property, so does the solution because it is non-negative. Point (2) is true because we have shown that $u \geq u_1 + u_2$ with u_1 and u_2 solutions of the problems (57) and (58) respectively and $\underline{u} = u_1 + u_2$ satisfies the required properties. \square

4.4. Proof of Corollary 1.13

In the case of (i) we can use the same supersolution (which we denote now by $\bar{u}(x)$) as for the stationary case. Since our initial condition is $u_0 = 0$, then, applying the comparison result, we get that $0 \leq u(t, x) \leq \bar{u}(x)$ for any $t > 0$ and a.e. $(x_1, x_2) \in \mathbb{R} \times [0, \infty)$.

The proof of part (ii) comes from the fact that $u_1(x_1, x_2) + u_2(x_1, x_2)$ is a subsolution for the parabolic problem for $t \in (0, T]$.

References

- [1] R.A. Adams, J.J.F. Fournier, Sobolev Spaces, second ed., in: Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003.
- [2] L. Alvarez, J.I. Díaz, On the retention of the interfaces in some elliptic and parabolic nonlinear problems, *Discrete Contin. Dyn. Syst.* 25 (1) (2009) 1–17.
- [3] S.N. Antontsev, J.I. Díaz, S. Shmarev, Energy Methods for Free Boundary Problems: Applications to Nonlinear PDEs and Fluid Mechanics, in: Progress in Nonlinear Differential Equations and their Applications, vol. 48, Birkhäuser Boston Inc., Boston, MA, 2002.
- [4] D.E. Apushkinskaya, N. Matevosyan, N.N. Uraltseva, The behavior of the free boundary close to a fixed boundary in a parabolic problem, *Indiana Univ. Math. J.* 58 (2) (2009) 583–604.
- [5] D.E. Apushkinskaya, N.N. Uraltseva, Uniform estimates near the initial state for solutions of the two-phase parabolic problem, *Algebra i Analiz* 25 (2) (2013) 63–74.
- [6] D.E. Apushkinskaya, N.N. Uraltseva, Kh. Shakhgolyan, On global solutions of a parabolic problem with an obstacle, *Algebra i Analiz* 14 (1) (2002) 3–25.
- [7] R. Aris, The Mathematical Theory of Diffusion and Reaction in Permeable Catalysis, Clarendon Press, Oxford, 1975.
- [8] P. Benilan, M.G. Crandall, A. Pazy, Nonlinear Evolution Governed by Accretive Operators, book in preparation.
- [9] Ph. Bénilan, M.G. Crandall, P. Sacks, Some L^1 existence and dependence results for semilinear elliptic equations under nonlinear boundary conditions, *Appl. Math. Optim.* 17 (3) (1988) 203–224.
- [10] Ph. Bénilan, P. Wittbold, Nonlinear evolution equations in Banach spaces: basic results and open problems, in: Functional Analysis (Essen, 1991), in: Lecture Notes in Pure and Appl. Math., vol. 150, Dekker, New York, 1994, pp. 1–32.
- [11] Ph. Benilan, P. Wittbold, On mild and weak solutions of elliptic–parabolic problems, *Adv. Differential Equations* 1 (6) (1996) 1053–1073.
- [12] H. Brézis, Une équation semilinéaire avec conditions aux limites dans \mathbb{R}^n , 1972.
- [13] H. Brézis, Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert, in: North-Holland Mathematics Studies, vol. 5, North-Holland Publishing Co., Amsterdam, 1973, Notas de Matemática (50).

- [14] H. Brézis, W.A. Strauss, Semi-linear second-order elliptic equations in L^1 , *J. Math. Soc. Japan* 25 (1973) 565–590.
- [15] M.G. Crandall, L.C. Evans, On the relation of the operator $\partial/\partial s + \partial/\partial \tau$ to evolution governed by accretive operators, *Israel J. Math.* 21 (4) (1975) 261–278.
- [16] M.G. Crandall, A. Pazy, Semi-groups of nonlinear contractions and dissipative sets, *J. Funct. Anal.* 3 (1969) 376–418.
- [17] J.I. Díaz, *Nonlinear Partial Differential Equations and Free Boundaries. Vol. I: Elliptic Equations*, in: *Research Notes in Mathematics*, vol. 106, Pitman (Advanced Publishing Program), London, 1985.
- [18] J.I. Díaz, F. de Thélin, On a nonlinear parabolic problem arising in some models related to turbulent flows, *SIAM J. Math. Anal.* 25 (4) (1994) 1085–1111.
- [19] J.I. Díaz, J.-M. Rakotoson, On the differentiability of very weak solutions with right-hand side data integrable with respect to the distance to the boundary, *J. Funct. Anal.* 257 (3) (2009) 807–831.
- [20] L.C. Evans, Nonlinear evolution equations in an arbitrary Banach space, *Israel J. Math.* 26 (1) (1977) 1–42.
- [21] D. Kröner, J.-F. Rodrigues, Global behaviour for bounded solutions of a porous media equation of elliptic–parabolic type, *J. Math. Pures Appl.* (9) 64 (2) (1985) 105–120.
- [22] J.-L. Lions, E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications. Vol. I*, in: *Die Grundlehren der mathematischen Wissenschaften, Band 181*, Springer-Verlag, New York, 1972, Translated from the French by P. Kenneth.
- [23] M. Marcus, L. Véron, Initial trace of positive solutions of some nonlinear parabolic equations, *Comm. Partial Differential Equations* 24 (7–8) (1999) 1445–1499.
- [24] M. Marcus, L. Véron, Semilinear parabolic equations with measure boundary data and isolated singularities, *J. Anal. Math.* 85 (2001) 245–290.
- [25] M. Marcus, L. Véron, *Nonlinear Second Order Elliptic Equations Involving Measures*, in: *De Gruyter Series in Nonlinear Analysis and Applications*, Berlin, 2013.
- [26] Y. Martel, P. Souplet, Small time boundary behavior of solutions of parabolic equations with noncompatible data, *J. Math. Pures Appl.* (9) 79 (6) (2000) 603–632.
- [27] V.P. Mikhailov, *Partial Differential Equations*, Mir., Moscú, 1978.
- [28] J.-F. Rodrigues, *Obstacle Problems in Mathematical Physics*, in: *North-Holland Mathematics Studies*, vol. 134, North-Holland Publishing Co., Amsterdam, 1987, *Notas de Matemática [Mathematical Notes]*, 114.
- [29] H. Shahgholian, N. Uraltseva, Regularity properties of a free boundary near contact points with the fixed boundary, *Duke Math. J.* 116 (1) (2003) 1–34.
- [30] C.J. van Duijn, L.A. Peletier, How the interface approaches the boundary in the dead core problem, *J. Reine Angew. Math.* 432 (1992) 1–21.
- [31] J.L. Vázquez, *The Porous Medium Equation: Mathematical Theory*, in: *Oxford Mathematical Monographs*, The Clarendon Press Oxford University Press, Oxford, 2007.
- [32] L. Véron, Elliptic equations involving measures, in: *Stationary Partial Differential Equations. Vol. I. Handb. Differ. Equ.*, North-Holland, Amsterdam, 2004, pp. 593–712.