# Electron beams: partially flat solutions of a nonlinear elliptic equation with a singular absorption term

Jesús Ildefonso Díaz Instituto de Matemática Interdisciplinar, Universidad Complutense de Madrid, Plaza de las Ciencias, nº 3, Madrid, 28040, Spain jidiaz@ucm.es

November 11, 2025

Dedicated to my Master, Haïm Brezis, who trusted me and gave me wings to fly higher.

Abstract. In the so-called Child-Langmuir law, established since 1911, an electron beam is formed linking two electrodes, which are assumed to be two parallel plates of area A, separated to a finite distance D. When  $D \ll \sqrt{A}$ , "edge effects" are negligible and the modelling is reduced to a nonlinear boundary problem for a singular ordinary differential equation in which a constant coefficient (the generated electric current j) must be found in order to get simultaneously Dirichlet and Neumann homogeneous boundary conditions in one of the extremes. If  $D > \sqrt{A}$ , then the problem becomes much more difficult since the "edge effects" arise in the plane (x,y) and the electric current (now j(x)) due to the presence of a very large perpendicular magnetic field) must be determined in order to get solutions u(x,y) of a singular semilinear equation which are partially flat ( $u = \frac{\partial u}{\partial n} = 0$  on a part of the boundary). In this paper, we offer a rigorous mathematical treatment of some former studies (Joel Lebowitz and Alexander Rokhenko (2003) and Alexander Rokhenko (2006)), where several open questions were left open: for instance, the need for a singularity of j(x) near the cathode edge to get such partially flat solutions.

Key words: Electron beams, space charge, Child-Langmuir law, singular semilinear equation, partially flat solution, super and subsolution method,  $H^1$ -matching and its generalizations. Subject Classification: 35Q99, 78A20, 35J75, 35J60.

### 1 Introduction

In 1911, the American physicist Clement D. Child (1868 – 1933) proposed in [11] an experimental law ("Child's law") that he deduced after modelling the electric current that flows between the plates of a vacuum tube: the main components in electronics from about 1905 to 1960, when transistors and integrated circuits mostly supplanted them. For some time (1907-1908), Child was visiting J.J. Thompson (1856-1940), 1906 Nobel Prize in Physics, who built the cathode ray tubes, proving in 1897 the existence and charge of the electron. Child's law is still a staple of textbooks treating charged particle motion in vacuum and in solids. After normalizing the variable and the unknown, and by considering the case in which the anode and cathode are simulated by planes at a finite distance, Child proposed a singular nonlinear ordinary differential equation which perhaps is one of the older singular equations considered in the literature. Its formulation was the following: find j > 0 (modelling the maximum generated electric

current) to get a positive solution of the nonlinear boundary problem

$$\begin{cases} -u''(y) + \frac{j}{\sqrt{u(y)}} = 0 & y \in (0, 1), \\ u(0) = 0 & u(1) = 1, \end{cases}$$
 (1)

such that

$$u'(0) = 0$$
 and  $u > 0$  on  $(0, 1)$ .

Notice that since u(0) = 0 the singularity of the equation is present on one of the boundaries (y = 0), representing the cathode. Condition u'(0) = 0 says that the electron flux is assumed to start from rest, since the electric field is zero there. This law was later considered in a series of papers (see, e.g., [38]), and improved by Irving Langmuir (1881-1957), 1932 Chemistry Nobel Prize. Since then, the above law is called the "Child-Langmuir law". In physical terms, it is known as the "three-halves-power law" (see Remark 2), since it says that the electric current J is proportional to the 3/2-power of the voltage,  $V^{3/2}$  (in contrast to Ohm's law on electric circuits for which J = kV for some k > 0). We will see, in Section 3, that in the renormalized formulation

$$j \equiv \frac{4}{9}$$
 and  $u(y) = y^{4/3}$  for  $x \in (0, 1)$ ,

in contrast with the 1823 Ohm's law [u(y) = ky, for some k > 0].

In the Child-Langmuir modelling, the electrodes are assumed to be two parallel plates of area A, separated by a finite distance D (D=1 after some change of variables). They assumed that  $D \ll \sqrt{A}$ , so that "edge effects" are negligible.

The study of the problem in which edge effects are taken into account is much more difficult and was a subject of research until our days (see, e.g., the survey [51]). A more realistic modelling, already considered by many authors (see, e.g., [45], [46]), takes place in the plane (X, y), in which the cathode is only an interval  $(-a, a) \times \{0\}$  of the device (the set of points  $(X, y) \in (-a - b, a + b) \times (0, 1)$ , for a given b > 0), and it is assumed the presence of a strong magnetic field. This makes the electron beam well confined in such a way that the electric current is only X-dependent, J(X, y) = j(X). The primitive question regarded by the Child-Langmuir law can now be stated in the following terms as an overdetermined problem: given a, b > 0, find sufficient conditions on a x-dependent function  $j : (-a - b, a + b) \rightarrow [0, +\infty)$ , with

$$\begin{cases}
j(X) > 0 & \text{if } X \in (-a, a), \ j \in L^1_{loc}(-a, a), \\
j(X) = 0 & \text{if } X \in (-a - b, -a) \cup (a, a + b),
\end{cases}$$
(2)

to get the solvability of the singular nonlinear boundary value problem

$$\widehat{P}_{a,b,j} = \begin{cases}
-\Delta u + \frac{j(X)}{\sqrt{u}} = 0 & X \in (-a - b, a + b), \ y \in (0, 1), \\
u(X, 0) = 0 & X \in (-a - b, a + b), \\
u(X, 1) = 1 & X \in (-a - b, a + b), \\
u(\pm (a + b), y) = y & y \in (0, 1),
\end{cases}$$
(3)

with the additional conditions

$$\widehat{AC}_{a,b} = \begin{cases} \frac{\partial u}{\partial y}(X,0) = 0 & X \in (-a,a), \\ u(X,y) > 0 & X \in (-a-b,a+b), \ y \in (0,1). \end{cases}$$

$$\tag{4}$$

The more important fact considered in the literature is the study of the adaptation of profiles  $u(\cdot, y)$  from the external profile (which is assumed to be given by u(b, y) = y, as in Ohm's law) to the profile in the center of the cathode u(0, y), where the authors expect to have the profile correspondent to the Child-Langmuir law,  $u(0, y) = y^{4/3}$  for  $y \in (0, 1)$ .

A very delicate question is the behaviour of j(X) near  $X = \pm a$  since it must generate complicated edge effects (for instance, by the *strong maximum principle*, we know that  $\frac{\partial u}{\partial y}(X,0) > 0$  if  $(-a-b,-a) \cup (a,a+b)$ , since u is harmonic on  $((-a-b,-a) \cup (a,a+b)) \times (0,1)$ ).

To simplify the formulation, due to the symmetry of the problem, it is usual to consider only (see, e.g., [45], [46], [44]) the transmission of the profiles  $u(X_0, y)$ ,  $y \in (0, 1)$ , when  $X_0 \in [0, a + b]$  in one of the two symmetrical parts of the device (for example, the one on the right). To give the edge behaviour (on the boundary of the cathode) a central presence in the formulation, it is traditional to introduce the change of variable x = X - a,  $X \in (0, a + b)$  and then  $x \in (-a, b)$  for a given b > 0. In this way, (-a, 0) becomes the center of the cathode and (0, 0) the edge of the right side of the cathode. We arrive then to the formulation which will be considered in this paper: given a, b > 0, we consider the problem of finding the coefficient j(x) such that there exists a solution of the singular boundary value problem on  $\Omega = (-a, b) \times (0, 1)$ 

$$P_{a,b,j} = \begin{cases} -\Delta u + \frac{j(x)}{\sqrt{u}} = 0 & \text{in } \Omega, \\ u(x,0) = 0 & x \in (-a,b), \\ u(x,1) = 1 & x \in (-a,b), \\ u(-a,y) = y^{4/3} & y \in (0,1), \\ u(b,y) = y & y \in (0,1), \end{cases}$$

$$(5)$$

satisfying the additional conditions

$$AC_{a,b} = \begin{cases} \frac{\partial u}{\partial y}(x,0) = 0 & x \in (-a,0), \\ u > 0 & \text{in } \Omega. \end{cases}$$
 (6)

Notice that the additional conditions imply a failure of the "unique continuation property" and the Hopf-Oleinik lemma related to the strong maximum principle (see, e.g., the exposition made in [25]): indeed, u>0 in  $(-a,b)\times(0,1)$  although  $u=\frac{\partial u}{\partial y}=0$  on  $(-a,0)\times\{0\}$ . The additional condition,  $\frac{\partial u}{\partial y}(x,0)=0$  for  $x\in(-a,0)$ , represents the vanishing of the electric field on the half of the cathode under consideration. This type of behaviour arises in many free boundary problems ([18]) for semilinear equations with a non-Lipschitz absorption. It is quite similar to the requirement of the so-called flat solutions arising, for instance, in the study of linear ([20], [25]) and nonlinear ([26]) Schrödinger equations. Here, the constraint on the function j and the singularity of the nonlinear absorption term  $(1/\sqrt{u})$  generate some important additional difficulties.

Despite the relevance of the space-charge-limited flows in many applications (vacuum and solid electronic devices, electron guns, particle accelerators, high current diodes, tubes, thyristors, etc., see, e.g., [37], [35], [34]) the study of 2-d simple geometries as here considered presents important difficulties, and it was the benchmark for almost a century, in particular for the study of the generated current j(x) in the vicinity of the cathode edges (see Figure 1). Many previous studies, based on numerical and asymptotic methods (see, e.g., [50], [45], [46], [44] and their many references) claim that due to the important constraint on j(x) (independence with respect to y) it is needed to assume that

$$\lim_{x \nearrow 0} j(x) = +\infty,$$

(see, in Figure 1, the singularities of the function j(x) at the extremes  $\pm a$ , for different widths a of the cathode). A rigorous mathematical proof of it was left as an open problem in the cited references.

One of the main goals of this paper is to prove, and make it precise, a conjecture proposed in many numerical analysis experiences (see, e.g., [44] and its references) concerning the formation of "wings" in the electric current (i.e., the formation of a singularity of j(x) near the cathode edge). In addition, we will prove a stronger version of the positivity of the solution: in fact, the solution is non-degenerate (see subsection 4.4).




Figure 1: Wings of j(x) in some numerical experiments for different values of the cathode width a: adapted from [48].

**Theorem 1** There exists  $A_0, b_0 > 0$  and  $\beta_0 \in (0, \frac{1}{2})$ , such that, if  $b \ge b_0 > 0$ , and we assume

$$\begin{cases} j(x) = \frac{A}{(-x)^{\beta}}, & \text{for } x \in (-a, 0), \\ j(x) = 0 & \text{for } x \in (0, b), \end{cases}$$
 (7)

with

$$0 \le \beta < \beta_0 \text{ and } A \in (0, A_0). \tag{8}$$

Then there exists a weak solution  $u \in L^2(\Omega; \delta)$  with  $\delta = d((x, y), \partial \Omega)$ , of problem  $P_{a,b,j}$  and the additional conditions  $AC_{a,b}$ . Moreover

$$0<\underline{C}\delta(x,y)^{4/3}\leq u(x,y)\leq \overline{C}\delta(x,y)^{\overline{\alpha}}\ a.e.\ (x,y)\in (-a,0)\times (0,1),$$

and  $Cy^{\alpha} \leq u(0,y), \ y \in (0,1), \ with \ \alpha = \frac{2}{3}(2-\beta), \ 1 < \overline{\alpha} \leq \frac{4}{3}, \ C,\underline{C},\overline{C} > 0.$  Finally, u is unique in the class of non-degenerate solutions.

The proof will be made by constructing suitable super and subsolutions for several auxiliary problems that are obtained by matching some strategic functions on different subdomains. In particular, we will need to consider the nonlinear singular eigenvalue type problem

$$\begin{cases}
-U''(s) + \frac{C}{\sqrt{U(s)}} = \lambda U(s) & s \in (-R, R), \\
U(\pm R) = 0,
\end{cases} \tag{9}$$

for which the global bifurcation diagram (in terms of the parameter  $\lambda$ ) will be completely characterized (see Figure 2). Here the positive constants C and R are given. We will show: i) there is a bifurcation from the infinity for  $\lambda$  near  $\lambda_1(R)$  (the first eigenvalue of the linear problem with C=0), ii) the bifurcation curve is strictly decreasing (which implies the uniqueness of the nonnegative solution U) and iii) the curve is not  $C^1$  for a suitable value  $\lambda = \lambda^* > \lambda_1(R)$  corresponding to a "flat solution" (i.e. the solution U is such that  $U'(\pm R) = 0$  and U(s) > 0 for any  $s \in (-R, R)$ ). This extends several results in the previous literature (see, e.g., [27], [24], and their references).




Figure 2: Bifurcation from infinity and critical values of  $\lambda$  for a flat solution.

Theorem 1 confirms many numerical experiences that lacked a rigorous proof. The progressive changes in the profiles can be illustrated by Figure 3.

We point out that the nature of the problem changes radically when the singular nonlinearity arises on the other side of the equality (as a forcing term): see, e.g. [13] and [28]. Most of the motivations to consider singular absorption terms have their foundation in fields such as chemical reactions, non-Newtonian flows and other applied subjects very different from the one of space-charge-limited flows (see, e.g., [31], [29], [12] and [18]). A survey on singular elliptic equations can be found in [33].




Figure 3: Progressive changes of profiles  $u(\cdot, y)$ .

We will need to construct several super and subsolutions satisfying suitable qualitative properties. This will require matching different functions. We point out that two remarkable matching studies in the literature were those of G. Gamov in [32] (when proving the tunneling effect) and the one by H. Brezis in [5] (proving the compact support of solutions of suitable variational inequalities: see the exposition made in [19]). Here we will need to use a matching method that is not the usual  $H^1$ -matching criterion but a

looser one that allows the formation of singularities as long as they have a certain sign (see Remark 25 below).

The organization of this paper is the following: Section 2 is devoted to the modelling of the problem. In particular, we will make mention to some optimization criterion on j(x) (see Remark 6). The one-dimensional case, including the Child-Langmuir law, is revisited in Section 3. The elegant study of the one-dimensional case is due to H. Brezis (personal communication in 2004). We also consider the non-autonomous case j = j(x) since it will be applied later. The main goal of this work is the study of the two-dimensional formulation that will be developed in the last Section 4 divided into different subsections. We will give details on the construction of a positive subsolution and a supersolution, both satisfying the additional condition, and prove the uniqueness of solutions in the class of non-degenerate solutions.

### 2 On the modelling of the 2-d space-charge-limited flow

Following [45] (see also [43]), a general formulation of the problem under consideration could start with the consideration of the stationary Maxwell system of equations for the electric and magnetic fields  $(\mathbf{E}, \mathbf{B})$  defined on a set  $\widetilde{\Omega} \subset \mathbb{R}^3$ ,  $\widetilde{\Omega} = \mathbb{R} \times (0, D) \times \mathbb{R}$ , with D > 0 given, separating two conducting electrodes placed on the planes Y = 0 (cathode) and Y = D (anode) [with  $\partial \widetilde{\Omega} = \Gamma_0 \cup \Gamma_1$ ]:

$$\begin{cases}
\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0} \\
\nabla \cdot \mathbf{B} = 0
\end{cases}$$

$$\nabla \times \mathbf{E} = \mathbf{0}$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{J}.$$
(10)

Here  $\rho(X, Y, Z)$  is the electron-charge density,  $\epsilon_0$  the free space permittivity and  $\mathbf{J}(X, Y, Z)$  denotes the current density (here we are using a variable notation (X, Y, Z) different from the one of the Introduction). Since we are assuming that  $\rho$  is stationary we get that

$$\operatorname{div} \mathbf{J} = 0 \text{ in } \widetilde{\Omega}.$$

Now we assume that the *cathode* is in the (X,Z) plane, Y=0, and it has a width 2A. We also assume that there is a *very strong magnetic field*  $\mathbf{B}$ , which is perpendicular to the electrodes ( $\mathbf{B}$  is in the Y-direction), inhibiting the transversal components of the electron velocities v(X,Y), and then  $\rho(X,Y,Z)=\rho(X,Y)\chi_{\{|X|\leq A\}}(X,Y)$ , where  $\chi_{\{|X|\leq A\}}(X,Y)$  is the characteristic function

$$\chi_{\{|X| \leq A\}}(X,Y) = \left\{ \begin{array}{ll} 1 & \text{if } |X| \leq A \\ 0 & \text{otherwise.} \end{array} \right.$$

The fact that  $\rho$  is not constant is the reason to call this type of problems as *space charge*. Due to the assumption on **B**, we know that the potential U of the electric field  $(\mathbf{E} = -\nabla U)$  is Z-independent, i.e., U = U(X,Y).

We assume that the emitted electrons leave the cathode with zero velocity and thus, if we take U = U(X,Y) = 0 in the cathode, the total mechanical energy is  $E_0 = 0$ . If e and m represent the charge and the mass of the electron, the conservation of the mechanical energy leads to the equation

$$\frac{m}{2}v^2(X,Y) = eU(X,Y). \tag{11}$$

i.e.,

$$v(X,Y) = \sqrt{\frac{2eU(X,Y)}{m}}.$$

Remember that the mechanical force (by a negative charge) is given by  $\mathbf{F} = (-e)\mathbf{E} = -e(-\nabla U) = e\nabla U$  and thus the potential energy is -eU. From this, we deduce that the current density is only dependent on X,  $\mathbf{J}(X,Y,Z) = J(X)\chi_{\{|X| \le A\}}(X,Y)\mathbf{e}_2$  and determines the velocity of electrons. We also recall that

$$\mathbf{J} = -\rho v \mathbf{e}_2 := J(X) \chi_{\{|X| \le A\}}(X, Y) \mathbf{e}_2.$$

Then

$$\rho(X,Y) = -\frac{J(X)\chi_{\{|X| \leq A\}}(X,Y)}{\sqrt{\frac{2eU(X,Y)}{m}}}.$$

We introduce now the *dimensionless variables* (once again, the notation is different from the one used in the Introduction)

$$x = \frac{X}{D}, y = \frac{Y}{D}$$
 and  $a = \frac{A}{D}$ ,

and the dimensionless functions

$$u(x,y) = \frac{U(X,Y)}{V} \text{ and } j(x) = \frac{9}{4} \sqrt{\frac{m}{2e}} \frac{J(X)}{\epsilon_0 V^{3/2}}.$$

Then, the first equation of the Maxwell system and the conservation of the mechanical energy lead to the singular nonlinear Poisson equation

$$\Delta u(x,y) = -4\pi \rho(x,y) = \frac{j(x)}{\sqrt{u(x,y)}},$$

with  $\rho(x, y) = 0$  (i.e. j(x) = 0) if |x| > a.

**Remark 2** In the dimensionless process, it is possible to choose a different new expression for J(X). So, for instance, in [45] it was used the function  $\tilde{j}(x) = \frac{4}{9}j(x)$ , so that the one-dimensional case corresponds to  $\tilde{j}(x) \equiv 1$ . In any case, the "three-halves-power law" Child-Langmuir law says that in the one-dimensional case

$$J = \frac{4}{9} \epsilon_0 \sqrt{\frac{2e}{m}} \frac{V^{3/2}}{D^2}.$$
 (12)

Remark 3 Most of the results of the following sections of this paper can be applied in the framework of two concentric cylindrical diodes (see the formulation considered in [10]). A different modelling was considered in [9] where they study a section through a thin wire at high potential contained within an earthed conductor at zero potential.

Remark 4 The study of the mathematical treatment of this problem was suggested to me by Haïm Brezis, during my visit to his Department of the University of Paris VI, in March 2003. Brezis attended a seminar by Joel Lebowitz (on his article [45]), at his workplace, Rutgers University (where Brezis would later have a contract as Distinguished Visiting Professor from 2004 until the date of his death in 2024). I received a more concrete formulation in the fax of Brezis on May, 21, 2004 (containing the Theorem 10 below) and I sent him a first set of my results on the two-dimensional problem in June and September of 2004. We made several working sessions during my stay in Paris, as Invited Professor of Paris VI, from February 15, to March 14 of 2005, but without finding a positive subsolution (see subsection 4.2 below). In this occasion, we received a visit from Joel Lebowitz to Paris VI University, and we exchanged opinions in a joint session the three of us together (see Remark 5 below and the picture at the end of this paper). I produced a new draft, sent to both of them, on June 27, 2006. I made a public presentation of the results in the Opening Workshop of the European FIRST Project on June 30, 2009, in Orsay (University of Paris Sud). Later, I presented a first version of the results of this article at the conference, "Recent and new perspectives in Nonlinear Analysis", held in Urbino (Italy), November 3-4, 2022 and at the IMEIO PhD Course, of the Universidad Politécnica de Madrid, on January 16, 2023.

Remark 5 (Suggestion by Joel Lebowitz, Paris, 2005). For other different geometrical electrode shapes, we can assume that the open (possibly unbounded) set  $\widetilde{\Omega} \subset \mathbb{R}^3$  has two components defining its finite boundary  $\partial \widetilde{\Omega} = \Gamma_0 \cup \Gamma_1$  and the problem is to find a pair  $(\mathbf{J}(x,y,z), u(x,y,z))$  satisfying (in some weak sense) the following set of conditions:

$$\begin{aligned} div\mathbf{J} &= 0 \text{ in } \widetilde{\Omega}, \\ u(x,y,z) &\geq 0 \text{ in } \widetilde{\Omega}, \\ \sqrt{u}\Delta u &= j(x,y,z) \text{ in } \widetilde{\Omega}, \\ j(x,y,z) &= |\mathbf{J}(x,y,z)|, \\ \frac{\mathbf{J}(x,y,z)}{|\mathbf{J}(x,y,z)|} &= \frac{\nabla u(x,y,z)}{|\nabla u(x,y,z)|} \\ u \mid_{\Gamma_0} &= 0 \text{ and } u \mid_{\Gamma_1} = 1. \end{aligned}$$

Remark 6 As a matter of fact, already in the pioneering works by Child and Langmuir it was mentioned that the interesting case corresponds to a current density  $J^*$  defined as the "largest current" that can be emitted without time-dependent behavior and thus independent of any thermodynamic effect on the cathode. This will be illustrated later for the one-dimensional case (when justifying why j = 4/9). For a general formulation, as the one presented in Remark 5, this kind of optimization criterion can be stated as follows (suggestion by Joel Lebowitz, Paris, 2005): from the first of the above conditions we deduce that there exists a constant  $\alpha = \alpha(\mathbf{J})$  such that for any simple curve  $\Gamma$  (when  $\widetilde{\Omega}$  is unbounded  $\Gamma$  can be unbounded without multiple intersection points) we have that

$$\int_{\Gamma} \mathbf{J} \cdot \mathbf{n} d\sigma = \alpha.$$

We define the set C of "admissible solutions"  $(\mathbf{J}(x,y,z),u(x,y,z))$  by means of the set of conditions given in Remark 5 and then the problem is now to find  $(\mathbf{J}^*,u^*)\in\mathcal{C}$  such that

$$\alpha(\mathbf{J}^*) = \max_{(\mathbf{J}, u) \in \mathcal{C}} \alpha(\mathbf{J}).$$

**Remark 7** As already mentioned in the Introduction, when  $a = +\infty$  the problem can be associated with a one-dimensional formulation for which we get  $\mathbf{J}^*(x,y,z) = \frac{4}{9}\mathbf{e}_2$  and, as we will prove in the next Section,  $u^*(y) = y^{4/3}$ . An interesting open question can be stated in this framework: is it true that if  $(\mathbf{J}, u) \in \mathcal{C}$  then necessarily  $\mathbf{J}(x, y, z) = j\mathbf{e}_2$  with j constant and u = u(y)?

**Remark 8** In the case of space-charge-limited flows  $(0 < a + b < +\infty)$  the boundary condition  $u(\pm(a + b), y) = y$  for  $y \in (0, 1)$  was first proposed in [45], since the external electric field **E** must behave, at least at very long distances, in a similar way to the case without any cathode (a = 0).

# 3 The one-dimensional Child-Langmuir law revisited

### 3.1 The autonomous case

The problem associated with the one-dimensional Child-Langmuir law can be formulated in the following terms: find j > 0 in order to get a function  $u \in W^{2,1}(0,1)$  solving the boundary problem

$$\begin{cases} -u''(y) + \frac{j}{\sqrt{u(y)}} = 0 & y \in (0,1), \\ u(0) = 0 & u(1) = 1, \end{cases}$$
 (13)

and such that

$$u'(0) = 0 \text{ and } u > 0.$$
 (14)

We recall that, in this physical case, the only interest is in positive solutions u(y) > 0 for any  $y \in (0, 1)$  but the problem also arises in other fields (for instance, in Chemical Engineering) where the possibility of having a dead core (a subinterval  $(0, \xi) \subset (0, 1)$  on which u = 0) makes perfect sense for suitable values of j.

Here  $W^{2,1}(0,1)$  denotes the usual Sobolev space requiring that  $u'' \in L^1(0,1)$  (see, e.g., [6]). Since  $W^{2,1}(0,1) \subset C^1([0,1])$  condition (14) is well justified

We point out that since the singular nonlinear term  $\frac{1}{\sqrt{u}}$  is neither monotonically non-decreasing, nor Lipschitz continuous, even in the case in which j > 0 is given, we cannot apply general results in the literature on the uniqueness of solutions.

As a matter of fact, given j > 0, some different notions of solution of (13) can be introduced (being compatible with the supplementary condition (14)). A natural notion of solution arises when problem (13) is understood in the framework of the Calculus of Variations. We define

$$K = \{u \in H^1(0,1) \text{ such that } u(0) = 0, u(1) = 1 \text{ and } u \ge 0 \text{ on } (0,1)\},\$$

and

$$J(u) = \int_0^1 \left(\frac{1}{2} u'(y)^2 + 2j\sqrt{u(y)}\right) dy,$$

and then we call variational solution of (13) to a function  $u \in K$  such that

$$J(u) = \min_{v \in K} \left\{ \int_0^1 \left( \frac{1}{2} v'(y)^2 + 2j\sqrt{v(y)} \right) dy \right\}.$$
 (15)

As said before, one of the reasons why the singular problem is quite relevant in the applications is for the possible occurrence of a *free boundary* (according to the values of j > 0) associated to a solution u of (13). In the one-dimensional setting, the *free boundary* is given simply by a point  $\xi \in [0,1)$  such that

$$\begin{cases} u(y) = 0 & \text{if } y \in [0, \xi], \\ u(y) > 0 & \text{if } y \in (\xi, 1], \text{ and } u'(\xi) = 0. \end{cases}$$
 (16)

Notice that the supplementary condition (14) corresponds to the case  $\xi = 0$ . Many authors used to say that, in that case, u is a "flat solution" (on y = 0). This is the more interesting case for us, since the electric fields should vanish on the cathode.

The treatment of free boundary solutions (not necessarily being a variational solution) leads to the notion of strong solutions with a free boundary:

**Definition 9** We say that  $u \in C^1([0,1)) \cap K$  is a strong solution with a free boundary if there exists  $\xi \in [0,1)$  such that (14) holds and, in addition,  $\frac{1}{\sqrt{u}} \in L^1(\xi,1)$ , and  $u''(y) = \frac{j}{\sqrt{u(y)}}$  a.e.  $y \in (\xi,1)$ .

Notice that, in this case, we must understand the differential equation as

$$-u''(y) + \frac{j}{\sqrt{u(y)}} \chi_{\{u>0\}} = 0 \quad y \in (0,1),$$

where  $\chi_{\{u>0\}}$  denotes the characteristic function of the set  $\{y \in (0,1) : u(y) > 0\}$ , since otherwise the singular term is not well defined on  $[0,\xi]$ .

The following result was communicated to me by Haïm Brezis on 2004. Although there are different similar treatments in the literature (see, e.g., [17]) the uniqueness part and the argument used in the proof is completely new.

**Theorem 10** (H. Brezis) There exists a unique variational solution. Moreover, if we define  $j^* = \frac{4}{9}$ , then:

a) if  $j = j^*$  the variational solution is a flat solution and it is given by

$$u(y) = y^{4/3}. (17)$$

b) if  $j > j^*$  the variational solution is a free boundary solution with

$$\xi = 1 - \frac{1}{\sqrt{\frac{9}{4}j}},\tag{18}$$

and it is given by  $u(y) = A(y - \xi)_+^{4/3}$  with  $A = (\frac{9}{4}j)^{2/3}$ . c) if  $0 < j < j^*$  then the variational solution is such that u > 0 on (0,1] and  $u'(0) = K_0 > 0$ , for some  $K_0 = K_0(j)$ .

**Remark 11** In fact, from the proof we will see that if  $j \to +\infty$  then  $\xi \to 1$  (a sort of limit of "boundary layer" type). Moreover, if  $j \to +0$  then  $u(y) \to y$  in  $C^0([0,1])$ .

*Proof.* Let us start by proving the uniqueness of variational free boundary type solutions. By well-known results, (15) admits always a minimizer which satisfies  $u'' = \frac{j}{\sqrt{u}}$  on the set  $[\xi < y < 1]$ . Then, by multiplying by u' we get

$$\begin{cases}
\frac{du}{dy} = \sqrt{4j} \left( u(y) \right)^{1/4} \\
u(1) = 1,
\end{cases}$$
(19)

a problem with the uniqueness of solutions since  $u^{1/4}$  is monotone increasing. By Leibniz's formula for separable ordinary differential equations, we get

$$u(y) = \left(\frac{3\sqrt{j}}{2}\right)^{4/3} (y - \xi)_{+}^{4/3},$$

which proves a) and b). Let us consider now the case of  $j \in (0, \frac{4}{9})$ . Note that if we arrive to prove that  $\frac{du}{du}(0) = K_0 > 0$  then after multiplying by u' we will get that

$$u'(y) = \sqrt{4j\sqrt{u(y)} + K_0^2},$$

and the Leibniz formula would lead to the implicit expression of the solution

$$\int_0^{u(y)} \frac{du}{\sqrt{4j\sqrt{u} + k_0^2}} = y. \tag{20}$$

A more direct way to apply the above argument is the following: let

$$H(\sigma) = \int_0^{\sigma} \frac{ds}{\sqrt{s^{1/2} + 1}}, \quad \text{for } \sigma > 0.$$
 (21)

Then H(0) = 0 and

$$H'(\sigma) = \frac{1}{\sqrt{\sqrt{\sigma} + 1}} > 0. \tag{22}$$

Thus H can be inverted,  $\sigma = H^{-1}(\tau)$ , and we get that  $H^{-1}(0) = 0$  and

$$\frac{d}{d\tau}H^{-1}(\tau) = \sqrt{\sqrt{H^{-1}(\tau)} + 1} > 0. \tag{23}$$

Let us prove that any variational solution can be written in the form  $u(y) = \frac{1}{A}H^{-1}(By)$ , for suitable constants A and B. Note that

$$u'(y) = \frac{B}{A}\sqrt{\sqrt{Au(y)} + 1} \quad \text{and thus } u'(0) = \frac{B}{A}. \tag{24}$$

Indeed, since H(Au(y)) = By we get

$$H'(Au)A\frac{du}{dy} = B,$$

i.e.,

$$\frac{1}{\sqrt{\sqrt{Au(y)}+1}}A\frac{du}{dy} = B,$$

which implies (24). Moreover

$$\frac{d^{2}u}{dy^{2}}(y) = \frac{B}{A} \frac{1}{2} \left( \sqrt{Au(y)} + 1 \right)^{-1/2} \frac{1}{2} \left( Au(y) \right)^{-1/2} A \frac{du}{dy}(y) 
= \frac{B}{A} \frac{1}{2} \left( \sqrt{Au(y)} + 1 \right)^{-1/2} \frac{1}{2} \left( Au(y) \right)^{-1/2} B \sqrt{\sqrt{Au(y)} + 1} 
= \frac{B^{2}}{4A} \frac{1}{\sqrt{Au(y)}} 
= \frac{B^{2}}{4A^{3/2}} \frac{1}{\sqrt{u(y)}}.$$
(25)

Thus, we must determine A and B by the conditions  $\frac{B^2}{4A^{3/2}} = j$  and u(1) = 1. Then we get the condition

$$H(A) = B = 2\sqrt{j}A^{3/4},\tag{26}$$

which has a unique solution A (and then a unique value of B) since H is strictly increasing, H(0) = 0 and  $\lim_{\sigma \to +\infty} H(\sigma) = +\infty$ . Then

$$u'(0) = \frac{B_0}{A_0} = \frac{H(A_0)}{A_0},$$

with  $A_0 > 0$  solution of

$$\frac{H(A_0)}{A_0^{3/4}} = 2\sqrt{j}.$$

The uniqueness of the constants A and B also proves the uniqueness of the variational solution, and part c) follows.

**Remark 12** In cases a) and b) we can improve the regularity: we have that  $u \in W^{2,p}(\xi,1) \quad \forall p \in [1,\frac{3}{2})$ , and in case c)  $u \in W^{2,p}(0,1) \quad \forall p \in [1,2)$ . In fact, we have the *sharp gradient estimate* 

$$|u'(y)| \le Cu^{1/4}(y)$$
, for some  $C > 0$ , for any  $y \in (0, 1)$ . (27)

Estimates of this nature play an important role in the study of the existence of solutions of the associated parabolic problem (the so-called "quenching problem"): see, e.g., [42] and [14], among others.

**Remark 13** Note also that  $j^* = \sup\{j > 0 : u'(0) > 0 \text{ with } u \text{ variational solution of (15)}\}$ . Something similar was already mentioned in the space charge literature (see, e.g., [37] p. 381 and [39] p. 2372).

#### 3.2 The distributed one-dimensional current density

For different purposes, it is useful to consider a similar problem to the above formulation but now for some j = j(y) with  $j \in L^1_{loc}(0,1), j \ge 0$ . The problem under consideration is

$$\begin{cases}
-u''(y) + \frac{j(y)}{\sqrt{u(y)}} \chi_{\{u>0\}} = 0 & y \in (0,1), u \ge 0 \\
u \ge 0 & y \in (0,1), \\
u(0) = 0 & u(1) = 1,
\end{cases}$$
(28)

where  $\chi_{\{u>0\}}$  denotes again the characteristic function of the set  $\{y \in (0,1) : u(y) > 0\}$ . Since the nonlinear term is decreasing, we will apply the method of super and subsolutions.

**Definition 14** Given  $p \ge 1$ , a function  $\overline{u} \in H^1(0,1)$  is a p-strong supersolution of problem (28), if

i) 
$$\overline{u} \geq 0$$
 and  $\frac{j}{\sqrt{\overline{u}}} \in L^p(0,1)$ ,

$$ii) \ \overline{u}'' \in L^1(0,1),$$

$$iii) - \overline{u}'' + \frac{j}{\sqrt{\overline{u}}} \chi_{\{\overline{u} > 0\}} \ge 0 \text{ a.e. in } (0, 1),$$

$$iv) \ \overline{u}(0) = 0 \ and \ \overline{u}(1) \ge 1$$

The notion of p-strong subsolution  $\underline{u} \in H^1(0,1)$  is similar, by replacing the inequalities  $\geq$  in iii) and iv) by the inequality  $\leq$ , but always with  $\underline{u}(0) = 0$ . We recall here a variation of the iterative method of super and subsolutions (see, e.g., the exposition and general ideas in the monograph [41]) which applies to our framework with a singular absorption. The proof is a special case of a more general statement, which will be given later (see Theorem 23 and Remark 24).

**Theorem 15** Let  $j \in L^1_{loc}(0,1), j \geq 0$ . Assume that there exists p > 1, a p-strong supersolution  $u^0$  and a p-strong subsolution  $u_0$ , such that

$$0 < u_0(y) \le u^0(y) \text{ a.e. } y \in (0,1).$$
(29)

Then, problem (28) has a maximal solution  $u^*$  and a minimal solution  $u_*$  on the "interval"  $[u_0, u^0]$  of  $H^1(0,1)$ , i.e., such that

$$u_0 \le u_* \le u^* \le u^0 \text{ a.e. } y \in (0,1).$$
 (30)

Moreover  $u''_*$ ,  $u''^* \in L^p(0,1)$ .

We can consider some special cases of j(y) that allow to get some results in the line of Theorem 10:

$$j(y) = \lambda y^q \text{ with } q \in \left(-\frac{1}{2}, 1\right), \ \lambda > 0.$$
(31)

Note that, obviously, q = 0 corresponds to the case treated before and that our study will consider cases in which j(0) = 0  $(q \in (0,1))$  as well as cases in which  $j(0) = +\infty$   $(q \in (-\frac{1}{2},0))$ .

**Theorem 16** Assume (31). Then if we define  $\lambda_q^* = \frac{2(1+2q)(2+q)}{9}$  we have: a) if  $\lambda = \lambda_q^*$  the function  $u(y) = y^{(4+2q)/3}$  is a flat solution of the problem.

b) if 
$$\lambda > \lambda_q^*$$
 the function  $u(y) = A(y - \xi)_+^{(4+2q)/3}$  is a free boundary solution with  $\xi = 1 - \frac{1}{\left[\frac{\lambda}{\lambda_s^*}\right]^{1/(2+q)}}$ ,

and  $A = (\frac{\lambda}{\lambda_a^*})^{2/3}$ . Moreover,

$$\frac{j(y)}{\sqrt{u(y)}}\chi_{\{u>0\}} = \frac{\lambda}{\sqrt{A}}(y-\xi)^{-\frac{(2-2q)}{3}}\chi_{\{y>\xi\}} \in L^1(0,1).$$

c) if  $0 < \lambda < \lambda_q^*$ , then the unique solution u is such that u > 0 on (0,1] and  $u'(0) = K_0 > 0$ , for some  $K_0 = K_0(\lambda)$ .

Remark 17 If we use a different notation,  $\beta = -q$  and  $\alpha = (4 - 2\beta)/3$ , then we get the existence of solutions of the form  $u(y) = y^{\alpha}$ , with  $\alpha \in (1, 4/3)$ , once we assume  $\beta \in (0, 1/2)$ . Curiously enough, the constraint  $3\alpha/2 + \beta = 2$  arises also in Theorem 1 dealing with the two-dimensional problem (see Section 4).

The key idea of the proof is to build a family of explicit solutions of the differential equation inspired by Chapter 2 of [18].

**Lemma 18** Let  $u_m(y) = Cy^{\frac{2}{1+m}}$ , with C > 0 and  $m \in (0,1)$ . Then

$$-u_m''(y) + \frac{j_m(y)}{\sqrt{u_m(y)}} = 0, (32)$$

where

$$j_m(y) = \frac{2C\sqrt{C}(1-m)}{(1+m)^2} y^{\frac{(1-2m)}{(1+m)}} \quad y \in (0,1).$$
(33)

*Proof.* It suffices to check that

$$(u_m)'(y) = C \frac{2}{1+m} y^{\left(\frac{2}{1+m}-1\right)} = \frac{2C}{1+m} y^{\frac{1-m}{1+m}},$$

$$(u_m)''(y) = \frac{2C(1-m)}{(1+m)^2} y^{\left(\frac{1-m}{1+m}-1\right)} = \frac{2C(1-m)}{(1+m)^2} y^{\frac{-2m}{1+m}}.$$
(34)

Since  $\sqrt{u_m(y)} = \sqrt{C}y^{\frac{1}{1+m}}$ , we get

$$-(u_m)''(y) = \frac{-j_m(y)}{\sqrt{u_m(y)}} = \frac{-1}{\sqrt{C}} y^{\frac{-1}{1+m}} j_m(y).$$

Applying (34) we arrive at (32) with  $j_m(y)$  given by (33).

Corollary 19 Given  $j(y) = \lambda y^q$  with  $q \in (-\frac{1}{2}, 1)$  and  $\lambda > 0$ , the function  $u_q(y) = C_q(\lambda)y^{\frac{4+2q}{3}}$ , with

$$C_q(\lambda) = \left[ \frac{9\lambda}{2(1+2q)(2+q)} \right]^{2/3},$$
 (35)

is a solution of

$$-u_q'' + \frac{\lambda y^q}{\sqrt{u_q}} = 0$$
, in  $(0,1)$ .

*Proof.* We impose  $\frac{(1-2m)}{(1+m)} = q$  and

$$\frac{2C\sqrt{C}(1-m)}{(1+m)^2} = \lambda.$$

Then we get  $m=\frac{1-q}{2+q}$  and  $C(\lambda)$  given by (35). Then the conclusion is a direct consequence of Lemma 18.

Proof of Theorem 16. Conclusions a) and b) are a direct consequence of Corollary 19. To prove part c) it suffices to build a subsolution  $\underline{u}(y) \leq y$  satisfying the searched properties, since  $\overline{u}(y) = y$  is a

supersolution satisfying that  $\underline{u}(y) \leq \overline{u}(y)$  a.e.  $y \in (0,1)$ , and then, by Theorem 15, there exists a minimal solution  $u_*$  of (28) satisfying

$$\underline{u}(y) \le u_*(y) \le \overline{u}(y)$$
 a.e.  $y \in (0,1)$ .

Then if  $\underline{u}'(0) > 0$  we conclude, as desired, that  $u'_*(0) > 0$ .

A construction of a suitable subsolution is the following: let  $\varepsilon > 0$  be small enough, to be determined later, let  $u_{\#}(y) = y^{(4+2q)/3}$  and define

$$\underline{u}(y) = \frac{u_{\#}(y) + \varepsilon y}{1 + \varepsilon}.$$

Then it is clear that  $\underline{u}(0) = 0$ ,  $\underline{u}(1) = 1$  and since  $\underline{u}'(y) = \frac{u'_{\#}(y) + \varepsilon}{1 + \varepsilon}$  we get  $\underline{u}'(0) = \frac{\varepsilon}{1 + \varepsilon} > 0$ . Moreover,

$$\underline{u}''(y) = \frac{u_{\#}''(y)}{1+\varepsilon} = \frac{\lambda_q^* y^q}{(1+\varepsilon)\sqrt{u_{\#}(y)}}.$$

Then, concerning the differential equation, we can have the condition for subsolution

$$-\underline{u}''(y) + \frac{\lambda y^q}{\sqrt{\underline{u}(y)}} = -\frac{\lambda_q^* y^q}{(1+\varepsilon)\sqrt{u_\#(y)}} + \frac{\sqrt{1+\varepsilon}\lambda y^q}{\sqrt{u_\#(y)+\varepsilon y}} \le 0$$

if we have

$$\frac{\sqrt{1+\varepsilon\lambda}}{\sqrt{u_{\#}(y)+\varepsilon y}} \le \frac{\lambda_q^*}{(1+\varepsilon)\sqrt{u_{\#}(y)}}.$$
(36)

But, obviously

$$\frac{\lambda_q^*}{(1+\varepsilon)\sqrt{u_\#(y)+\varepsilon y}} \le \frac{\lambda_q^*}{(1+\varepsilon)\sqrt{u_\#(y)}}.$$

Then we arrive at the searched inequality (36) if we have

$$\lambda \le \frac{\lambda_q^*}{(1+\varepsilon)^{3/2}}.$$

This is clearly true since  $\lambda < \lambda_q^*$ : it suffices to take  $\varepsilon > 0$  such that

$$\varepsilon < \left(\frac{\lambda_q^*}{\lambda}\right)^{2/3} - 1.$$

The functions  $u_q(y)$  given in Corollary 19 can be used as super and subsolutions to get the existence of a flat solution for a more general function j(y). The result that follows is just a sample of many other possible results.

**Theorem 20** Let  $-\frac{1}{2} < r < q < 0$ . Assume

$$\lambda_r y^r \ge j(y) \ge \lambda_q y^q$$
 a.e.  $y \in (0,1)$ ,

with

$$\lambda_q = \lambda_q^* \text{ and } \lambda_r > \lambda_r^* \text{ with } \lambda_r \text{ large enough,}$$
 (37)

or

$$\lambda_q < \lambda_q^* \text{ and } \lambda_r = \lambda_r^* \text{ with } \lambda_q \ge 0 \text{ small enough.}$$
 (38)

Then there exists a maximal solution  $u^*$  and a minimal solution  $u_*$  of problem (28). Moreover, in case of (37)

$$A_r(y-\xi_r)_+^{\frac{4+2r}{3}} \le u_*(y) \le u^*(y) \le y^{\frac{4+2q}{3}}$$
 a.e.  $y \in (0,1)$ ,

for some  $A_r > 0$  and  $\xi_r \in (0,1)$ , and in case (38)

$$y^{\frac{4+2r}{3}} \le u_*(y) \le u^*(y) \le u_q(y)$$
 a.e.  $y \in (0,1)$ ,

with  $u_q(y)$  the solution of problem (28) corresponding to  $j(y) = \lambda_q y^q$  (when  $\lambda_q \in (0, \lambda_q^*)$ ) mentioned in part c) of Theorem 16. Moreover if  $\lambda_q = 0$  we can take as  $u_q$  the function  $u_q(y) = y$ .

*Proof.* We recall that for  $a > b \ge 0$  we have  $0 < y^a < y^b < 1$  for any  $y \in (0,1)$ . Thus, equivalently, if  $-a < -b \le 0$  then  $y^{-a} > y^{-b} > 1$  for any  $y \in (0,1)$ .

We take now -a = r and -b = q so  $y^r > y^q$ . The conclusion will be obtained through the application of Theorem 16 to different choices of the super and subsolution. In case (37) we take as supersolution  $u^0(y) = y^{\frac{4+2q}{3}}$  since

$$-u^{0"}(y) + \frac{j(y)}{\sqrt{u^0(y)}} \ge -u^{0"}(y) + \frac{\lambda_q^* y^q}{\sqrt{u^0(y)}} = 0.$$

On the other hand, as a subsolution we can take  $u_0(y) = A_r(y - \xi_r)_+^{\frac{4+2r}{3}}$  for some  $A_r > 0$  and  $\xi_r \in (0,1)$  as indicated in part b) of Theorem 16 (remember that  $\lambda_r > \lambda_q^*$ ) since

$$-u_0''(y) + \frac{j(y)}{\sqrt{u_0(y)}} \le -u_0''(y) + \frac{\lambda_r y^r}{\sqrt{u_0(y)}} = 0.$$

We have  $u_0(0) = u^0(0) = 0$ ,  $u_0(1) = u^0(1) = 1$ . Moreover, if  $\lambda_r$  is large enough we have

$$u_0(y) = A_r(y - \xi_r)^{\frac{4+2r}{3}} < y^{\frac{4+2q}{3}} = u^0(y)$$
 for any  $y \in (0,1)$ ,

(in spite that  $0 < \frac{4+2r}{3} < \frac{4+2q}{3}$  and thus  $y^{\frac{4+2q}{3}} < y^{\frac{4+2r}{3}}$  for  $y \in (0,1)$ ). Indeed, we can take  $A_r > 0$  and  $\xi_r \in (0,1)$  so that  $u_0'(1) > u^{0'}(1) = \frac{4+2q}{3}$  and thus both functions are well-ordered. In case (38) we take as subsolution the function  $u_0(y) = y^{\frac{4+2r}{3}}$  and as supersolution the function  $u^0(y) = y^{\frac{4+2r}{3}}$ 

In case (38) we take as subsolution the function  $u_0(y) = y^{\frac{q+2r}{3}}$  and as supersolution the function  $u^0(y) = u_q(y)$  the solution of problem (28) corresponding to  $j(y) = \lambda_q y^q$  when  $\lambda_q \in (0, \lambda_q^*)$  mentioned in part c) of Theorem 16. The conditions in terms of the differential equation are satisfied, as before. Moreover, again,  $u_0(0) = u^0(0) = 0$ ,  $u_0(1) = u^0(1) = 1$ , and the inequality  $u_0(y) = y^{\frac{4+2r}{3}} < u_q(y) = u^0(y)$  for any  $y \in (0,1)$ , holds once that  $\lambda_q > 0$  is small enough (since we know that  $u_q(y) \to y$  when  $\lambda_q \to 0$ ). If  $\lambda_q = 0$  the function  $u_q(y) = y$  is a supersolution and the proof ends.

**Remark 21** The gradient estimate mentioned when j is a constant is no longer valid when  $j(y) = \lambda y^{-\beta}$  with  $\beta \in (0, \frac{1}{2})$ . If  $\lambda \ge \lambda_{\beta}$ , for some  $\lambda_{\beta} > 0$ 

$$|u'(y)| \le Cu^{\frac{1+2\beta}{4+2\beta}}(y)$$
, for some  $C > 0$ , for any  $y \in (0,1)$ . (39)

# 4 The 2-d parallel-plate geometry

As mentioned in the Introduction, Theorem 1 makes precise, a conjecture by A. Rokhlenko [44]: if j(x) behaves as  $A/|x|^{\beta}$ , for some  $\beta \in (0, 1/2)$ , for  $x \in (-a, 0)$  then there exists a weak solution u(x, y) of (3), and u "behaves" (near the cathode  $[-a, a] \times \{0\}$ ) as  $y^{\alpha}$  for some  $\alpha \in (1, 4/3)$ , with  $\alpha = 4/3 - 2\beta$ . We will obtain other auxiliary results for the more general case of  $j: (-a, b) \to [0, +\infty)$  such that

$$\begin{cases} j(x) > 0 & \text{if } x \in (-a,0), \ j \in L^1_{loc}(-a,0), \\ j(x) = 0 & \text{if } x \in (0,b), \end{cases}$$
(40)

with a possible singularity of the current density function j(x) only expected at the origin x = 0. Our main goal is, therefore, to get the transition, near x = 0, between flat and linear profiles u(.,y).

As mentioned before, since in other frameworks the solutions may vanish in some parts of the spatial domain (dead cores), we can reformulate the PDE as

$$P_{a,b,j} = \begin{cases}
-\Delta u + \frac{j(x)}{\sqrt{u}} \chi_{\{u>0\}} = 0 & x \in (-a,b), \ y \in (0,1), \\
u(x,0) = 0 & x \in (-a,b), \\
u(x,1) = 1 & x \in (-a,b), \\
u(-a,y) = y^{4/3} & y \in (0,1), \\
u(b,y) = y & y \in (0,1).
\end{cases}$$
(41)

This Section will be structured in several subsections. In subsection 4.1 we present the adaptation of the super and subsolutions method to the problem  $P_{a,b,j}$ . The construction of positive subsolutions satisfying the additional conditions  $AC_{a,b}$  will be presented in subsection 4.2. The proof is made by constructing suitable super and subsolutions for several auxiliary problems and matching suitably the corresponding solutions. In particular, we will use the global bifurcation diagram (in terms of the parameter  $\lambda$ ) associated to the auxiliary problem (9). In subsection 4.3 we will end the proof of Theorem 1 by constructing a suitable supersolution which is flat on  $(-a,0) \times \{0\}$ . Finally, in subsection 4.4 we will prove the uniqueness of non-degenerate solutions.

### 4.1 On the existence of solutions of $P_{a,b,j}$ via super and subsolutions method

Due to the singularity of the nonlinear term, the super and subsolutions method needs to be applied under some adequate conditions. Let  $\Omega = (-a, b) \times (0, 1)$  and j as in (40).

**Definition 22** A function  $u^0 \in W^{1,1}(\Omega)$ , is said a p-positive supersolution of  $P_{a,b,j}$  if  $u^0 \ge 0$ ,  $\frac{j}{\sqrt{u^0}} \in L^p(\Omega)$ , for some  $p \ge 1$ , and it verifies in a very weak sense that

$$\begin{cases}
-\Delta u^{0} \ge -\frac{j(x)}{\sqrt{u^{0}}} & in \Omega, \\
u^{0}(x,0) \ge 0 & x \in (-a,b), \\
u^{0}(x,1) \ge 1 & x \in (-a,b), \\
u^{0}(-a,y) \ge y^{4/3} & y \in (0,1), \\
u^{0}(b,y) \ge y & y \in (0,1).
\end{cases}$$

The notion of p-positive subsolution  $u_0$  is introduced similarly, i.e.,  $u_0 \in W^{1,1}(\Omega)$  with  $u_0 > 0$  and  $\frac{j}{\sqrt{u_0}} \in L^p(\Omega)$ , satisfies that

$$\begin{cases}
-\Delta u_0 \le -\frac{j(x)}{\sqrt{u_0}} & in \Omega, \\
u_0(x,0) = 0 & x \in (-a,b), \\
u_0(x,1) \le 1 & x \in (-a,b), \\
u_0(-a,y) \le y^{4/3} & y \in (0,1), \\
u^0(b,y) \le y & y \in (0,1).
\end{cases}$$

Theorem 23 Assume that

$$\begin{cases}
\text{there exists } p > 1, \text{ a } p-positive supersolution } u^0 \text{ and a } p-positive subsolution } u_0 \\
\text{of } P_{a,b,j} \text{ such that } u_0 \leq u^0 \text{ a.e. in } \Omega.
\end{cases}$$
(42)

Then problem  $P_{a,b,j}$  possesses a minimal and a maximal solutions  $u_*$  and  $u^*$  in the interval  $[u_0, u^0]$ , i.e.,

$$u_0 \le u_* \le u^* \le u^0$$
 a.e.in  $\Omega$ .

*Proof.* We define the iterative schemes (starting with  $u^0$  and  $u_0$ )

$$\begin{cases}
-\Delta u^n = -\frac{j(x)}{\sqrt{u^{n-1}}} & \text{in } \Omega, \\
u^n(x,0) = 0 & x \in (-a,b), \\
u^n(x,1) = 1 & x \in (-a,b), \\
u^n(-a,y) = y^{4/3} & y \in (0,1), \\
u^n(b,y) = y & y \in (0,1),
\end{cases}$$

and

$$\begin{cases}
-\Delta u_n = -\frac{j(x)}{\sqrt{u_{n-1}}} & \text{in } \Omega, \\
u_n(x,0) = 0 & x \in (-a,b), \\
u_n(x,1) = 1 & x \in (-a,b), \\
u_n(-a,y) = y^{4/3} & y \in (0,1), \\
u_n(b,y) = y & y \in (0,1).
\end{cases}$$

By using the comparison principle for the Laplace operator, we get that

$$0 < u_0 \le u_1 \le ... \le u_n \le ... \le u^n \le ... \le u^1 \le u^0$$
 a.e. in  $\Omega$ ,

and so the sequences  $\{u^n\}, \{u_n\}$  converge (monotonically) in  $L^p(\Omega)$  to some functions  $u_*$  and  $u^*$  and the sequences  $\{\frac{j}{\sqrt{u^{n-1}}}\}, \{\frac{j}{\sqrt{u_{n-1}}}\}$  are bounded in  $L^p(\Omega)$  and converge also (monotonically) in  $L^p(\Omega)$ .

Remark 24 With some minor modifications, the above result holds when the super and subsolutions are in the weighted space  $\frac{j}{\sqrt{u^0}} \in L^p(\Omega, \delta)$ , for some  $p \geq 1$ , with  $\delta = d((x, y), \partial\Omega)$ . See, e.g., [8], [47], [30] and [40] among many other papers (see also some applications in [1]). This allows a greater generality to treat the singular term: for instance, the function  $u^0(x, y) = y$  is such that  $\frac{1}{\sqrt{u^0}} \in L^p(\Omega)$  for  $p \in [1, 2)$ ,  $\frac{1}{\sqrt{u^0}} \notin L^2(\Omega)$  but  $\frac{1}{\sqrt{u^0}} \in L^2(\Omega, \delta)$ .

Remark 25 By some standard approximating arguments, it is well-known that the above notion of p-super and subsolutions of  $P_{a,b,j}$  can be extended to the case in which the diffusion term generates an additional distribution over a simple curve  $\Gamma$  separating  $\Omega$  in two different parts (matching without a  $W^{1,p}$ -contact: see [36] and [4]). Then, for instance, the subsolution  $u_0$  of problem  $P_{a,b,j}$  is allowed to satisfy

$$-\Delta u_0 + \frac{j(x)}{\sqrt{u_0}} \le 0 \text{ in } \mathcal{D}'(\Omega).$$

The existence of a (not-flat) supersolution  $u^0$  can be easily proved.

**Lemma 26** Let j(x) satisfying (40). Then the function  $u^0(x,y) = y$  is a positive p-supersolution of  $P_{a,b,j}$  for any  $p \in [1,2)$ .

*Proof.* It is a trivial fact since

$$-\Delta u^0 = 0 \ge -\frac{j(x)}{\sqrt{u^0}} \quad x \in (-a, b), \ y \in (0, 1),$$

and

$$\begin{cases} u^{0}(x,0) = 0 & x \in (-a,b), \\ u^{0}(x,1) = 1 & x \in (-a,b), \\ u^{0}(-a,y) \ge y^{4/3} & y \in (0,1), \\ u^{0}(b,y) = y & y \in (0,1). \end{cases}$$

Moreover,  $\frac{j(x)}{\sqrt{u^0}} \in L^p(\Omega)$  for any  $p \in [1, 2)$ .

The construction of a positive subsolution is a very delicate task, which will be presented in the following subsection. Before proving Theorem 1 we can get an existence and uniqueness result of a positive solution under a more general assumption on j(x) than in Theorem 1 but without ensuring that it is flat on  $(-a, 0) \times \{0\}$ . We define

$$\delta(x, y) := \operatorname{dist}((x, y), \partial\Omega),$$

which sometimes we shall denote simply as  $\delta$ .

**Theorem 27** Assume that j(x) satisfies

$$j(x) = \frac{A}{(-x)^{\beta}}$$
, for  $x \in (-a, 0)$  and  $j(x) = 0$  if  $x \in (0, b)$ ,

with

$$0 < \beta < 1/2, \beta \text{ and } A > 0 \text{ small enough.}$$
 (43)

Then there exists a weak solution  $u \in L^2(\Omega; \delta)$  of

$$P_{a,b,j} = \begin{cases}
-\Delta u + \frac{j(x)}{\sqrt{u}} = 0 & x \in (-a,b), \ y \in (0,1), \\
u(x,0) = 0 & x \in (-a,b), \\
u(x,1) = 1 & x \in (-a,b), \\
u(-a,y) = y^{4/3} & y \in (0,1), \\
u(b,y) = y & y \in (0,1),
\end{cases}$$
(44)

such that

$$C\delta(x,y)^{4/3} \le u(x,y) \le y \text{ a.e. } (x,y) \in (-a,b) \times (0,1),$$
 (45)

for some C > 0, and  $Cy^{\alpha} \le u(0, y), y \in (0, 1)$ , with  $\alpha = \frac{2}{3}(2 - \beta)$ .

The solutions satisfying inequalities of the type (45) are called *nondegenerate* solutions. The uniqueness of solutions given in Theorem 27 is a consequence of the techniques introduced in the paper [21]. Indeed, let  $\nu \in (0, 4/3]$ , and define the class of functions

$$\mathcal{M}(\nu) := \Big\{ u \in L^2(\Omega; \delta) \mid \text{such that } u(x, y) \ge C\delta(x, y)^{\nu} \quad \text{in } \Omega, \text{ for some } C > 0 \Big\}.$$
 (46)

We have

**Theorem 28** Assume j(x) as in Theorem 27. Then, there exists at most a solution  $u \in \mathcal{M}(\nu)$  of  $P_{a,b,j}$ .

The proof will be obtained through some smoothing estimates for some suitable parabolic problem (see subsection 4.4). Finally, the proof of the main result of this paper (Theorem 1) will be a direct consequence of the method of super and subsolutions, and the following result on partially flat supersolutions:

**Theorem 29** There exist  $A_0, b_0 > 0$  and  $\beta_0 \in (0, \frac{1}{2})$  such that, if  $b \ge b_0 > 0$ , and if we assume (7) then there exists a partially flat supersolution  $\overline{u}(x,y)$  of problem  $P_{a,b,j}$ , i.e., such that  $\overline{u} \in L^2(\Omega; \delta)$ ,

$$\begin{cases}
-\Delta \overline{u} + \frac{j(x)}{\sqrt{\overline{u}}} \ge 0 & x \in (-a, b), \ y \in (0, 1), \\
\overline{u}(x, 0) \ge 0 & x \in (-a, b), \\
\overline{u}(x, 1) \ge 1 & x \in (-a, b), \\
\overline{u}(-a, y) \ge y^{4/3} & y \in (0, 1), \\
\overline{u}(b, y) \ge y & y \in (0, 1),
\end{cases}$$
(47)

and

$$u(x,y) \le \overline{u}(x,y) \le C\delta(x,y)^{\overline{\alpha}} \text{ a.e. } (x,y) \in (-a,0) \times (0,1), \tag{48}$$

for some C > 0, with  $\overline{\alpha} \in (1, \frac{4}{3})$  and u(x, y) the solution given in Theorem 27.

**Remark 30** In order to have the zero flux condition on (-a,0) we need some special unbounded behaviour on j(x) near x = 0. Given  $x_0 \in (-a,0)$  we can construct (as in [3], [22], [18]) a "barrier function" of the form

$$\overline{u}(x,y:x_0) = K \|(x,y) - (x_0,0)\|^{4/3} = K[(x-x_0)^2 + y^2]^{2/3},$$
(49)

for a suitable K>0 in such a way that  $\overline{u}$  let a local supersolution once we assume

$$j(x) \ge C(-x)_{+}^{-\delta}, \ x \in (-a, a)$$

for some suitable  $\delta > 0$ . More precisely we can assume

$$j(x) \ge 4 \max(1, \frac{\sqrt{2}}{\sqrt{-x}}), \ x \in (-a, 0) \ \text{and} \ j(x) = 0 \text{ for } x \in (0, a),$$
 (50)

and that

$$j(x)[(x-x_0)^2+y^2]^{-1/3} \in L^p(\Omega_{x_0})$$
, for some  $p>1$ ,

where  $\Omega_{x_0} = \{(x,y) \in (-a,0) \times (0,1) : (x-x_0)^2 + y^2 < (-x_0)^2\}$ , for any  $x_0 \in (-a,0)$ . Assume also that there exists a p-nonnegative subsolution  $u_0$  such that  $u_0(x,y) \leq y$  a.e. in  $\Omega$ . Then, if u satisfies the problem in the interval  $[u_0,y]$ , i.e., such that

$$u_0 \le u \le y$$
 a.e.in  $\Omega$ 

we have that

$$u(x,y) \le \max(1, \frac{2^{1/3}}{(-x_0)^{1/3}})[(x-x_0)^2 + y^2]^{2/3},$$
 (51)

for any  $(x,y) \in (-a,0) \times (0,1)$  and  $x_0 \in (-a,0)$  such that  $(x-x_0)^2 + y^2 < (-x_0)^2$ . In particular  $\frac{\partial u}{\partial y}(x,0) = 0$  for  $x \in (-a,0)$ . The difficulty in this approach is to prove the positivity of the solution.

# 4.2 On the construction of some strict positive subsolutions with zero flux on $(-a,0) \times \{0\}$

The key point in the proof of Theorem 27 is the construction of a flat positive subsolution. We start by reducing the difficulty of this task by splitting the domain into two different subdomains. We define the subsets

$$\Omega_{-} := \{(x,y) \in \Omega : x \in (-a,0)\}, \text{ and } \Omega_{+} := \{(x,y) \in \Omega : x \in (0,b)\}.$$

We will need to introduce an artificial boundary condition on the points (0, y),  $y \in (0, 1)$ , corresponding to the external boundary of this half of the cathode. We will prove a rigorous result that, in some sense, is connected to the study made in [44] by using asymptotic techniques and numerical analysis. Obtaining the correct profile function in the external boundary condition, at x = 0, is already an important conclusion, since several choices were already considered in many papers in the literature on space charge problems. It was expected that the correct behaviour at the external border of the cathode is greater than the profile  $y^{4/3}$  in the middle of the cathode, corresponding to the Child-Langmuir law, but no concrete function has been proposed for this external profile. The precise artificial boundary condition on the exterior of the cathode, which we will introduce (as a Dirichlet boundary condition), is the following:

$$u(0,y) = hy^{\alpha}, \ y \in (0,1), \text{ for some } h \in (0,1] \text{ and for some } \alpha \in (1,4/3).$$
 (52)

The following result will simplify the task of finding a global positive subsolution since it will allow us to pass from the study of a discontinuous absorption coefficient j(x) to a problem with a strictly positive one.

**Proposition 31** Let j(x) satisfying (40). Let  $\alpha \in (1,4/3)$ . Given  $h \in (0,1]$ , consider the problem on  $\Omega_{-}$ 

$$P_{a,0,j} = \begin{cases} -\Delta u_{-} + \frac{j(x)}{\sqrt{u_{-}}} = 0 & in \Omega_{-}, \\ u_{-}(-a, y) = y^{4/3} & y \in (0, 1), \\ u_{-}(0, y) = hy^{\alpha} & y \in (0, 1), \\ u_{-}(x, 0) = 0 & x \in (-a, 0), \\ u_{-}(x, 1) = 1 & x \in (-a, 0). \end{cases}$$

$$(53)$$

Assume that there exists  $u_{0,-}(x,y)$ , a  $p_0$ -subsolution of problem  $P_{a,0,j}$ , for some  $p_0 \ge 1$ , such that

$$\frac{\partial u_{0,-}}{\partial x}(0,y) \le 0 \text{ for } y \in (0,1), \tag{54}$$

satisfying the additional conditions

$$AC_{a,0} = \begin{cases} \frac{\partial u_{0,-}}{\partial y}(x,0) = 0 & x \in (-a,0), \\ u_{0,-}(x,y) > 0 & x \in (-a,0), \ y \in (0,1). \end{cases}$$
 (55)

Then problem  $P_{a,b,j}$  has a  $p_0$ -subsolution  $\underline{u}$  satisfying the additional conditions  $AC_{a,b}$ .

Proof. Let  $u_+(x,y)$  be the unique classical solution  $u_+ \in C^2(\Omega_+) \cap C^0(\overline{\Omega_+})$  of the linear problem

$$P_{0,b,0} = \begin{cases}
-\Delta u_{+} = 0 & \text{in } \Omega_{+}, \\
u_{+}(0,y) = hy^{\alpha} & y \in (0,1), \\
u_{+}(b,y) = y & y \in (0,1), \\
u_{+}(x,0) = 0 & x \in (0,b), \\
u_{+}(x,1) = 1 & x \in (0,b).
\end{cases}$$
(56)

Define the function

$$\underline{u}(x,y) = \begin{cases} u_{-}(x,y) & \text{if } (x,y) \in \Omega_{-}, \\ u_{+}(x,y) & \text{if } (x,y) \in \Omega_{+}. \end{cases}$$

It is clear that  $\underline{u}$  is a continuous function  $\underline{u} \in C^0(\Omega)$  but its gradient has a discontinuity in the segment  $x = 0, y \in (0, 1)$ , since j(x) is discontinuous in that segment. All the boundary conditions of  $P_{a,b,j}$  are fulfilled and also the additional conditions  $AC_{a,b}$ . In order to check that  $\underline{u}(x, y)$  is a  $p_0$ -subsolution of problem  $P_{a,0,j}$  we will apply Corollary I.1 of [4]. To do this, if we define the segment  $\Gamma = \{(0, y), y \in (0, 1)\}$  then it suffices to check that

$$\frac{\partial u_{-}}{\partial \mathbf{n}} \le \frac{\partial u_{+}}{\partial \mathbf{n}} \quad \text{on } \Gamma, \tag{57}$$

where  $\mathbf{n}$  is the unit exterior normal vector to  $\Omega_{-}$ . In our case,  $\mathbf{n} = \mathbf{e_1}$  and then condition (57) is expressed as

$$\frac{\partial u_{-}}{\partial x}(0,y) \le \frac{\partial u_{+}}{\partial x}(0,y) \qquad y \in (0,1). \tag{58}$$

By the assumption (54), it suffices to check that

$$\frac{\partial u_+}{\partial x}(0,y) \ge 0 \text{ for } y \in (0,1).$$

To do that, let us consider the function  $U_{+}(x,y) = u_{+}(x,y) - y$ . Then

$$\begin{cases}
-\Delta U_{+} = 0 & \text{in } \Omega_{+}, \\
U_{+}(0, y) = y^{\alpha} - y & y \in (0, 1), \\
U_{+}(b, y) = 0 & y \in (0, 1), \\
U_{+}(x, 0) = 0 & x \in (0, b), \\
U_{+}(x, 1) = 0 & x \in (0, b).
\end{cases}$$
(59)

We can define, now, the auxiliary function,

$$\underline{U}(x,y) = (b-x)(hy^{\alpha} - y), (x,y) \in \Omega_{+}.$$

Then we get that

$$\begin{cases}
-\Delta \underline{U} \leq 0 & \text{in } \Omega_+, \\
\underline{U}(0,y) = \omega(y) = hy^{\alpha} - y & y \in (0,1), \\
\underline{U}(b,y) = 0 & y \in (0,1), \\
\underline{U}(x,0) = 0 & x \in (0,b), \\
\underline{U}(x,1) = 0 & x \in (0,b).
\end{cases}$$

Thus, by the maximum principle we get that  $\underline{U}(x,y) \leq U_+(x,y)$  on  $\Omega_+$ . But since  $\underline{U}(0,y) = U_+(0,y)$  and we have that  $\frac{\partial \underline{U}}{\partial x}(0,y) \geq 0$  for  $y \in (0,1)$ , we deduce that necessarily,  $\frac{\partial U_+}{\partial x}(0,y) \geq 0$  for  $y \in (0,1)$ , which leads to the required inequality.

According Proposition 31, to finish with the construction of the global subsolution  $u_0$ , we must justify the existence of a function  $u_{0,-}(x,y)$  solution of the nonlinear problem  $P_{a,0,j}$ , raised on  $\Omega_-$ , and to check that  $u_{0,-}(x,y)$  satisfies the additional conditions (55) and (54).

In order to construct the subsolution for the case of a possible unbounded j(x), satisfying that  $j(x) \leq \frac{A}{(-x)^{\beta}}$  on (-a,0), we will use some ideas coming from the study of Fluid Mechanics in the consideration of spatial domains with corners (see, e.g., [2]). We will try to find the subsolution  $u_{0,-}(x,y)$  in the form

$$u_{0,-}(x,y) = \phi(r,\theta) = kr^{\alpha}U(\theta), \text{ for some } k > 0, \alpha > 1,$$
(60)

where  $r = \sqrt{x^2 + y^2}$  and  $\theta = \arctan(y/x)$ . Then, the partial differential inequality becomes

$$-\triangle \phi + \frac{j(r\cos\theta)}{\sqrt{\phi}} = -\frac{1}{r}\frac{\partial}{\partial r}(r\frac{\partial \phi}{\partial r}) - \frac{1}{r^2}\frac{\partial^2 \phi}{\partial \theta^2} + \frac{j(r\cos\theta)}{\sqrt{\phi}} \le 0.$$
 (61)

We will assume  $r \in [0, R)$ , for a suitable R > 0, and  $\theta \in (\frac{\pi}{2}, \pi)$ . The additional conditions (55) will require  $\phi(r, \theta) > 0$  if r > 0 and

$$\phi(r,\pi) = \frac{\partial \phi}{\partial \theta}(r,\pi) = 0 \text{ for } r \in [0,R).$$

We make now the structural condition (40) on j(x) and thus

$$j(r\cos\theta) \le \frac{A}{(-r\cos\theta)^{\beta}}$$
, if  $\theta \in (\frac{\pi}{2}, \pi)$ , for some  $A > 0$  and  $\beta \in (0, 1)$ . (62)

The partial differential inequality (61) leads to the study of the ordinary differential equation

$$-U''(\theta) + \frac{V(\theta)}{\sqrt{U(\theta)}} = \lambda U(\theta) \quad \theta \in (\frac{\pi}{2}, \pi), \tag{63}$$

once we assume the constraint

$$\alpha + 2\beta = 4/3,\tag{64}$$

and then with

$$V(\theta) = \frac{A}{(-\cos\theta)^{\beta}} \text{ and } \lambda = \alpha^2.$$
 (65)

The complementary conditions  $AC_{a,0}$  become now

$$\left\{ \begin{array}{ll} U(\theta)>0 & \theta\in(\frac{\pi}{2},\pi), \\ U(\pi)=U'(\pi)=0. \end{array} \right.$$

Notice that the potential  $V(\theta)$  is singular only for  $\theta = \frac{\pi}{2}$ . On the other hand, we will need to match this subsolution with another function which is positive for  $\theta = \frac{\pi}{2}$ . Then, we will construct the subsolution in two different pieces (see Figures 4 and 8)

$$U(\theta) = \begin{cases} v_1(\theta) & \text{if } \theta \in (\frac{\pi}{2}, \pi - R_0), \\ v_2(\theta) & \text{if } \theta \in (\pi - R_0, \pi), \end{cases}$$

for some  $R_0 \in (0, \frac{\pi}{2})$ , and with  $v_2(\theta) \in [0, 1]$  such that

$$\begin{cases}
-v_2''(\theta) + \frac{V_0}{\sqrt{v_2(\theta)}} = \lambda v_2(\theta) & \theta \in (\pi - R_0, \pi), \\
v_2(\pi) = v_2'(\pi) = 0,
\end{cases}$$
(66)

where

$$V(\theta) \ge V_0 \text{ if } \theta \in (\frac{\pi}{2}, \pi - R_0).$$




Figure 4: Boundary layer

On the other hand,  $v_1(\theta)$  must take into account the singularity of the potential  $V(\theta)$  on the interval  $(\frac{\pi}{2}, \pi - R_0)$  and this will require a careful matching (see Figure 4). In addition, we must guarantee a good match with the function  $u_+(x,y)$  defined on  $\Omega_+$ , as indicated in Proposition 31. This means that we want to have

$$v_1'(\frac{\pi}{2}) \ge 0,$$

since we require  $\frac{\partial u_{0,-}}{\partial x}(0,y) \leq 0$  for  $y \in (0,1)$ , and from  $u_{0,-}(0,y) = kr^{\alpha}U(\theta)|_{\theta=\frac{\pi}{2}}$ , we have

$$\frac{\partial u_{0,-}}{\partial x}(0,y) = \dots = -Ar^{\alpha-1}U'(\theta)\big|_{\theta=\frac{\pi}{2}} \le 0.$$

Before presenting the details on the construction of  $v_1(\theta)$  and  $v_2(\theta)$ , it is very useful to consider the auxiliary nonlinear ODE of eigenvalue type (9) presented in the Introduction.

### 4.2.1 Bifurcation curve and flat solution for an auxiliary related nonlinear eigenvalue ordinary differential problem

Since the equation (66) can be understood as a nonlinear eigenvalue problem, it is useful to start by considering the following auxiliary related problem:

$$\begin{cases}
-U''(s) + \frac{V_0}{\sqrt{U(s)}} = \lambda U(s) & s \in (-R, R), \\
U(\pm R) = 0,
\end{cases}$$
(67)

where the positive constants  $V_0$  and R are given. The following result extends several results in the previous literature (see, e.g., [23], [27], [24] and some of the expositions made in [1]).

**Theorem 32** Given the positive constants  $V_0$  and R then:

- i) there is a bifurcation from infinity for  $\lambda$  near  $\lambda_1(R) = (\frac{\pi}{2R})^2$  (the first eigenvalue of the linear problem with  $V_0 = 0$ ),
  - ii) the bifurcation curve is strictly decreasing (which implies the uniqueness of nonnegative solutions),
- iii) the curve is not  $C^1$  for a suitable value  $\lambda = \lambda^* > \lambda_1(R)$  corresponding to a "flat solution" (i.e. the solution U is such that  $U'(\pm R) = 0$  and U(s) > 0).

To show the qualitative behaviour of solutions of problem (67) we make the change of variables

$$u_{\lambda,V_0}(x) = \left(\frac{V_0}{\lambda}\right)^{\frac{2}{3}} u(\sqrt{\lambda}x),$$

where u is now the solution of the renormalized problem

$$P(L) \begin{cases} -u'' = f(u) & \text{in } (-L, L), \\ u(\pm L) = 0, \end{cases}$$
 (68)

with

$$f(u) = u - \frac{1}{\sqrt{u}}$$

and  $L = \sqrt{\lambda}R$ . By multiplying by u and integrating by parts, we get that nontrivial solutions may exist only if  $\lambda > \lambda_1 = \frac{\pi^2}{4R^2}$ , the first eigenvalue to the linear problem

$$\begin{cases} -u'' = \lambda u & \text{in } (-R, R), \\ u(\pm R) = 0. \end{cases}$$
 (69)

Notice that now the role of the "eigenvalue"  $\lambda$  is transferred to the length of the new interval  $L = \sqrt{\lambda}R$ . We introduce

$$F(r) = \int_0^r f(s)ds = \frac{r^2}{2} - 2\sqrt{r},$$

and note that f(s) < 0 if  $0 < s < 1 := r_f$  and f(s) > 0 if 1 < s. On the other hand F(s) < 0 if  $0 < s < r_F = 2\sqrt[3]{2}$  ( $\approx 2,51$ ) and F(s) > 0 for  $s > r_F$  (see Figure 5).




Figure 5: Functions f and F.

By multiplying by u', integrating by parts and denoting  $\mu := ||u||_{L^{\infty}}$  for  $\mu \in [r_F, \infty)$  we get that a function u is a positive solution of problem P(L) if and only if

$$\frac{1}{\sqrt{2}} \int_{u(x)}^{\mu} \frac{dr}{(F(\mu) - F(r))^{1/2}} = |x| \,, \ \, \text{for} \ \, |x| \leq L,$$

and  $\mu$  and L > 0 are related by the equation

$$\gamma(\mu) = L$$
,

where  $\gamma: [r_F, +\infty) \to \mathbb{R}$  is given by

$$\gamma(\mu) := \frac{1}{\sqrt{2}} \int_0^\mu \frac{dr}{(F(\mu) - F(r))^{1/2}}.$$
 (70)

Now we use the following fact, whose proof is exactly as in [23] and [27]: a function u is a positive solution of problem P(L) if and only if

$$\frac{1}{\sqrt{2}} \int_{u(x)}^{\mu} \frac{dr}{(F(\mu) - F(r))^{1/2}} = |x|, \text{ for } |x| \le L.$$

Moreover

$$u'(\pm L) = \mp \sqrt{2}\sqrt{F(\mu)}.\tag{71}$$

Thus,  $u'(\pm R) = 0$  corresponds to the case in which the maximum of the solution is  $r_F$ . It can be computed (using the Gauss-Lobatto rules: a remark made by G. Díaz to the author) that  $\gamma(r_F) \approx 2,09$ .

We will start by proving the existence of a branch of positive solutions for a bounded interval of the parameter,  $\lambda \in ((\frac{\pi}{2R})^2, \lambda_1^*)$ .

**Theorem 33** We define with  $F(r) = \frac{r^2}{2} - 2\sqrt{r}$ . Let  $r_F = 2\sqrt[3]{2}$ . Then the mapping  $\gamma: [r_F, +\infty) \to \mathbb{R}$ has the following properties

- (i)  $\gamma \in C[r_F, \infty) \cap C^1(r_F, \infty)$ ;
- (ii) For any  $\mu > r_F \gamma'(\mu) < 0$ ,
- (iii)  $\gamma'(\mu) \to -\xi$  as  $\mu \downarrow r_F$ , for some  $\xi > 0$ ,
- (iv)  $\lim_{\mu \to +\infty} \gamma(\mu) = \frac{\pi}{2}$ .

Concerning the flat solution and the possible solutions with compact support we have:

### Theorem 34 Let

$$\lambda_1^* = \frac{1}{2R^2} \left( \int_0^{r_F} \frac{dr}{(F(\mu) - F(r))^{1/2}} \right)^2 \tag{72}$$

then we have:

a) if  $\lambda \in (0, (\frac{\pi}{2R})^2)$  there is no positive solution, b) if  $\lambda \in ((\frac{\pi}{2R})^2, \lambda_1^*)$  there is a unique positive solution  $u_{\lambda, V_0}$ . Moreover  $\partial u_{\lambda, V_0}/\partial n(\pm R) < 0$  $\|u_{\lambda,V_0}\|_{L^{\infty}(-R,R)} = \left(\frac{V_0}{\lambda}\right)^{\frac{2}{3}} \gamma^{-1}(\sqrt{\lambda}R),$ c) if  $\lambda = \lambda_1^*$  there is only one positive solution  $u_{\lambda_1^*,V_0}$ . Moreover  $u'_{\lambda_1^*,V_0}(\pm R) = 0$ 

$$\|u_{\lambda_1^*,V_0}\|_{L^{\infty}(-R,R)} = \left(\frac{4V_0}{\lambda_1^*}\right)^{\frac{2}{3}}.$$

d) if  $\lambda > \lambda_1^*$ , there is a family of nonnegative solutions that are generated by extending by zero the function  $u_{\lambda_1^*,V_0}$  outside (-R,R) (and which we label again as  $u_{\lambda_1^*,V_0}$ ). In particular, if  $\lambda=\lambda_1^*\omega$  with  $\omega > 1$  we have a family  $S_1(\lambda)$  of compact support nonnegative solutions with connected support defined by

$$u_{\lambda,V_0}(x) = \frac{1}{\omega^{\frac{2}{3}}} u_{\lambda_1^*,V_0}(\sqrt{\omega}x - z),$$

where the shifting argument z is arbitrary among the points  $z \in (-R, R)$  such that support of  $u_{\lambda, V_0}(.) \subset$ (-R,R). Moreover, for  $\lambda > \lambda_1^*$  large enough we can build, similarly, a subset of  $S_i(\lambda)$  of compact support nonnegative solutions with the support formed by j components, with  $j \in \{1, 2, ..., N\}$ , for some suitable  $N = N(\lambda)$  and then the set of nontrivial and nonnegative solutions of  $P(\lambda)$  is formed by  $S(\lambda) = \bigcup_{i=1}^{N} S_i(\lambda)$ . In any case those solutions satisfy that

$$\|u_{\lambda,V_0}\|_{L^{\infty}(-R,R)} = \frac{1}{\omega^{\frac{2}{3}}} \|u_{\lambda_1^*,V_0}\|_{L^{\infty}(-R,R)} = \frac{1}{\omega^{\frac{2}{3}}} \left(\frac{4V_0}{\lambda_1^*}\right)^{\frac{2}{3}}, \text{ for any } \omega = \lambda/\lambda_1^* > 1.$$

*Proofs of Theorems 33 and 34.* The proof of property i) is exactly the same as the one presented in [27]. For the proof of (ii) and (iii), we have

$$\gamma'(\mu) = \int_0^{\mu} \frac{\theta(\mu) - \theta(r)}{(F(\mu) - F(r))^{1/2}} dr$$

where  $\theta(t) = 2F(t) - tf(t) = -3\sqrt{t}$ , and differentiating we get for, any t > 0

$$\theta'(t) = -\frac{3}{2\sqrt{t}} < 0.$$

Hence  $\gamma'(\mu) < 0$  for any  $\mu > r_F$  (which proves (ii)).

For the proof of (iii) it suffices to see that as  $\mu \downarrow r_F$ , the integrand of  $\gamma'(\mu)$  converges pointwise to  $(-F(r))^{-3/2}$  near r=0 and in our case  $(-F(r))^{-3/2}$  behaves as  $r^{-3/4}$  near r=0 and thus  $\gamma'(\mu)$ converges to a number  $-\xi$  as  $\mu \downarrow r_F$ , for some  $\xi > 0$ .

Finally, to prove (iv), we note that

$$\gamma(\mu) \le \frac{\mu}{2} \int_0^1 \frac{dt}{(\frac{\mu^2}{2}(1-t^2))^{1/2}} = \int_0^1 \frac{dt}{\sqrt{1-t^2}} = \frac{\pi}{2}.$$
 (73)

Moreover, we have

$$\gamma(\mu) = \frac{\mu}{\sqrt{2}} \int_0^1 \frac{dt}{\left(\frac{\mu}{\sqrt{2}}((1-t^2) - \frac{1}{\mu^{3/2}}(1-\sqrt{t})\right)},\tag{74}$$

and if  $\mu \to +\infty$ , by using Lebesgue's Theorem, we get

$$\lim_{\mu \to +\infty} \gamma(\mu) = \frac{\pi}{2}.$$

Now we define  $L_0 = \frac{\pi}{2}$  and  $L^*$  given by

$$L^* = \gamma(r_F) = \frac{1}{\sqrt{2}} \int_0^{r_F} \frac{dr}{(-F(r))^{1/2}} = \frac{1}{\sqrt{2}} \int_0^{r_F} \frac{dr}{(2\sqrt{r} - \frac{r^2}{2})^{1/2}}.$$

We know that  $u'(\pm R) = 0$  corresponds to the value  $L^*$  and that the maximum of the solution is  $r_F$ . So, qualitatively, function  $\gamma$  is described by the Figure 6.




Figure 6: Critical value  $L^*$ .

If we now go back with our change of variables, we get

$$\|u_{\lambda,V_0}\|_{L^{\infty}(-R,R)} = \left(\frac{V_0}{\lambda}\right)^{\frac{2}{3}} r_F,$$

and we obtain, finally, the bifurcation diagram given by the first branch of Figure 2, where solutions for  $\lambda > \lambda^*$  are compact supported solutions originated as in [27] from the extension by zero of the free boundary solution  $u_{\lambda^*}$  satisfying

$$\begin{cases} -u_{\lambda^*}'' + \frac{V_0}{\sqrt{u_{\lambda^*}}} = \lambda^* u_{\lambda^*} & \text{in } (-R, R), \\ u_{\lambda^*}(\pm R) = u_{\lambda^*}'(\pm R) = 0. \end{cases}$$
 (75)

The rest of the details are completely analogous to the similar parts of the paper [27].

**Remark 35** Once that we know that for  $\lambda > \lambda_1^*$  we have that

$$u_{\lambda,V_0}(x) = \frac{1}{\omega^{\frac{2}{3}}} u_{\lambda_1^*,V_0}(\sqrt{\omega}x - z),$$

then we get that the bifurcating curve  $\Lambda$  is not  $C^1$  at  $\lambda = \lambda_1^*$  since  $\Lambda'(\lambda_1^*-) = \xi < 0$  and  $\Lambda'(\lambda_1^*+) = -\frac{2C}{3(\lambda_1^*)^{5/3}}$ , with  $C = 4^{2/3}(V_0)^{2/3}$ . In addition, for  $\lambda > \lambda_1^*$  we can express other norms (different from the  $L^{\infty}$ -norm) in terms of  $\lambda$ . For instance, we have that

$$\|u_{\lambda,V_0}'\|_{L^{\infty}(-R,R)} = C\lambda^{-\frac{1}{6}}$$

for a suitable constant C > 0 independent of  $\lambda$ . This proves that  $\left\| u'_{\lambda,V_0} \right\|_{L^{\infty}(-R,R)} \to 0$  as  $\lambda \to +\infty$ .

### 4.2.2 On the construction of the subsolution over $\Omega_{-}$ : continuation

Proof of Theorem 27. Now, let us come back to the construction of the positive subsolution over  $\Omega_{-}$  mentioned in Proposition 31. We recall that, for functions of the form (60) the partial differential equation (61) leads to the ordinary differential equation

$$-U''(\theta) + \frac{V(\theta)}{\sqrt{U(\theta)}} = \lambda U(\theta) \quad \theta \in (\frac{\pi}{2}, \pi), \tag{76}$$

once we assume the constraint

$$\alpha + 2\beta = 4/3,\tag{77}$$

with

$$V(\theta) = \frac{A}{(-\cos\theta)^{\beta}} \text{ and } \lambda = \alpha^2.$$
 (78)

Notice that the potential  $V(\theta)$  is singular only for  $\theta = \frac{\pi}{2}$ . On the other hand, we need to match this subsolution with another function which is positive for  $\theta = \frac{\pi}{2}$ . Then, given  $\varepsilon > 0$  small enough, we will construct the subsolution in two different pieces

$$U(\theta) = \begin{cases} v_1(\theta) & \text{if } \theta \in (\frac{\pi}{2}, \pi - R_0), \\ v_2(\theta) & \text{if } \theta \in (\pi - R_0, \pi), \end{cases}$$

for some  $R_0$  with  $\pi - R_0 \in (\frac{\pi}{2} + \varepsilon, \frac{\pi}{2} + 2\varepsilon)$ , and with  $v_2(\theta)$  such that

$$\begin{cases}
-v_2''(\theta) + \frac{V_{\varepsilon}}{\sqrt{v_2(\theta)}} = \lambda v_2(\theta) & \theta \in (\pi - R_0, \pi), \\
v_2(\pi) = v_2'(\pi) = 0,
\end{cases}$$
(79)

where

$$V(\theta) \le V_{\varepsilon} \text{ if } \theta \in (\frac{\pi}{2} + \varepsilon, \pi).$$

See Figure 7.

Proof of Theorem 27. We recall the change of variable

$$u_{\lambda,V_0}(x) = \left(\frac{V_0}{\lambda}\right)^{\frac{2}{3}} u(\sqrt{\lambda}x),$$

which links the question of the distinguished eigenvalue  $\lambda^* = (\alpha^*)^2 = \frac{1}{2R^2} \gamma(r_F)^2$  with a special length  $L^*$  for which the solution is flat on the boundary.




Figure 7: Singular potential function  $V(\theta)$ .

The computations  $r_F = 4^{2/3}$  ( $\approx 2,51$ ) and  $\gamma(r_F) \approx 2,09$  allow to see that the corresponding  $L^*$  leads to  $\lambda^* = \frac{16}{9}$  and thus  $\alpha^* = \frac{4}{3}$  and  $R = \frac{\pi}{2}$ . This corresponds to the case  $\beta = 0$  thanks to the reciprocal relation

$$\beta = 2 - \frac{3\alpha}{2}.$$

Moreover, we must take A small enough (in fact  $A = \frac{4}{9}$ ) and there is a kind of fortunate coincidence.

Notice that although  $||u_{\lambda,V_0}||_{L^{\infty}} > 1$  there is no difficulty with this since we can take  $u_{0,-}(x,y) = \phi(r,\theta) = kr^{\alpha}U(\theta)$ , for some k > 0 small enough.

If we consider the case of  $\beta \in (0, \frac{1}{2})$  (i.e.  $\alpha^* \in (1, \frac{4}{3})$ ) then the problem is not autonomous (it appears  $V(\theta) = \frac{A}{(-\cos\theta)^{\beta}}$ ), the corresponding R > 0, given by the equation  $\lambda^* = (\alpha^*)^2 = \frac{1}{2R^2}\gamma(r_F)^2$ , is such that  $R > \frac{\pi}{2}$  and we must truncate  $V(\theta)$  on an interval  $(\pi - R_0, \pi)$ , for instance by taking  $R_0 < R$  such that  $u_{\lambda,V_0}(\pi - R_0) = 1$  (see Figures 7 and 8).

We see then that the matching between  $v_1(\theta)$  and  $v_2(\theta)$  must take place at  $\pi - R_0$  (see Figure 8).

The construction of  $v_1(\theta)$ , in the so-called "boundary layer zone", can be carried out as in the onedimensional problem with a distributed potential  $V(\theta)$ . Indeed, given  $h \in (0,1)$  we take  $v_1(\theta)$  as the solution of

$$\begin{cases}
-v_1''(\theta) + \frac{V(\theta)}{\sqrt{v_1(\theta)}} \le 0 & \theta \in (\frac{\pi}{2}, \pi - R_0), \\
v_1(\frac{\pi}{2}) = h, v_1(\pi - R_0) = K \in (h, 1].
\end{cases}$$
(80)

Notice that using that

$$\cos \theta \le \frac{2}{\pi} (\frac{\pi}{2} - \theta), \text{ if } \theta \in (\frac{\pi}{2}, \pi),$$




Figure 8: Matching functions  $v_1(\theta)$  and  $v_2(\theta)$  to get a subsolution.

then the structural assumption on  $V(\theta)$  implies that

$$V(\theta) \le \frac{A\pi^{\beta}}{2^{\beta}(\theta - \frac{\pi}{2})^{\beta}} := \frac{C}{(\theta - \frac{\pi}{2})^{\beta}}.$$

and we can take

$$-v_1''(\theta) + \frac{\frac{C}{(\theta - \frac{\pi}{2})^{\beta}}}{\sqrt{v_1(\theta)}} \le 0$$

with the above boundary conditions. Notice that this is an important key point in the proof, since in Corollary 19, the possible singular coefficient was depending on the y variable but the coefficient j in the two-dimensional problem depends of the other variable j=j(x). By applying Corollary 19 to  $q=-\beta\in(-\frac{1}{2},0)$ , we can define

$$v_1(\theta) = u_q(\theta - \frac{\pi}{2}) + h$$

and thus,  $v_1(\theta) \ge u_q(y)$ , if  $y = \theta - \frac{\pi}{2}$ ,

$$-v_1''(\theta) + \frac{V(\theta)}{\sqrt{v_1(\theta)}} \le -v_1''(\theta) + \frac{\frac{C}{(\theta - \frac{\pi}{2})^{\beta}}}{\sqrt{v_1(\theta)}} \le -u_q''(y) + \frac{Cy^q}{\sqrt{u_q}} = 0$$

Thus, we have

$$v_1'(\frac{\pi}{2}) = 0.$$

We get that the graph of the matched function presents a cusp at  $\theta = \pi - R_0$ , which is far from being *natural*, but it is sufficient for our purposes. Indeed, we can regularize the peaked function by means of a third transition function  $w(\theta)$  if  $\theta \in (\frac{\pi}{2}, \frac{\pi}{2} + 2\varepsilon)$ , so that the resulting subsolution will be of the form

$$U(\theta) = \begin{cases} v_1(\theta) & \text{if } \theta \in (\frac{\pi}{2}, \frac{\pi}{2} + \varepsilon], \\ w(\theta) & \text{if } \theta \in (\frac{\pi}{2} + \varepsilon, \frac{\pi}{2} + 2\varepsilon), \\ v_2(\theta) & \text{if } \theta \in [\frac{\pi}{2} + 2\varepsilon, \pi). \end{cases}$$
(81)

We start by recalling that the change of variables made in the proof of Theorem 32 shows that the parameter  $\lambda^* = (\alpha^*)^2 = \frac{1}{2R^2} \gamma(r_F)^2$  arising in the problem definition of  $v_2(\theta)$  is independent of the coefficient  $V_{\varepsilon}$  also arising in the nonlinear term of this problem. Thus, we will take  $\varepsilon > 0$  such that

$$V(\theta) \le V_{\varepsilon} < \lambda^*, \text{ if } \theta \in (\frac{\pi}{2} + \varepsilon, \pi).$$

We also recall that, from the above construction, we assume  $\beta > 0$  small enough (which implies that also  $R - \frac{\pi}{2}$  is small enough) and then there exists  $\delta > 0$  such that, if  $\theta \in (\frac{\pi}{2} + \varepsilon, \frac{\pi}{2} + 2\varepsilon)$ ,

$$\max(1 - v_1(\theta), 1 - v_2(\theta)) \leq \delta, \text{ and}$$
$$\max(v_1'(\theta), -v_2'(\theta)) \leq \delta.$$

We take  $R_0 = \frac{\pi}{2} + \frac{3\varepsilon}{2}$ . Then, we define the concave function

$$w(\theta) = k_0 + k_1(\theta - R_0) - k_2(\theta - R_0)^2$$
, if  $\theta \in (\frac{\pi}{2} + \varepsilon, \frac{\pi}{2} + 2\varepsilon) = (R_0 - \frac{\varepsilon}{2}, R_0 + \frac{\varepsilon}{2})$ ,

for some constants  $k_0, k_1, k_2$  to be chosen now, with  $k_0, k_2 > 0$ . On the interval  $(\frac{\pi}{2} + \varepsilon, \frac{\pi}{2} + 2\varepsilon)$  we have

$$w(\theta) \ge k_0 - |k_1| \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4}.$$

Then, the subsolution condition, concerning the differential equation, leads to the inequality

$$-w''(\theta) + \frac{V_{\varepsilon}}{\sqrt{w(\theta)}} - \lambda^* w(\theta) \le 2k_2 + \frac{V_{\varepsilon}}{\sqrt{k_0 - |k_1| \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4}}} - \lambda^* (k_0 - |k_1| \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4}) \le 0.$$
 (82)

On the boundary, we must demand the continuity of the matching

$$\begin{cases} w(\frac{\pi}{2} + \varepsilon) = k_0 - k_1 \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4} = v_1(\frac{\pi}{2} + \varepsilon), \\ w(\frac{\pi}{2} + 2\varepsilon) = k_0 + k_1 \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4} = v_2(\frac{\pi}{2} + 2\varepsilon), \end{cases}$$
(83)

and the good sign conditions on the measures generated by the derivatives matching

$$\begin{cases} w'(\frac{\pi}{2} + \varepsilon) = k_1 + 2k_2 \frac{\varepsilon}{2} \ge v'_1(\frac{\pi}{2} + \varepsilon), \\ w'(\frac{\pi}{2} + 2\varepsilon) = k_1 - 2k_2 \frac{\varepsilon}{2} \le v'_2(\frac{\pi}{2} + 2\varepsilon). \end{cases}$$
(84)

Let

$$v_1^L := v_1(\frac{\pi}{2} + \varepsilon), \qquad v_2^R := v_2(\frac{\pi}{2} + 2\varepsilon).$$

From (83) we have

$$k_0 - k_1 \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4} = v_1^L, \ k_0 + k_1 \frac{\varepsilon}{2} - \frac{k_2 \varepsilon^2}{4} = v_2^R.$$

Adding and subtracting these equations gives

$$k_1 = \frac{v_2^R - v_1^L}{\varepsilon}, \qquad k_0 = \frac{v_1^L + v_2^R}{2} + \frac{k_2 \varepsilon^2}{4}.$$

Hence  $k_1$  is fixed by the endpoint difference and  $k_0$  depends linearly on  $k_2 > 0$ . From (84) we have

$$k_1 + k_2 \varepsilon \ge v_1'(L), k_1 - k_2 \varepsilon \le v_2'(R).$$

Rearranging gives the lower bounds

$$k_2 \ge \frac{v_1'(L) - k_1}{\varepsilon}, \qquad k_2 \ge \frac{k_1 - v_2'(R)}{\varepsilon}.$$

Thus any admissible  $k_2$  must satisfy

$$k_2 \ge k_{2,\min} := \max \left\{ 0, \frac{v_1'(L) - k_1}{\varepsilon}, \frac{k_1 - v_2'(R)}{\varepsilon} \right\}.$$

Using the hypotheses  $|v_1'(L)|, |v_2'(R)| \leq \delta$  and  $|v_2^R - v_1^L| \leq \delta$ , we have  $|k_1| \leq \delta/\varepsilon$ , so  $k_{2,\min} = O(\delta/\varepsilon)$ , typically small. By substituting the relations for  $k_0, k_1$  into (??) we can define:

$$\Phi(k_2) := 2k_2 + \frac{V_{\varepsilon}}{\sqrt{k_0 - |k_1|(\varepsilon/2) - \frac{k_2 \varepsilon^2}{4}}} - \lambda^* \left(k_0 - |k_1|(\varepsilon/2) - \frac{k_2 \varepsilon^2}{4}\right).$$

Note that

$$k_0 - |k_1|(\varepsilon/2) - \frac{k_2 \varepsilon^2}{4} = \frac{v_1^L + v_2^R}{2} - |k_1| \frac{\varepsilon}{2},$$

which is independent of  $k_2$ . Hence, the desired inequality becomes

$$\Phi(k_2) = 2k_2 + C \le 0,$$

where

$$C := \frac{V_{\varepsilon}}{\sqrt{\frac{v_1^L + v_2^R}{2} - |k_1|(\varepsilon/2)}} - \lambda^* \left(\frac{v_1^L + v_2^R}{2} - |k_1|(\varepsilon/2)\right).$$

Using  $\max(1-v_1,1-v_2) \leq \delta$ , we have  $v_1^L, v_2^R \geq 1-\delta$  and thus

$$\frac{v_1^L + v_2^R}{2} - |k_1| \frac{\varepsilon}{2} \ge 1 - \frac{3\delta}{2} > 0$$

for small  $\delta$ . Therefore

$$-C \approx V_{\varepsilon} + \lambda^* < 0.$$

By continuity, C remains positive for sufficiently small  $\delta$ . Because C < 0, the function  $\Phi(k_2) = 2k_2 + C$  is increasing and linear in  $k_2$ . Thus the inequality  $\Phi(k_2) \leq 0$  holds for any

$$0 < k_2 \le \frac{-C}{2}.$$

To satisfy both the derivative and main inequalities, it suffices to have

$$k_{2,\min} \le k_2 \le \frac{-C}{2}.$$

Since  $k_{2,\min} = O(\delta/\varepsilon)$  and  $-C/2 \approx (\lambda^* - V_{\varepsilon})/2 > 0$ , this interval is nonempty for small enough  $\delta$ . In conclusion, defining

$$B := \frac{v_1^L + v_2^R}{2} - |k_1| \frac{\varepsilon}{2}, \qquad C := \frac{V_{\varepsilon}}{\sqrt{B}} - \lambda^* B,$$

a convenient, explicit choice is

$$\begin{cases} k_1 = \frac{v_2^R - v_1^L}{\varepsilon}, & k_{2,\min} = \max \left\{ 0, \frac{v_1'(L) - k_1}{\varepsilon}, \frac{k_1 - v_2'(R)}{\varepsilon} \right\}, \\ k_2 = k_{2,\min} + \min \left\{ \eta, \frac{-C}{4} \right\}, & k_0 = \frac{v_1^L + v_2^R}{2} + \frac{k_2 \varepsilon^2}{4}, \end{cases}$$

where  $\eta > 0$  is any small tolerance (e.g.,  $\eta = \frac{1}{10}\varepsilon^3$ ). Then,  $k_0, k_2 > 0$ , all the required inequalities are satisfied and the function  $U(\theta)$  defined in (81) is a subsolution as required,  $0 < U(\theta) \le 1$  for  $\theta \in (\frac{\pi}{2}, \pi)$ . Finally, the boundary inequality

$$u_{0,-}(x,y) = kr^{\alpha}U(\theta) \le 1 \text{ if } y = 1$$

holds for k small enough such that

$$kH^{\alpha}U(\theta) \le 1$$
, where  $H = \sqrt{a^2 + 1}$ . (85)

and the proof of the existence of  $u_{0,-}(x,y)$ , a  $p_0$ -subsolution of problem  $P_{a,0,j}$ , assumed in Proposition 31 is now completed.

With respect to the nondegenerate estimate (45) of Theorem 27, we point out that the estimate from above on  $\Omega_+$  is trivially satisfied since  $u_{0,-}(x,y)$  is a harmonic function). Thus, the important part is to show the lower estimate for points  $(x,y) \in \Omega_-$  which are very near the part of the boundary in which y = 0. For those points, their distance to the boundary is  $\delta(x,y) = y$ . Since r > y, from the constructed subsolution,  $u_{0,-}(x,y) = kr^{\alpha}U(\theta)$ , and the fact that we have

$$c_1(\pi - \theta)^{4/3} \le U(\theta) \le c_2(\pi - \theta)^{4/3},$$

for  $\theta$  sufficiently close to  $\pi$ , with positive constants  $c_1, c_2$  (as we can easely prove via the comparison with sub/supersolutions of the form  $\varphi_{\pm}(x) := A_{\pm}x^{4/3}$ ,  $x = \pi - \theta$ ), we get the nondegenerate estimate (45) of Theorem 27. Notice that the assumptions of the uniqueness result Theorem 28 are satisfied.

### 4.3 Construction of a partially flat supersolution

We make now the structural assumptions, (7), stated in Theorem 1. Thus, in particular, we already know, by Theorems 27 and 28 that the mere existence of a flat nondegenerate subsolution implies the uniqueness of the solution. We concentrate our attention now in proving that the above conditions on j(x) are sufficient conditions for the existence of a partially flat supersolution, i.e., being flat only on the half of the cathode region under study  $[-a, 0] \times \{0\}$ .

Proof of Theorem 29. Since we only need to work with the inequality  $\geq$  in the equation, it is enough to use (as before) that

$$j(r,\theta) = \frac{A}{(-r\cos\theta)^{\beta}} \ge V_0 \text{ for } x \in (-a,0).$$

We will search for the supersolution by some matching arguments, now in three different regions, as indicated in the Figure 9.




Figure 9: Decomposition in subdomains.

In the first region  $\Omega_{-}$  we will construct the supersolution  $u_{-}^{0}(x,y)$  in the same form than the subsolution,

$$u_{-}^{0}(x,y) = \overline{\phi}(r,\theta) = \overline{k}r^{\overline{\alpha}}\overline{U}(\theta)$$
, for some  $\overline{k} > 0$ ,  $\overline{\alpha} > 1$ ,

and thus with  $U(\theta)$  solution of

$$-\overline{U}''(\theta) + \frac{V_0}{\sqrt{(\overline{U}(\theta)}} = \lambda^* \overline{U}(\theta) \quad \theta \in (\frac{\pi}{2}, \pi).$$
 (86)

So that no boundary layer needs to be taken into account now. This means that, in terms of the proof of Theorem 27,  $\overline{U}(\theta)$  is very similar to  $v_2(\theta)$ . Nevertheless, since we should ensure that  $u_-^0(x,y) \ge u_{0,-}(x,y)$  we will take the parameter  $\lambda^* = (\overline{\alpha})^2 = \frac{1}{2\overline{R}^2} \gamma(r_F)^2$ , such that  $\overline{\alpha} \le \alpha$  (the exponent used in the definition of the subsolution), which is obtained if we take  $\frac{\pi}{2} \leq \overline{R} \leq R$  (with R the value choosen for the part of the subsolution given by  $v_2(\theta)$ ). In this way, we will also have that

$$1 - \delta \le \overline{U}(\frac{\pi}{2}) \le 1$$
, and  $-\delta \le \overline{U}'(\frac{\pi}{2}) \le 0$ ,

for some  $\delta > 0$  small enough. Note that even if  $V_0 \leq V_{\varepsilon}$ , the solution structure of these semilinear problems allows us to take k > k large enough (with k the value taken for the subsolution) in order to have  $u_{-}^{0}(x,y) \geq u_{0,-}(x,y)$  on  $\Omega_{-}$ .

The matching with the second region will follow a different argument. In fact, on  $\Omega_+$ , the supersolution  $u_+^0$  must be superharmonic. This will ensure that  $u_+^0(x,y) \ge u_{0,+}(x,y)$  also on  $\Omega_+ = \Omega_+^2 \cup \Omega_+^3$  (see Figure 9), since  $u_{0,+}$  was taken as subharmonic and the inequality  $u_+^0 \ge u_{0,+}$  is realized on the boundary of  $\Omega_+$ . We start by searching  $u_+^0(x,y)$  on  $\Omega_+^2$  in the form

$$u_{+,2}^0(x,y) = Kr^{\alpha}\sin(\alpha\theta), \ \theta \in (\theta_b, \frac{\pi}{2}),$$

with  $\tan \theta_b = b$ . It is not difficult to check that

$$-\Delta u_{+,2}^0 = K\alpha(\alpha - 1)r^{\alpha - 2}\sin(\alpha\theta) \ge 0.$$

The matching with  $u_{-}^{0}(x,y)$  is a delicate question. For a  $H^{1}$ -matching we must have:

$$\left\{ \begin{array}{ll} u_{-}^{0}(0,y) = u_{+,2}^{0}(0,y) & y \in (0,1), \\ \nabla u_{-}^{0}(0,y) = \nabla u_{+,2}^{0}(0,y) & y \in (0,1). \end{array} \right.$$

The first condition holds if

$$\frac{k}{K} = \frac{\sin(\frac{\alpha\pi}{2})}{\overline{U}(\frac{\pi}{2})}.$$

The second condition (once we allow the formation of singularities with a good sign, i.e., with a nonnegative generated implicit distribution) leads to

$$\overline{U}'(\frac{\pi}{2}) \ge \frac{K}{k}\alpha\cos(\frac{\alpha\pi}{2}) = \alpha\overline{U}(\frac{\pi}{2})\frac{\cos(\frac{\alpha\pi}{2})}{\sin(\frac{\alpha\pi}{2})} = \alpha\overline{U}(\frac{\pi}{2})\cot(\frac{\alpha\pi}{2}).$$

But,  $\frac{\alpha\pi}{2} \in (\frac{\pi}{2}, \frac{2\pi}{3}) \subset (0, \pi)$  so that  $\cot(\frac{\alpha\pi}{2}) < 0$  (see the Figure 10). Thus we must investigate for which  $\alpha \in (1, \frac{4}{3}]$  we have some kind of Robin type boundary inequality on  $\overline{U}(\theta)$  for  $\theta = \frac{\pi}{2}$ :

$$\frac{-\overline{U}'(\frac{\pi}{2})}{\overline{U}(\frac{\pi}{2})} \le \alpha(-\cot(\frac{\alpha\pi}{2})). \tag{87}$$

This condition trivially holds if  $\alpha = \frac{4}{3}$  (since, in that case,  $\overline{U}'(\frac{\pi}{2}) = 0$ ). Then by continuity in  $\alpha$ , there exists  $\alpha_0 \in (1, \frac{4}{3})$  such that (87) holds for any  $\alpha \in (1, \frac{4}{3})$ . This justifies the assumption (7) in Theorem 29 (and in Theorem 1).




Figure 10: Representation of the functions  $-\alpha \cot(\frac{\alpha\pi}{2})$ 

Finally, in the third region  $\Omega^3_+$ , we extend the above function  $u^0_{+,2}(x,y)$ , by a kind of interpolation argument (remember that we must have the boundary condition  $u^0_{+,3}(b,y) \geq y$ , for any  $y \in (0,1)$ ). Then we define

$$u_{+,3}^0(x,y) = C_3 r \sin(\alpha \theta) + \widehat{C}_3 r^{\alpha} \sin(\alpha \theta_b) \quad \text{if } (x,y) \in \Omega_{+,3},$$

with  $\tan \theta_b = b$ . It is easy to prove that  $u_{+,3}^0(x,y)$  is a super-harmonic function  $(-\Delta u_{+,3}^0 \ge 0)$  and that (if b is large enough) the positive constants  $C_3$  and  $\widehat{C}_3$  can be chosen such that  $u_{+,3}^0(x,y)$  satisfies the correct inequalities on the boundaries

$$\begin{cases} u_{+,3}^{0}(b,y) \ge y, & \text{for any } y \in (0,1), \\ u_{+,3}^{0}(x,0) \ge 0 & \text{for any } x \in (0,b), \end{cases}$$

and it correctly matches with  $u_{+,2}^0(x,y)$  (generalizing, at most, a "good signed" measure) on the matching boundary  $\theta = \theta_b$ . This proves Theorem 29 (and also Theorem 1, except the proof of the uniqueness of solution which will be presented in the next subsection).

### 4.4 Proof of the uniqueness of nondegenerate solutions

The proof of Theorem 28 will be a consequence of some previous results concerning the parabolic problem

$$PP_{a,b,j,u_0} = \begin{cases} u_t - \Delta u + \frac{j(x)}{\sqrt{u}} \chi_{\{u>0\}} = 0 & t > 0, \ x \in (-a,b), \ y \in (0,1), \\ u(t,x,0) = 0 & t > 0, \ x \in (-a,b), \\ u(t,x,1) = 1 & t > 0, \ x \in (-a,b), \\ u(t,-a,y) = y^{4/3} & t > 0, \ y \in (0,1), \\ u(t,b,y) = y & t > 0, \ y \in (0,1), \\ u(0,x,y) = u_0(x,y) & x \in (-a,b), \ y \in (0,1). \end{cases}$$

$$(88)$$

Notice that the above parabolic problem is not related to the evolution problem associated with the electron beam formulation. It would be related to some associated evolution problem if the modelling under study were concerned with the Chemical Engineering problem dealing with a chemical kinetics with a reaction of order -1/2. Here, we are only interested in the fact that this artificial parabolic problem leads us (under suitable assumptions) to the uniqueness of solutions of the mathematical model for a stationary electron beam.

The existence of solutions to problem  $PP_{a,b,j,u_0}$  is an easy variation of results dealing with parabolic quenching type problems (see, e.g., [16], [15], [21] and [14]). We point out that no flat condition is required in this formulation. As in the elliptic problem, we can prove that for certain initial data, the solutions are non-degenerate in the sense that for any t > 0 the solution u is such that u(t) belongs to the following set

$$\mathcal{M}(\nu) := \left\{ u \in L^2(\Omega; \delta) \mid \text{such that } u(x,y) \geq C \delta(x,y)^{\nu} \quad \text{in } \Omega, \text{ for some } C > 0 \right\}, \ \nu \in \left(0, \frac{4}{3}\right].$$

The following result is an adaptation to this framework of the main theorem of [21]. It gives the continuous dependence of solutions for the initial data (implying, obviously, the uniqueness of solutions), as well as a smoothing effect concerning the initial datum. In the proof, we will use some Hilbertian techniques. We consider two initial data, and we will prove that if the respective solutions are such that  $u(t), v(t) \in \mathcal{M}(\nu)$  (i.e. with  $\delta^{-\nu}u, \delta^{-\nu}v \geq C$ ) then we can estimate the  $L^2(\Omega)$ -norm of  $\delta^{-\gamma}[u(t) - v(t)]_+$  for some suitable  $\gamma \in (0, 1]$  in terms of the  $L^2(\Omega)$ -norm of  $[u_0 - v_0]_+$ . Notice that this automatically implies an estimate on the  $L^2(\Omega)$ -norm of  $[u(t) - v(t)]_+$ .

**Theorem 36** Let  $u_0, v_0 \in L^1(\Omega) \cap L^2(\Omega; \delta)$ . Let u, v be weak solutions of  $PP_{a,b,j,u_0}$  and  $PP_{a,b,j,v_0}$ , respectively such that  $u(t), v(t) \in \mathcal{M}(\nu)$  for some  $\nu \in (0, \frac{4}{3}]$ . Then, for any  $t \in (0, \infty)$ , we have

$$\left\| \delta^{-\gamma} [u(t) - v(t)]_{+} \right\|_{L^{2}(\Omega)} \le C t^{-\frac{2\gamma + 1}{4}} \left\| [u_{0} - v_{0}]_{+} \right\|_{L^{2}(\Omega;\delta)}, \tag{89}$$

with

$$\gamma := \min\left\{\frac{3\nu}{2}, 1\right\} \tag{90}$$

and for some constant C>0 independent of t. In particular,  $u_0\leq v_0$  implies that for any  $t\in[0,+\infty)$ ,

$$u(t,\cdot) \leq v(t,\cdot)$$
 a.e. in  $\Omega$ 

and

$$\left\| \delta^{-\gamma} \left( u(t) - v(t) \right) \right\|_{L^{2}(\Omega)} \le C t^{-\frac{2\gamma + 1}{4}} \left\| u_0 - v_0 \right\|_{L^{2}(\Omega; \delta)}. \tag{91}$$

Before referring to the proof of Theorem 36, we will prove that this implies the uniqueness of the positive solution for the stationary problem  $P_{a,b,j}$  presented in Theorem 28.

Proof of Theorem 28. Let us call  $u_{\infty}$  and  $v_{\infty}$  two possible solutions of  $P_{a,b,j}$  in the class  $\mathcal{M}(\nu)$ . By taking  $u_0 = u_{\infty}$  and  $v_0 = v_{\infty}$  as initial data in  $PP_{a,b,j,u_0}$  and  $PP_{a,b,j,v_0}$ , since  $u_{\infty}$  and  $v_{\infty}$  are, obviously, solutions of the respective parabolic problems, we get that  $u_{\infty} - v_{\infty}$  satisfies

$$\left\| \delta^{-1} \left( u_{\infty} - v_{\infty} \right)_{+} \right\|_{L^{2}(\Omega)} \le C t^{-\frac{2\gamma+1}{4}} \| (u_{\infty} - v_{\infty})_{+} \|_{L^{2}(\Omega, \delta)}.$$

Making  $t \nearrow +\infty$  and reversing the role of  $u_{\infty}$  and  $v_{\infty}$ , we get that  $u_{\infty} = v_{\infty}$ .

The proof of Theorem 36 follows some slight modifications of the paper [21]. Nonetheless, for the sake of completeness, we present here the main lines of the proof.

Without loss of generality, we can use the notion of mild solution on  $L^1(\Omega)$ , i.e.  $u \in \mathcal{C}([0,T); L^1(\Omega))$ , for any T > 0:  $j(x)u^{-1/2} \in L^1(\Omega \times (0,T))$  and u fulfills the identity

$$u(\cdot,t) = S(t)u_0(\cdot) - \int_0^t S(t-s)j(\cdot)(\chi_{\{u>0\}}u^{-1/2}(\cdot,s))ds, \quad \text{in } L^1(\Omega),$$
(92)

where S(t) is the  $L^1(\Omega)$ -semigroup corresponding to the Laplace operator with the corresponding Dirichlet (stationary) boundary conditions (see, e.g., [14]). We shall need some well-known auxiliary results. The first one is a singular version of the Gronwall's inequality (due to Brezis and Cazenave [7]), which is especially useful in the study of non-global Lipschitz perturbations of the heat equation.

**Lemma 37** Let T > 0,  $A \ge 0$ ,  $0 \le a, b \le 1$  and let f be a non-negative function with  $f \in L^p(0,T)$  for some p > 1 such that  $\max\{a,b\} < 1/p'$  (where  $\frac{1}{p} + \frac{1}{p'} = 1$ ). Consider a non-negative function  $\varphi \in L^{\infty}(0,T)$  such that, for almost every  $t \in (0,T)$ ,

$$\varphi(t) \le At^{-a} + \int_0^t (t-s)^{-b} f(s)\varphi(s) \, ds. \tag{93}$$

Then, there exists C > 0 only depending on T, a, b, p and  $||f||_{L^p(0,T)}$  such that, for almost every  $t \in (0,T)$ ,

$$\varphi(t) \le ACt^{-a}. (94)$$

We shall also use some regularizing effects properties satisfied by the semigroup S(t) of the heat equation with zero Dirichlet boundary conditions (since we will apply it to the difference of two solutions of the parabolic problem  $PP_{a,b,j,u_0}$ ) due to different authors, among them [49], [16] and [47]: see details, in [21].

### Lemma 38

1. There exists C > 0 such that, for any t > 0 and any  $u_0 \in L^2(\Omega)$ ,

$$\|\nabla S(t)u_0\|_{L^2(\Omega)} \le Ct^{-\frac{1}{2}} \|u_0\|_{L^2(\Omega)}.$$
(95)

2. There exists C>0 such that, for any t>0 and any  $u_0\in L^1(\Omega)$ ,

$$||S(t)u_0||_{L^2(\Omega)} \le Ct^{-\frac{N}{4}} ||u_0||_{L^1(\Omega)}. \tag{96}$$

3. There exists C > 0 such that, for any t > 0, any  $m \in (0,1]$  and any  $u_0 \in L^2(\Omega; \delta^{2m})$ ,

$$||S(t)u_0||_{L^2(\Omega)} \le Ct^{-\frac{m}{2}} ||u_0||_{L^2(\Omega,\delta^{2m})}. \tag{97}$$

4. There exists C > 0 such that, for any t > 0, any  $p \in [1, +\infty)$  and any  $u_0 \in L^p(\Omega, \delta)$ ,

$$||S(t)u_0||_{L^p(\Omega)} \le Ct^{-\frac{1}{2p}} ||u_0||_{L^p(\Omega,\delta)}.$$
 (98)

Finally, we can give the main arguments of the proof.

*Proof of Theorem 36.* By the constant variations formula, we know that for any  $t \in [0,T]$ ,

$$u(t) - v(t) = S(t)(u_0 - v_0) + \int_0^t S(t - s) \left( h(u(s)) - h(v(s)) \right) ds \quad \text{in } \Omega, \tag{99}$$

where  $h(x,u) := j(x)u^{-1/2}$ . By the convexity of the function  $u \mapsto u^{-1/2}$  and the assumption that  $u(t), v(t) \in \mathcal{M}(\nu)$ , we deduce that

$$h(x,u) - h(x,v) \le Cj(x)\delta^{-3\nu/2}(u-v)_{+} \quad \text{in } \Omega.$$
 (100)

Thus, if we denote w := u - v, we get for any  $\tau, t \in [0, T]$  with  $\tau \leq t$ 

$$w_{+}(t) \le S(t-\tau)w_{+}(\tau) + C \int_{\tau}^{t} S(t-s)(j(x)\delta^{-3\nu/2}w_{+}(s)) ds.$$
 (101)

We multiply (101) by the weight  $\delta^{-\gamma}$ , with  $\gamma \in [0,1]$  to be chosen later, and take the  $L^2$ -norms. Then,

$$\|\delta^{-\gamma}w_{+}(t)\|_{L^{2}(\Omega)} \leq \|\delta^{-\gamma}S(t-\tau)w_{+}(\tau)\|_{L^{2}(\Omega)} + C\int_{\tau}^{t} \|S(t-s)j(x)\delta^{-[(\beta+1)\nu+\gamma]}w_{+}(s)\|_{L^{2}(\Omega)} ds.$$

Let us fix s, t > 0 and let us call  $\psi := S(t-s)j(x)\delta^{-(\beta+1)\nu}w_+(s)$ . Then, by Hölder inequality,

$$\|\delta^{-\gamma}\psi\|_{L^2(\Omega)}^2 = \int_{\Omega} \frac{\psi^2}{\delta^{2\gamma}} \, dx \le \left(\int_{\Omega} \frac{\psi^2}{\delta^2} \, dx\right)^{\gamma} \left(\int_{\Omega} \psi^2 \, dx\right)^{1-\gamma}$$

(note that the limit cases  $\gamma \equiv 0$  and  $\gamma \equiv 1$  are allowed). Then, applying the Hardy inequality,

$$\|\delta^{-\gamma}\psi\|_{L^2(\Omega)} \le C\|\nabla\psi\|_{L^2(\Omega)}^{\gamma}\|\psi\|_{L^2(\Omega)}^{1-\gamma}$$

By property 1 of Lemma 38, for  $\frac{t-s}{2}$ , we get

$$\|\delta^{-[(\beta+1)\nu+\gamma]}S(t-s)w_{+}(s)\|_{L^{2}(\Omega)} \le C(t-s)^{-\frac{\gamma}{2}}\|S\left(\frac{t-s}{2}\right)j(x)\delta^{-(\beta+1)\nu}w_{+}(s)\|_{L^{2}(\Omega)}. \tag{102}$$

Analogously, using property 4 of Lemma 38, and that  $j \in L^2(\Omega)$  (remember the assumptions made in Theorem 27)

$$\|\delta^{-\gamma}S(t)w_{+}(0)\|_{L^{2}(\Omega)} \le Ct^{-\frac{\gamma}{2}}\|S\left(\frac{t}{2}\right)j(x)w_{+}(0)\|_{L^{2}(\Omega)} \le Ct^{-\left(\frac{\gamma}{2} + \frac{1}{4}\right)}\|w_{+}(0)\|_{L^{2}(\Omega;\delta)}.$$
(103)

In order to apply the singular Gronwall's inequality, we must relate the weights  $\delta^{-\gamma}$  and  $\delta^{-3\nu/2}$  keeping in mind that  $\gamma \in [0,1]$ . To do that, we apply property 3 of Lemma 38 for some  $m \in [0,1]$ . We shall take

$$3\nu/2 = \gamma + m. \tag{104}$$

Indeed, if  $(\beta + 1)\nu \in (1, 2]$ , then we take  $\gamma = 1$ ,  $m = 3\nu/2 - 1$  and we apply point 3 of Lemma 38 to the initial datum:

$$\|S\left(\frac{t-s}{2}\right)j(x)\delta^{-(\beta+1)\nu}w_{+}(s)\|_{L^{2}(\Omega)} = \|S\left(\frac{t-s}{2}\right)j(x)\delta^{-(m+1)}w_{+}(s)\|_{L^{2}(\Omega)} \le C(t-s)^{-\frac{m}{2}}\|\delta^{-\gamma}w_{+}(s)\|_{L^{2}(\Omega)}.$$
(105)

On the other hand, if  $3\nu/2 \in [0,1]$ , we can take  $\gamma = 3\nu/2$  and thus, since S(t-s) is a contraction in  $L^2(\Omega)$ , we get

$$\|S\left(\frac{t-s}{2}\right)j(x)\delta^{-(\beta+1)\nu}w_{+}(s)\|_{L^{2}(\Omega)} = \|S\left(\frac{t-s}{2}\right)j(x)\delta^{-\gamma}w_{+}(s)\|_{L^{2}(\Omega)} \le \|\delta^{-\gamma}w_{+}(s)\|_{L^{2}(\Omega)}, \quad (106)$$

which corresponds to (104) with m=0. In other words,

$$\gamma = \min\{1, 3\nu/2\}$$

and

$$m = \max\{3\nu/2 - 1, 0\}.$$

Collecting the previous inequalities, we arrive to

$$\|\delta^{-\gamma}w_{+}(t)\|_{L^{2}(\Omega)} \leq Ct^{-\frac{2\gamma+1}{4}}\|w_{+}(0)\|_{L^{2}(\Omega;\delta)} + C\int_{0}^{t} (t-s)^{-\frac{m}{2}}\|\delta^{-\gamma}w_{+}(s)\|_{L^{2}(\Omega)}.$$

Thus, we can apply Lemma 37 with  $a=\frac{2\gamma+1}{4}\in\left[\frac{1}{4},\frac{3}{4}\right],\,b=\frac{m}{2}$  and  $A=C\|w_+(0)\|_{L^2(\Omega;\delta)}$  to deduce that

$$\|\delta^{-\gamma}w_+(t)\|_{L^2(\Omega)} \le Ct^{-\frac{2\gamma+1}{4}}\|w_+(0)\|_{L^2(\Omega;\delta)}.$$

Remark 39 As in the proof of Theorem 28, if  $u_{\infty}$  is the non-degenerate solution of problem  $P_{a,b,j}$ , for any  $u_0 \in L^2(\Omega; \delta)$  leading to a non-degenerate solution u(t) of the parabolic problem  $PP_{a,b,j,u_0}$ , we get a quantitative estimate on the asymptotic stability of  $u_{\infty}$  in the class of non-degenerate stationary solutions, since, for any  $t \in (0, \infty)$ , we have

$$\|\delta^{-\gamma}[u(t) - u_{\infty}]\|_{L^{2}(\Omega)} \le Ct^{-\frac{2\gamma+1}{4}} \|u_{0} - u_{\infty}\|_{L^{2}(\Omega;\delta)},$$

with  $\gamma := \min\left\{\frac{3\nu}{2}, 1\right\}$ , for some constant C > 0 independent of t.

Acknowledgement Research partially supported by the project PID2023-146754NB-I00 funded by MCIU/AEI/10.13039/501100011033 and FEDER, EU. MCIU/AEI/10.13039/-501100011033/FEDER, EU. The author thanks Haïm Brezis and Joel Lebowitz for several conversations on this subject. Thanks are also extended to my brother, Gregorio Díaz, and my friend, Laurent Véron, for their careful reading of an earlier version of the manuscript. They suggested several improvements to the exposition. In particular, L. Véron showed me an alternative proof of Theorem 2.



Figure 11: Haïm Brezis, Joel Lebowitz and the author, in Paris, 2005.

### References

- [1] H. aBrezis and J. I. Díaz eds., Mathematics and Environment, Proceedings of the meeting between the Académie des Sciences and the Real Academia de Ciencias, Paris, 23-24 may, 2002, Rev. R. Acad. Cien.Serie A Matem.(RACSAM) 96, 2002.
- [2] G. K. Batchelor, An introduction to fluid dynamics, Cambridge University Press, 1967.
- [3] A. Bensoussan, H. Brezis, and A. Friedman, Estimates on the free boundary for quasi-variational inequalities, Comm. Partial Differential Equations 2 (1977), no. 3, 297–321.
- [4] H. Berestycki and P.-L. Lions, Some applications of the method of super and subsolutions, Bifurcation and Nonlinear Eigenvalue Problems: Proceedings, Université de Paris XIII, Villetaneuse, France, October 2–4, 1978, Springer, 2006, pp. 16–41.
- [5] H. Brezis, Solutions of variational inequalities with compact support, Uspekhi Mat. Nauk 129 (1974), 103–108.
- [6] \_\_\_\_\_\_, Functional analysis, Sobolev spaces and partial differential equations, Springer. New York, 2011
- [7] H. Brezis and Th. Cazenave, A nonlinear heat equation with singular initial data, Journal D'Analyse Mathematique **68** (1996), no. 1, 277–304.
- [8] H. Brezis, Th. Cazenave, Y. Martel, and A. Ramiandrisoa, Blow up for  $u_t \Delta u = g(u)$  revisited, Adv. Differential Equations 1 (1974), no. 1, 73–90.
- [9] C. J. Budd, A. Friedman, B. McLeod, and A. A. Wheeler, *The space charge problem*, SIAM Journal on Applied Mathematics **50** (1990), no. 1, 181–198.
- [10] X. Chen, K. Dickens, E. H. Choi, and M. Kristiansen, Space-charge limited current for 1-d cylindrical diodes, Digest of Technical Papers. PPC-2003. 14th IEEE International Pulsed Power Conference (IEEE Cat. No. 03CH37472), vol. 1, IEEE, 2003, pp. 467–470.
- [11] C. D. Child, Discharge from hot CaO, Physical Review (Series I) 32 (1911), no. 5, 492.
- [12] Y. S. Choi, A. C. Lazer, and P. J. McKenna, Some remarks on a singular elliptic boundary value problem, Nonlinear Analysis: Theory, Methods & Applications 32 (1998), no. 3, 305–314.
- [13] M. G. Crandall, P. H. Rabinowitz, and L. Tartar, On a Dirichlet problem with a singular nonlinearity, Communications in Partial Differential Equations 2 (1977), no. 2, 193–222.
- [14] A. N. Dao, J. I. Díaz, and P. Sauvy, Quenching phenomenon of singular parabolic problems with L<sup>1</sup> initial data, Electronic Journal of Differential Equations 2016 (2016), no. 2016, 1–16.
- [15] J. Dávila and M. Montenegro, *Positive versus free boundary solutions to a singular elliptic equation*, Journal d'Analyse Mathématique **90** (2003), no. 1, 303–335.
- [16] \_\_\_\_\_\_, Existence and asymptotic behavior for a singular parabolic equation, Trans. Amer. Math. Soc. **357** (2005), no. 5, 1801–1828.
- [17] P. Degond and P.-A. Raviart, An asymptotic analysis of the one-dimensional Vlasov-Poisson system: the Child-Langmuir law, Asymptotic Analysis 4 (1991), no. 3, 187–214.
- [18] J. I. Díaz, Nonlinear partial differential equations and free boundaries, Pitman, London, 1985.
- [19] \_\_\_\_\_\_, On the Haïm Brezis pioneering contributions on the location of free boundaries, Elliptic and Parabolic Problems: A Special Tribute to the Work of Haïm Brezis, Progress in Nonlinear Differential Equations and Their Applications, vol. 63, Birkhäuser Verlag Basel/Switzerland, 2005, pp. 217–234.

- [20] \_\_\_\_\_, On the ambiguous treatment of the Schrödinger equation for the infinite potential well and an alternative via flat solutions: the one-dimensional case, Interfaces and Free Boundaries 17 (2015), no. 3, 333–351.
- [21] J. I. Díaz and J. Giacomoni, Monotone continuous dependence of solutions of singular quenching parabolic problems, Rendiconti del Circolo Matematico di Palermo Series 2 72 (2023), no. 4, 2593– 2602.
- [22] J. I. Díaz and J. Hernández, On the existence of a free boundary for a class of reaction-diffusion systems, SIAM J. Math. Anal. 15 (1984), no. 4, 670–685.
- [23] \_\_\_\_\_\_, Global bifurcation and continua of nonnegative solutions for a quasilinear elliptic problem, Comptes Rendus de l'Académie des Sciences-Series I-Mathematics **329** (1999), no. 7, 587–592.
- [24] \_\_\_\_\_, Positive and nodal solutions bifurcating from the infinity for a semilinear equation: solutions with compact support, Portugaliae Mathematica 72 (2015), no. 2, 145–160.
- [25] \_\_\_\_\_, Beyond the classical strong maximum principle: sign-changing forcing term and flat solutions, Advances in Nonlinear Analysis 13 (2024), no. 1, 20230128.
- [26] J. I. Díaz, J. Hernández, and Y. Ilyasov, Flat solutions of some non-Lipschitz autonomous semilinear equations may be stable for  $n \geq 3$ , Chinese Ann. Math 38 (2017), 345–378.
- [27] J. I. Díaz, J. Hernández, and F. J. Mancebo, Branches of positive and free boundary solutions for some singular quasilinear elliptic problems, Journal of Mathematical Analysis and Applications 352 (2009), no. 1, 449–474.
- [28] J. I. Díaz, J. Hernández, and J. M. Rakotoson, On very weak positive solutions to some semilinear elliptic problems with simultaneous singular nonlinear and spatial dependence terms, Milan Journal of Mathematics **79** (2011), 233–245.
- [29] J. I. Díaz, J.-M. Morel, and L. Oswald, An elliptic equation with singular nonlinearity, Communications in Partial Differential Equations 12 (1987), no. 12, 1333–1344.
- [30] J. I. Díaz and J. M. Rakotoson, On very weak solutions of semilinear elliptic equations with right-hand side data integrable with respect to the distance to the boundary, Discrete and Continuum Dynamical Systems 27 (2010), no. 3, 1037.
- [31] W. Fulks and J. S. Maybee, A singular non-linear equation, Osaka J. Math. 12 12 (1960), 1–19.
- [32] G. Gamow, Zur quantentheorie des atomkernes, Zeitschrift für Physik 51 (1928), no. 3, 204–212.
- [33] J. Hernández and F. J. Mancebo, Singular elliptic and parabolic equations, Handbook of differential equations: stationary partial differential equations, vol. 3, Elsevier, 2006, pp. 317–400.
- [34] I. Hofmann, Space charge physics for particle accelerators, Springer, 2017.
- [35] S. Humphries, Charged particle beams, Courier Corporation, 2013.
- [36] A. M. Il'in, A. S. Kalashnikov, and O. A. Oleinik, *Linear equations of the second order of parabolic type*, Russian Mathematical Surveys **17** (1962), no. 3, 1.
- [37] P. T. Kirstein, G. S. Kino, and W. E. Waters, Space-charge flow, McGraw Hill, 1967.
- [38] I. Langmuir, The effect of space charge and residual gases on thermionic currents in high vacuum, Physical Review 2 (1913), no. 6, 450.

- [39] J. W. Luginsland, Y.Y. Lau, R.J. Umstattd, and J.J. Watrous, Beyond the Child-Langmuir law: A review of recent results on multidimensional space-charge-limited flow, Physics of Plasmas 9 (2002), no. 5, 2371–2376.
- [40] M. Montenegro and A. Ponce, The sub-supersolution method for weak solutions, Proceedings of the American Mathematical Society 136 (2008), no. 7, 2429–2438.
- [41] C.-V. Pao, Nonlinear parabolic and elliptic equations, Springer Science & Business Media, 2012.
- [42] D. Philips, Existence of solutions of quenching problems, Applicable Analysis 24 (1987), no. 4, 253–264.
- [43] G. L. Pollack, D. R. Stump, and C. Inglefield, *Electromagnetism*, American Association of Physics Teachers, 2005.
- [44] A. Rokhlenko, Space charge limited flow in a rectangular region: profile of the current density, Journal of Applied Physics **100** (2006), no. 1 article no. 013305.
- [45] A. Rokhlenko and J. L. Lebowitz, Space-charge-limited 2d electron flow between two flat electrodes in a strong magnetic field, Physical Review Letters 91 (2003), no. 8, 085002.
- [46] \_\_\_\_\_, Space charge limited flow of a thin electron beam confined by a strong magnetic field, Physics of Plasmas 11 (2004), no. 10, 4559–4563.
- [47] Ph. Souplet, Optimal regularity conditions for elliptic problems via  $L^p_{\delta}$ -spaces, Duke Math. J. 127 (2005), no. 1, 175–192.
- [48] R. J. Umstattd and J. W. Luginsland, Two-dimensional space-charge-limited emission: beam-edge characteristics and applications, Physical Review Letters 87 (2001), no. 14, 145002.
- [49] L. Véron, Effets régularisants de semi-groupes non linéaires dans des espaces de Banach, Annales de la Faculté des Sciences de Toulouse: Mathématiques, vol. 1, 1979, pp. 171–200.
- [50] J. J. Watrous, J. W. Luginsland, and M. H. Frese, Current and current density of a finite-width, space-charge-limited electron beam in two-dimensional, parallel-plate geometry, Physics of Plasmas 8 (2001), no. 9, 4202–4210.
- [51] P. Zhang, A. Valfells, L. K. Ang, J. W. Luginsland, and Y. Y. Lau, 100 Years of the Physics of Diodes, Applied Physics Reviews 4 (2017), no. 1, 4.011304-1-4.011304-29.