



A note on the dynamics of an oscillator in the presence of strong friction

H. Amann^a, J.I. Diaz^{b,*}

^a*Institut für Mathematik, Universität Zürich, Winterthurerstr. 190, Zürich CH-8057, Switzerland*

^b*Departamento de Matemática Aplicada, Facultad de CC. Matemáticas,
Universidad Complutense de Madrid, Ciudad Universitaria, Madrid 28040, Spain*

Received 14 February 2003; accepted 14 February 2003

Abstract

We study the longtime behavior of the solutions of a second-order autonomous differential equation, differing from the one of a harmonic oscillator by a nonlinear friction term being only Hölder continuous. In particular, we show that there are two solution curves reaching the rest point in finite time.

© 2003 Elsevier Ltd. All rights reserved.

MSC: primary 34C05; 34D05; secondary 34C15; 34C35

Keywords: Phase plane analysis; Stability; Asymptotic behavior; Hölder continuous nonlinearity

1. Introduction

In this note we study the phase plane flow of the equation

$$\ddot{x} + \varphi(\dot{x}) + x = 0 \tag{1.1}$$

with the friction function φ possessing the following properties:

- it is continuous on \mathbb{R} and locally Lipschitz continuous on $\mathbb{R} \setminus \{0\}$,
- odd, and positive for positive arguments;
- there exist $a, \varepsilon > 0$ and $\alpha \in (0, 1)$ such that $\varphi(\eta) = a\eta^\alpha$ for $0 < \eta \leq \varepsilon$;
- $\lim_{\eta \rightarrow \infty} \varphi(\eta)/\eta < \frac{1}{2}$. (1.2)

* Corresponding author. Fax: +34-91-394-4407.

E-mail addresses: amann@math.unizh.ch (H. Amann), ji.diaz@mat.ucm.es (J.I. Diaz).

Clearly, if φ is defined by

$$\varphi(\eta) := \text{sign}(\eta) |\eta|^\alpha, \quad \eta \in \mathbb{R}, \tag{1.3}$$

then it satisfies (1.2) whenever $\alpha \in (0, 1)$, and the corresponding differential equation is

$$\ddot{x} + |\dot{x}|^{\alpha-1} \dot{x} + x = 0. \tag{1.4}$$

A nonmonotone example is given by

$$\varphi(\eta) := \begin{cases} a\eta^\alpha, & 0 \leq \eta \leq \varepsilon, \\ 1 + 1/\eta, & \varepsilon < \eta < \infty, \\ -\varphi(-\eta), & -\infty < \eta < 0, \end{cases} \tag{1.5}$$

where $a = \varepsilon^{-1-\alpha} + \varepsilon^{-\alpha}$. This nonlinearity appears in the literature because of the pioneering work of Lord Rayleigh (see [4,5]).

Functions satisfying (1.2) are often used to provide continuous approximations of the Coulomb friction problem

$$\ddot{x} + \text{sign}(\dot{x}) + x \ni 0, \tag{1.6}$$

where sign is meant to represent the maximal monotone graph in \mathbb{R}^2 , given by -1 for $\eta < 0$, by $[-1, 1]$ for $\eta = 0$, and by 1 for $\eta > 0$.

It is easy to see (e.g., [4]) that, in the Coulomb friction problem, every point in the phase plane reaches a critical point $(\xi, 0)$ with $\xi \in [-1, 1]$ in finite time by the flow induced by (1.6). In the case of Eq. (1.1) the origin is the only critical point of the phase flow, that is, the flow generated by

$$\dot{x} = y, \quad \dot{y} = -x - \varphi(y). \tag{1.7}$$

From (1.2) and (1.7) we deduce that

$$(x^2 + y^2)' = -2\varphi(y)y < 0, \quad y \neq 0.$$

Thus, given any $R > 0$, the disc $\bar{B}_R := \{(x, y) \in \mathbb{R}^2; x^2 + y^2 \leq R^2\}$ is positively invariant. Hence the origin is globally asymptotically stable, that is, it is stable and every point $(x, y) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ is transported to the origin by the flow. (Henceforth we simply say that ‘the orbit through (x, y) reaches the origin’, etc.)

Since, as mentioned above, Eq. (1.1) can be considered to be an approximation of the Coulomb friction problem, there arises the question whether the critical point is reached in finite time in this case as well. In [2,3] it is shown that for problem (1.4) there exist at least two orbits reaching the origin in finite time and there is an albeit small subset M of $\mathbb{R}^2 \setminus \{(0, 0)\}$ such that each orbit meeting M needs infinite time to reach the rest state. The proofs are based on asymptotic methods and comparison theorems and make use of the explicit form (1.3) of φ .

In this paper we show by completely different and elementary techniques from the theory of ordinary differential equations that, given assumptions (1.2), problem (1.7) also possesses at least two orbits converging to the rest state in finite time. Furthermore, we give a rather detailed analysis of the flow in the neighborhood of the critical point, showing that most orbits reach it in infinite time only. In fact, on the basis of our results we conjecture that there are precisely two orbits converging to zero in finite time.

2. The semiflow on the upper half-plane

First, we observe that the oddness of φ implies that the phase portrait of (1.7) is invariant under the reflection $(x, y) \mapsto (-x, -y)$. Thus it suffices to study the semiflow on the closure of the upper half-plane $\mathbb{H} := \{(x, y) \in \mathbb{R}^2; y > 0\}$, induced by restriction from the flow generated by (1.7). Consequently, in the rest of this section all assertions on invariant regions, etc. pertain to this semiflow. (We refer to [1] for the elementary facts about semiflows and invariant regions which we use without further mention).

From (1.7) we see that every orbit meets the real axis off zero vertically. For $0 < s \leq 1$ we denote by Γ_s the curve

$$\Gamma_s := \{(x, y) \in \mathbb{H}; -x = s\varphi(y)\},$$

oriented so that y is decreasing. Then every orbit in \mathbb{H} meets $\Gamma := \Gamma_1$ horizontally, and the direction field has a positive x -component and a negative y -component on the right of Γ . Thus no orbit approaches the origin through the first quadrant. Furthermore, given any $\eta > 0$,

$$\{(x, y) \in \mathbb{H}; -x \leq \varphi(y), y \leq \eta\}$$

is positively invariant.

We denote by $C(x, y)$ the solution curve of (1.7) passing through $(x, y) \in \mathbb{H}$ and being oriented in positive time direction.

Lemma 2.1. *There exists $\eta^* > 0$ such that $C(x, y)$ meets the positive y -axis whenever $(x, y) \in \Gamma_s$ for some $s \in (0, 1]$ and satisfies $y \geq \eta^*$.*

Proof. Suppose that $(x_0, y_0) \in \Gamma_s$ and $C(x_0, y_0)$ stays for positive time in

$$\mathbb{H}^- := \{(x, y) \in \mathbb{H}; x < 0\}.$$

From (1.2) we deduce the existence of $\alpha < \frac{1}{2}$ and $\bar{\eta} > 0$ such that

$$\varphi(\eta) \leq \alpha\eta, \quad \eta \geq \bar{\eta}. \tag{2.1}$$

Fix $\beta \in (\frac{1}{2}, 1 - \alpha)$ and suppose that

$$y_0 \geq \bar{\eta}/\beta =: \eta^*. \tag{2.2}$$

Let t^* be the time at which $C(x_0, y_0)$ reaches the η -level, that is, $y(t^*) = \beta y_0$. Then we deduce from the second equation in (1.7), (2.1), and (2.2) that

$$t^* = \int_{\beta y_0}^{y_0} \frac{dy}{x + \varphi(y)} \geq \int_{\beta y_0}^{y_0} \frac{dy}{\varphi(y)} \geq \frac{1 - \beta}{\alpha}, \tag{2.3}$$

owing to $x(t) < 0$ for $0 \leq t \leq t^*$. By integrating the first equation of (1.7) we obtain

$$x(t^*) - x_0 = \int_0^{t^*} y \, dt \geq \beta y_0 t^*.$$

Thus, using $x_0 = -s\varphi(y_0)$ and (2.1), we deduce from (2.3) that

$$x(t^*) \geq x_0 + \frac{\beta(1-\beta)}{\alpha} y_0 \geq \varphi(y_0) \left(\frac{\beta(1-\beta)}{\alpha^2} - s \right) > 0,$$

owing to $\alpha < \frac{1}{2} < \beta < 1 - \alpha$. But this contradicts the assumption that $C(x_0, y_0)$ stays in \mathbb{H}^- and thus proves the assertion. \square

For $0 < s < 1$ and $(x, y) \in \Gamma_s$ the positive unit tangent vectors to $C(x, y)$ and Γ_s at (x, y) are given by

$$(y, (1-s)\varphi(y))/r(s, y), \quad (s\varphi'(y), -1)/\rho(s, y), \tag{2.4}$$

respectively, where

$$r(s, y) := \sqrt{y^2 + (1-s)^2[\varphi(y)]^2}, \quad \rho(s, y) := \sqrt{1 + s^2[\varphi'(y)]^2},$$

provided φ is differentiable at y . In this case these vectors are equal iff

$$y = \psi(s, y)s\varphi'(y), \quad (1-s)\varphi(y) = \psi(s, y),$$

where $\psi(s, y) := r(s, y)/\rho(s, y)$, that is, iff

$$y = s(1-s)\varphi(y)\varphi'(y). \tag{2.5}$$

Note that (1.2) implies

$$\varphi(y) = ay^2, \quad \varphi'(y) = \alpha\varphi(y)/y, \quad 0 < y \leq \varepsilon. \tag{2.6}$$

Hence we deduce from (2.5), setting

$$\beta := \frac{1}{2}(1-\alpha), \quad \lambda(s) := \alpha a^2 s(1-s),$$

that

$$y = k_2(s) := [\lambda(s)]^\beta, \tag{2.7}$$

provided $0 < y \leq \varepsilon$. Furthermore, $(x, y) \in \Gamma_s$ implies in this case

$$x = k_1(s) := -s\varphi(k_2(s)) = -s\alpha[\lambda(s)]^{2\beta}. \tag{2.8}$$

Set

$$s_\varepsilon := \begin{cases} \frac{1}{2} - \frac{1}{2} \sqrt{1 - \frac{4}{\alpha a^2} \varepsilon^{2(1-\alpha)}} & \text{if } \varepsilon < \left(\frac{\alpha a^2}{4}\right)^{1/2(1-\alpha)}, \\ \frac{1}{2} & \text{otherwise.} \end{cases}$$

Define the oriented curve $K := K_\varepsilon$ in \mathbb{H}^- by means of the parametrization

$$(0, s_\varepsilon] \rightarrow \mathbb{H}^-, \quad s \mapsto k(s) := (k_1(s), k_2(s)).$$

Then, by the preceding considerations,

$$k(s) \in \Gamma_s, \quad 0 < s \leq s_\varepsilon,$$

and the positive unit tangent vectors to $C(x, y)$ and Γ_s at $(x, y) = k(s)$ coincide for $0 < s \leq s_e$.

Lemma 2.2. *The region M in \mathbb{H}^- , bounded by $\{(k_1(s_e), \eta); 0 < \eta \leq k_2(s_e)\}$ and K , is positively invariant.*

Proof. It is clear that the direction field of (1.7) is pointing inward along the vertical boundary of M .

From (2.6) and (2.8) we deduce that

$$\dot{k}_1(s) = -\varphi(k_2(s))[1 + \alpha s \dot{k}_2(s)/k_1(s)].$$

Note that (2.7) implies

$$\frac{\dot{k}_2(s)}{k_2(s)} = \beta \frac{\dot{\lambda}(s)}{\lambda(s)} = \beta \left(\frac{1}{s} - \frac{1}{1-s} \right).$$

Hence,

$$\dot{k}_1(s) = k_1(s) \left[\frac{1}{s} + \alpha \beta \frac{\dot{\lambda}(s)}{\lambda(s)} \right].$$

The positive normal to K at $k(s)$ has the direction of $(-\dot{k}_2(s), \dot{k}_1(s))$. Consequently, the inner product of the positive unit normal to K and the positive unit tangent vector to $C(x, y)$ at $(x, y) = k(s)$ is a positive multiple of

$$\begin{aligned} &(-\dot{k}_2(s), \dot{k}_1(s)) \cdot (k_2(s), -k_1(s) - \varphi(k_2(s))) \\ &= -[k_2(s)]^2 \beta \frac{\dot{\lambda}(s)}{\lambda(s)} + [\varphi(k_2(s))]^2 s(1-s) \left(\frac{1}{s} + \alpha \beta \frac{\dot{\lambda}(s)}{\lambda(s)} \right) \\ &= -\alpha a^2 \beta [\lambda(s)]^{2\beta-1} (1-2s) + a^2 \lambda^{2\alpha\beta} (1-s + \alpha\beta[1-s-s]) \\ &= a^2 [\lambda(s)]^{2\alpha\beta} (1-s) > 0. \end{aligned}$$

Thus the direction field of (1.7) is inward pointing along K as well. This implies the assertion. \square

Proposition 2.3. *There exists a positive semiorbit in \mathbb{H}^- reaching the origin in finite time.*

Proof. Let $s \in (0, s_e]$ be fixed and put

$$S := \{(x, y) \in \Gamma_s; C(x, y) \text{ meets the } y\text{-axis in } \mathbb{H}\}.$$

Then $S \neq \emptyset$ by Lemma 1.1. Put

$$\bar{y} := \inf \{y \in \mathbb{R}^+; (x, y) \in S\}.$$




Fig. 1.

If $(x, y) \in \Gamma_s$ and $y \leq k_2(s)$ then it follows from Lemma 2.2 that $(x, y) \notin S$. Hence $\bar{y} \geq k_2(s) > 0$.

For $(x, y) \in S$ we denote by $t(y)$ the time at which $C(x, y)$ meets the positive y -axis first, at $(0, y_0)$ say. Then, since $x(t) \geq -s\varphi(y(t))$ for $0 \leq t \leq t(y)$, it follows by integrating the first equation in (1.7) that

$$t(y) = \int_0^{t(y)} dt = \int_y^{y_0} \frac{dy}{-x - \varphi(y)} \leq \frac{1}{1-s} \int_0^y \frac{dy}{\varphi(y)}.$$

Thus fix $\hat{y} \in S$. Then

$$t(y) \leq \frac{1}{1-s} \int_0^{\hat{y}} \frac{dy}{\varphi(y)} =: T < \infty, \quad \bar{y} \leq y \leq \hat{y}.$$

Hence, we deduce from the continuous dependence of the flow on the initial value and the compactness of the interval $[0, T]$ that $\bar{y} \in S$. This proves the assertion since $t(\bar{y}) \leq T$. \square

In the following picture we depict \mathbb{H}^- in the neighborhood of the origin together with K , Γ , and Γ_s for three values of s , as well as seven orbits for problem (1.5) with $\alpha = 0.4$ and $\varepsilon = 1.2$. (See Fig. 1).

Finally, we prove that there are orbits needing infinite times to reach the origin.

Proposition 2.4. *Every orbit meeting M needs infinite time to reach the origin.*

Proof. Choose $s \in (0, s_c]$ and suppose that $(\xi, \eta) \in \Gamma_s \cap M$. From (2.4) we deduce that the inner product of the unit tangent vector to $C(\xi, \eta)$ with the negative unit

normal to Γ_s at (ξ, η) is a positive multiple of

$$\begin{aligned} \gamma(s, \eta) &:= s(1 - s)\varphi(\eta)\varphi'(\eta) - \eta = \eta[\alpha a^2 s(1 - s)\eta^{2(x-1)} - 1], \\ &= \eta[\lambda(s)\eta^{2(x-1)} - 1]. \end{aligned}$$

From $0 < \eta \leq k_2(s) = [\lambda(s)]^{1/2}$ it follows that $\gamma(s, \eta) \geq \gamma(s, k_2(s)) = 0$. This shows that, given any s , the subregion of M lying below Γ_s is positively invariant.

For $(x, y) \in M$ there is a unique $s \in (0, s_x]$ such that Γ_s meets K in (x, y_s) , where y_s satisfies $s\varphi(y_s) = -x$. Thus, the positive orbit starting at time zero in (x, y) lies below Γ_s , that is, $s\varphi(y(t)) \leq -x(t)$ for $0 \leq t < t^+$, where t^+ is the positive escape time of (x, y) . Consequently, by integrating the first equation of (1.7), we see that

$$t^+ = \int_0^{t^+} dt = \int_x^0 \frac{dx}{y} \geq \int_x^0 \frac{dx}{(-x/a)^{1/\alpha}} = \int_0^{-x/a} \frac{d\xi}{\xi^{1/\alpha}} = \infty,$$

owing to $1/\alpha > 1$. \square

3. The final result

After the preceding preparations we can now prove the main theorem of this paper. It gives a rather precise picture of the flow on \mathbb{R}^2 generated by (1.7).

Theorem 3.1. *The origin is the only critical point of the phase flow of (1.1), and it is globally asymptotically stable. More precisely, given any $R > 0$, the disc $\bar{\mathbb{B}}_R$ is positively invariant and every orbit converges to the rest state. The phase portrait is symmetric with respect to the origin.*

There exist real numbers satisfying $0 < \bar{x} \leq \hat{x}$ with the following properties:

- (i) *Every orbit meets $[-\hat{x}, \hat{x}] \times \{0\}$.*
- (ii) *If an orbit meets $[-\hat{x}, 0] \times \{0\}$, resp. $(0, \hat{x}] \times \{0\}$, then it stays in \mathbb{H}^- , resp.*

$$(-\mathbb{H})^+ := \{(x, y) \in \mathbb{R}^2; x > 0, y < 0\}.$$

If an orbit meets $(-\infty, -\hat{x}) \times \{0\}$, resp. $(\hat{x}, \infty) \times \{0\}$, then it leaves \mathbb{H}^- , resp. $(-\mathbb{H})^+$, after finite time.

- (iii) *The orbits through $(-\hat{x}, 0)$ and $(0, \hat{x})$ reach the rest point in finite time.*
- (iv) *Every orbit meeting $[(-\bar{x}, 0) \cup (0, \bar{x})] \times \{0\}$ needs infinite time to reach the critical point.*

Proof. The first part of the assertion has already been observed in the Introduction and at the beginning of Section 2.

By Proposition 2.3 and its proof there exists $(\hat{x}_0, \hat{y}_0) \in \mathbb{H}^-$ such that the positive semiorbit starting in (\hat{x}_0, \hat{y}_0) stays in \mathbb{H}^- , converges to $(0, 0)$ in finite time, and every positive semiorbit through (\hat{x}_0, η) with $\eta > \hat{y}_0$ leaves \mathbb{H}^- by intersecting the positive y -axis away from zero. Let $\gamma^-(\hat{x}_0, \hat{y}_0)$ be the negative semiorbit starting at time zero at (\hat{x}_0, \hat{y}_0) . Then it is easily seen that there exists $\hat{x} > 0$ such that $\gamma^-(\hat{x}_0, \hat{y}_0)$ meets $(-\infty, 0) \times \{0\}$ for the first time in $(-\hat{x}, 0)$. Thus assertions (i)–(iii) are clear.




Fig. 2.

Assertion (iv) is now an easy consequence of what has just been shown and of Proposition 2.4. \square

In the next picture a portion of the phase plane of problem (1.1) is depicted, where φ is given by (1.5) with $\alpha = 0.4$ and $\varepsilon = 1.2$. (See Fig. 2).

We conjecture, but do not know, that $\bar{x} = \hat{x}$ in which case there are precisely two orbits reaching the origin in finite time.

References

- [1] H. Amann, *Ordinary Differential Equations*, de Gruyter Studies in Mathematics, Vol. 13, Walter de Gruyter & Co., Berlin, 1990. (An introduction to nonlinear analysis, Translated from German by G. Metzen).
- [2] J.I. Díaz, A. Linán, On the asymptotic behavior for a damped oscillator under a sublinear friction, *RACSAM Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.* 95 (1) (2001) 155–160.
- [3] J.I. Díaz, A. Linán, On the asymptotic behavior of solutions of a damped oscillator under a sublinear friction term: from the exceptional to the generic behaviors, in: *Partial Differential Equations, Lecture Notes in Pure and Applied Mathematics*, Vol. 229, Marcel Dekker, New York, 2002, pp. 163–170.
- [4] D.W. Jordan, P. Smith, *Nonlinear Ordinary Differential Equations*, 2nd Edition, Oxford Applied Mathematics and Computing Science Series, The Clarendon Press, Oxford University Press, New York, 1987.
- [5] J.W. Rayleigh, B. Strutt, *The Theory of Sound*, 2nd Edition, Dover Publications, New York, NY, 1945.